

AN EXAMPLE ARTICLE*

DIANNE DOE[†], PAUL T. FRANK[‡], AND JANE E. SMITH[‡]

Abstract. This is an example SIAM L^AT_EX article. This can be used as a template for new articles. Abstracts must be able to stand alone and so cannot contain citations to the paper's references, equations, etc. An abstract must consist of a single paragraph and be concise. Because of online formatting, abstracts must appear as plain as possible. Any equations should be inline.

Key words. example, L^AT_EX

MSC codes. 68Q25, 68R10, 68U05

1. Introduction. The introduction introduces the context and summarizes the manuscript. It is important to clearly state the contributions of this piece of work.

For $\Omega = (0, 2T)$, $1 < \alpha < 2$, suppose $f \in C^2(\Omega)$

$$(1.1) \quad \begin{cases} (-\Delta)^{\frac{\alpha}{2}} u(x) = f(x), & x \in \Omega \\ u(x) = 0, & x \in \mathbb{R} \setminus \Omega \end{cases}$$

where

$$(1.2) \quad (-\Delta)^{\frac{\alpha}{2}} u(x) = -\frac{\partial^\alpha u}{\partial |x|^\alpha} = -\kappa_\alpha \frac{d^2}{dx^2} \int_\Omega \frac{|x-y|^{1-\alpha}}{\Gamma(2-\alpha)} u(y) dy$$

$$(1.3) \quad \kappa_\alpha = -\frac{1}{2 \cos(\alpha\pi/2)} > 0$$

THEOREM 1.1. *Let u be a solution of (1.1) on Ω . Then, for any $x \in \Omega$ and $l = 0, 1, 2, 3, 4$*

$$(1.4) \quad |u^{(l)}(x)| \leq C[x(2T-x)]^{\alpha/2-l}$$

The paper is organized as follows. Our main results are in section 3, experimental results are in section 6, and the conclusions follow in section 8.

2. Numeric Format.

$$(2.1) \quad x_i = \begin{cases} T \left(\frac{i}{N} \right)^r, & 0 \leq i \leq N \\ 2T - T \left(\frac{2N-i}{N} \right)^r, & N \leq i \leq 2N \end{cases}$$

where $r \geq 1$. And let

$$(2.2) \quad h_j = x_j - x_{j-1}, \quad 1 \leq j \leq 2N$$

*Submitted to the editors DATE.

Funding: This work was funded by the Fog Research Institute under contract no. FRI-454.

[†]Imagination Corp., Chicago, IL (ddoe@imag.com, <http://www.imag.com/~ddoe/>).

[‡]Department of Applied Mathematics, Fictional University, Boise, ID (ptfrank@fictional.edu, jesmith@fictional.edu).

Let $\{\phi_j(x)\}_{j=1}^{2N-1}$ be standard hat functions, which are basis of the piecewise linear function space.

$$\phi_j(x) = \begin{cases} \frac{1}{h_j}(x - x_{j-1}), & x_{j-1} \leq x \leq x_j \\ \frac{1}{h_{j+1}}(x_{j+1} - x), & x_j \leq x \leq x_{j+1} \\ 0, & \text{otherwise} \end{cases}$$

And then, we can approximate $u(x)$ with

$$u_h(x) := \sum_{j=1}^{2N-1} u(x_j)\phi_j(x)$$

For convience, we denote

$$I_h^{2-\alpha}(x) := \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x-y|^{1-\alpha} u_h(y) dy$$

And now, we can approximate the operator (1.2) at x_i with

$$\begin{aligned} D_h^\alpha u_h(x_i) &:= D_h^2 I_h^{2-\alpha}(x_i) \\ &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} I_h^{2-\alpha}(x_{i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) I_h^{2-\alpha}(x_i) + \frac{1}{h_{i+1}} I_h^{2-\alpha}(x_{i+1}) \right) \end{aligned}$$

Finally, we approximate the equation (1.1) with

$$-\kappa_\alpha D_h^\alpha u_h(x_i) = f(x_i), \quad 1 \leq i \leq 2N-1$$

The discrete equation (2.7) can be written in matrix form

$$AU = F$$

where U is unknown, $F = (f(x_1), \dots, f(x_{2N-1}))$. The matrix A is constructed as follows: Since

$$\begin{aligned} I_h^{2-\alpha}(x_i) &= \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u_h(y) dy \\ &= \sum_{j=1}^{2N-1} \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u(x_j) \phi_j(y) dy \\ &= \sum_{j=1}^{2N-1} u(x_j) \frac{1}{\Gamma(2-\alpha)} \int_{x_{j-1}}^{x_{j+1}} |x_i - y|^{1-\alpha} \phi_j(y) dy \\ &= \sum_{j=1}^{2N-1} \frac{u(x_j)}{\Gamma(4-\alpha)} \left(\frac{|x_i - x_{j-1}|^{3-\alpha}}{h_j} - \frac{h_j + h_{j+1}}{h_j h_{j+1}} |x_i - x_j|^{3-\alpha} + \frac{|x_i - x_{j+1}|^{3-\alpha}}{h_{j+1}} \right) \\ &=: \sum_{j=1}^{2N-1} \tilde{a}_{ij} u(x_j), \quad 0 \leq i \leq 2N \end{aligned}$$

Then, substitute in (2.6), we have

$$(2.10) \quad -\kappa_\alpha D_h^\alpha u_h(x_i) = \sum_{j=1}^{2N-1} a_{ij} u(x_j)$$

where

$$(2.11) \quad a_{ij} = -\kappa_\alpha \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \tilde{a}_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{a}_{i,j} + \frac{1}{h_{i+1}} \tilde{a}_{i+1,j} \right)$$

3. Main results. Here we state our main results; the proof is deferred to section 4 and section 5.

Let's denote $h = \frac{1}{N}$, we have

THEOREM 3.1 (Truncation Error). *If $f \in C^2(\Omega)$ and $\alpha \in (1, 2)$, and $u(x)$ is a solution of the equation (1.1), then there exists a constant $C = C(T, \alpha, r, \|f\|_{C^2(\Omega)})$, such that the truncation error of the discrete format satisfies*

$$(3.1) \quad \begin{aligned} |-\kappa_\alpha D_h^\alpha u_h(x_i) - f(x_i)| \leq & C(h^{r\alpha/2+r}(x_i^{-1-\alpha} + (2T - x_i)^{-1-\alpha}) \\ & + h^2(x_i^{-\alpha/2-2/r} + (2T - x_i)^{-\alpha/2-2/r}) \\ & + h^2 \begin{cases} |T - x_{i-1}|^{1-\alpha}, & 1 \leq i \leq N \\ |T - x_{i+1}|^{1-\alpha}, & N < i \leq 2N-1 \end{cases} \end{aligned}$$

THEOREM 3.2 (Convergence). *The discrete equation (2.7) has solution U , and there exists a positive constant $C = C(T, \alpha, r, \|f\|_{C^2(\Omega)})$ such that the error between the numerical solution U with the exact solution $u(x_i)$ satisfies*

$$(3.2) \quad \max_{1 \leq i \leq 2N-1} |U_i - u(x_i)| \leq Ch^{\min\{\frac{r\alpha}{2}, 2\}}$$

That means the numerical method has convergence order $\min\{\frac{r\alpha}{2}, 2\}$.

4. Proof of Theorem 3.1. For convenience, let's denote

$$(4.1) \quad I^{2-\alpha}(x) = \frac{1}{\Gamma(2-\alpha)} \int_\Omega |x-y|^{1-\alpha} u(y) dy$$

Then, the truncation error of the discrete format can be written as

$$(4.2) \quad \begin{aligned} -\kappa_\alpha D_h^\alpha u_h(x_i) - f(x_i) &= -\kappa_\alpha (D_h^2 I_h^{2-\alpha}(x_i) - \frac{d^2}{dx^2} I^{2-\alpha}(x_i)) \\ &= -\kappa_\alpha D_h^2 (I_h^{2-\alpha}(x_i) - I^{2-\alpha}(x_i)) - \kappa_\alpha (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) \end{aligned}$$

THEOREM 4.1. *There exists a constant $C = C(T, r, \|f\|_{C^2(\Omega)})$ such that*

$$(4.3) \quad -\kappa_\alpha (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) \leq Ch^2(x_i^{1-2/r} + (2T - x_i)^{1-2/r})$$

Proof. Since $f \in C^2(\Omega)$ and

$$(4.4) \quad \frac{d^2}{dx^2}(-\kappa_\alpha I^{2-\alpha}(x)) = f(x), \quad x \in \Omega,$$

we have $I^{2-\alpha} \in C^4(\Omega)$. Therefore, using equation (A.2) of Lemma A.1, for $1 \leq i \leq 2N-1$, we have

$$(4.5) \quad -\kappa_\alpha(D_h^2 - \frac{d^2}{dx^2})I^{2-\alpha}(x_i) = \frac{h_{i+1} - h_i}{3}f'(x_i) + \frac{1}{4!} \frac{2}{h_i + h_{i+1}}(h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2))$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$. By Lemma B.2, we have 1.

$$(4.6) \quad \left| \frac{h_{i+1} - h_i}{3}f'(x_i) \right| \leq \frac{\|f\|_{C^1(\Omega)}}{3} 2^{|r-2|} r(r-1) T^{2/r} h^2 (x_i^{1-2/r} + (2T - x_i)^{1-2/r})$$

2.

$$(4.7) \quad \begin{aligned} & \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \\ & \leq \frac{\|f\|_{C^2(\Omega)}}{12} (h_i^2 - h_i h_{i+1} + h_{i+1}^2) \end{aligned} \quad \square$$

5. Proof of Theorem 3.2. aaaaaaaaaa

6. Experimental results. Figure 1 shows some example results. Additional results are available in the supplement in Table 1. Table 1 shows additional supporting




FIG. 1. Example figure using external image files.

evidence.

TABLE 1
Example table.

Species	Mean	Std. Dev.
1	3.4	1.2
2	5.4	0.6
3	7.4	2.4
4	9.4	1.8

7. Discussion of $Z = X \cup Y$.

8. Conclusions. Some conclusions here.

Appendix A. Approximate of difference quotients.

LEMMA A.1. *If $g(x)$ is twice differentiable continuous function on open set Ω , there exists $\xi \in [x_{i-1}, x_{i+1}]$ such that*

$$(A.1) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) = g''(\xi), \quad \xi \in [x_{i-1}, x_{i+1}]$$

And if $g(x) \in C^4(\Omega)$, then

$$(A.2) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) = g''(x_i) + \frac{h_{i+1} - h_i}{3} g'''(x_i) + \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 g''''(\eta_1) + h_{i+1}^3 g''''(\eta_2))$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$.

Proof.

$$g(x_{i-1}) = g(x_i) - (x_i - x_{i-1})g'(x_i) + \frac{(x_i - x_{i-1})^2}{2} g''(\xi_1), \quad \xi_1 \in [x_{i-1}, x_i]$$

$$g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i)g'(x_i) + \frac{(x_{i+1} - x_i)^2}{2} g''(\xi_2), \quad \xi_2 \in [x_i, x_{i+1}]$$

Substitute them in the left side of (A.1), we have

$$\frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) = \frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2)$$

Now, using intermediate value theorem, there exists $\xi \in [\xi_1, \xi_2]$ such that

$$\frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2) = g''(\xi)$$

And for the second equation, similarly

$$g(x_{i-1}) = g(x_i) - h_i g'(x_i) + \frac{h_i^2}{2} g''(x_i) - \frac{h_i^3}{3!} g'''(x_i) + \frac{h_i^4}{4!} g''''(\eta_1)$$

$$g(x_{i+1}) = g(x_i) + h_{i+1} g'(x_i) + \frac{h_{i+1}^2}{2} g''(x_i) + \frac{h_{i+1}^3}{3!} g'''(x_i) + \frac{h_{i+1}^4}{4!} g''''(\eta_2)$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$. Substitute them to the left side of (A.2), we can get the result. \square

LEMMA A.2. *If $y \in [x_{j-1}, x_j]$, denote $y = \theta x_{j-1} + (1 - \theta)x_j$, $\theta \in [0, 1]$,*

$$(A.3) \quad u(y_j^\theta) - u_h(y_j^\theta) = -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [x_{j-1}, x_j]$$

(A.4)

$$u(y_j^\theta) - u_h(y_j^\theta) = -\frac{\theta(1-\theta)}{2}h_j^2u''(y_j^\theta) + \frac{\theta(1-\theta)}{3!}h_j^3(\theta^2u'''(\eta_1) - (1-\theta)^2u'''(\eta_2))$$

where $\eta_1 \in [x_{j-1}, y_j^\theta], \eta_2 \in [y_j^\theta, x_j]$.

Proof. By Taylor expansion, we have

$$u(x_{j-1}) = u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(\xi_1), \quad \xi_1 \in [x_{j-1}, y_j^\theta]$$

$$u(x_j) = u(y_j^\theta) + (1-\theta)h_j u'(y_j^\theta) + \frac{(1-\theta)^2 h_j^2}{2!} u''(\xi_2), \quad \xi_2 \in [y_j^\theta, x_j]$$

Thus

$$\begin{aligned} u(y_j^\theta) - u_h(y_j^\theta) &= u(y_j^\theta) - (1-\theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1-\theta)}{2}h_j^2(\theta u''(\xi_1) + (1-\theta)u''(\xi_2)) \\ &= -\frac{\theta(1-\theta)}{2}h_j^2u''(\xi), \quad \xi \in [\xi_1, \xi_2] \end{aligned}$$

The second equation is similar,

$$\begin{aligned} u(x_{j-1}) &= u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(y_j^\theta) - \frac{\theta^3 h_j^3}{3!} u'''(\eta_1) \\ u(x_j) &= u(y_j^\theta) + (1-\theta)h_j u'(y_j^\theta) + \frac{(1-\theta)^2 h_j^2}{2!} u''(\xi_2) + \frac{(1-\theta)^3 h_j^3}{3!} u'''(\eta_2) \end{aligned}$$

where $\eta_1 \in [x_{j-1}, y_j^\theta], \eta_2 \in [y_j^\theta, x_j]$. Thus

$$\begin{aligned} u(y_j^\theta) - u_h(y_j^\theta) &= u(y_j^\theta) - (1-\theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1-\theta)}{2}h_j^2u''(y_j^\theta) + \frac{\theta(1-\theta)}{3!}h_j^3(\theta^2u'''(\eta_1) - (1-\theta)^2u'''(\eta_2)) \end{aligned}$$

Appendix B. Inequality.

LEMMA B.1.

$$(B.1) \quad h_i \leq rT^{1/r}h \begin{cases} x_i^{1-1/r}, & 1 \leq i \leq N \\ (2T - x_{i-1})^{1-1/r}, & N < i \leq 2N-1 \end{cases}$$

Proof. For $1 \leq i \leq N$,

$$\begin{aligned} h_i &= T \left(\left(\frac{i}{N} \right)^r - \left(\frac{i-1}{N} \right)^r \right) \\ &\leq rT \frac{1}{N} \left(\frac{i}{N} \right)^{r-1} = rT^{1/r}h x_i^{1-1/r} \end{aligned}$$

For $N < i \leq 2N-1$,

$$\begin{aligned} h_i &= T \left(\left(\frac{2N-i}{N} \right)^r - \left(\frac{2N-i+1}{N} \right)^r \right) \\ &\leq rT \frac{1}{N} \left(\frac{2N-i+1}{N} \right)^{r-1} = rT^{1/r}h (2T - x_{i-1})^{1-1/r} \end{aligned}$$

LEMMA B.2. *There is a constant $C = 2^{|r-2|}r(r-1)T^{2/r}$ such that for all $i \in \{1, 2, \dots, 2N-1\}$*

$$(B.2) \quad |h_{i+1} - h_i| \leq Ch^2(x_i^{1-2/r} + (2T - x_i)^{1-2/r})$$

Proof.

$$h_{i+1} - h_i = \begin{cases} T \left(\left(\frac{i+1}{N} \right)^r - 2 \left(\frac{i}{N} \right)^r + \left(\frac{i-1}{N} \right)^r \right), & 1 \leq i \leq N-1 \\ 0, & i = N \\ -T \left(\left(\frac{2N-i-1}{N} \right)^r - 2 \left(\frac{2N-i}{N} \right)^r + \left(\frac{2N-i+1}{N} \right)^r \right), & N+1 \leq i \leq 2N-1 \end{cases}$$

For $i = 1$,

$$h_2 - h_1 = T(2^r - 2) \left(\frac{1}{N} \right)^r = (2^r - 2)T^{2/r}h^2x_1^{1-2/r}$$

For $2 \leq i \leq N-1$,

$$h_{i+1} - h_i = r(r-1)T N^{-2}\eta^{r-2}, \quad \eta \in \left[\frac{i-1}{N}, \frac{i+1}{N} \right]$$

If $r \in [1, 2)$,

$$\begin{aligned} h_{i+1} - h_i &\leq r(r-1)T N^{-2}\eta^{r-2} \leq r(r-1)T h^2 \left(\frac{i-1}{N} \right)^{r-2} \\ &\leq r(r-1)T h^2 2^{2-r} \left(\frac{i}{N} \right)^{r-2} \\ &= 2^{2-r}r(r-1)T^{2/r}h^2x_i^{1-2/r} \end{aligned}$$

else if $r > 2$,

$$\begin{aligned} h_{i+1} - h_i &\leq r(r-1)T N^{-2}\eta^{r-2} \leq r(r-1)T h^2 \left(\frac{i+1}{N} \right)^{r-2} \\ &\leq r(r-1)T h^2 2^{r-2} \left(\frac{i}{N} \right)^{r-2} \\ &= 2^{r-2}r(r-1)T^{2/r}h^2x_i^{1-2/r} \end{aligned}$$

Since

$$2^r - 2 \leq 2^{|r-2|}r(r-1), \quad r \geq 1$$

we have

$$h_{i+1} - h_i \leq 2^{|r-2|}r(r-1)T^{2/r}h^2x_i^{1-2/r}, \quad 1 \leq i \leq N-1$$

For $i = N$, $h_{N+1} - h_N = 0$. For $N < i \leq 2N-1$, it's central symmetric to the first half of the proof, which is

$$h_i - h_{i+1} \leq 2^{|r-2|}r(r-1)T^{2/r}h^2(2T - x_i)^{1-2/r}$$

148 Summarizes the inequalities, we can get

149 (B.3) $|h_{i+1} - h_i| \leq 2^{|r-2|} r(r-1) T^{2/r} h^2(x_i^{1-2/r} + (2T - x_i)^{1-2/r}) \quad \square$

150 **Acknowledgments.** We would like to acknowledge the assistance of volunteers
 151 in putting together this example manuscript and supplement.

152

REFERENCES