

AN EXAMPLE ARTICLE*

DIANNE DOE[†], PAUL T. FRANK[‡], AND JANE E. SMITH[‡]

Abstract. This is an example SIAM L^AT_EX article. This can be used as a template for new articles. Abstracts must be able to stand alone and so cannot contain citations to the paper's references, equations, etc. An abstract must consist of a single paragraph and be concise. Because of online formatting, abstracts must appear as plain as possible. Any equations should be inline.

Key words. example, L^AT_EX

MSC codes. 68Q25, 68R10, 68U05

1. Introduction. The introduction introduces the context and summarizes the manuscript. It is importantly to clearly state the contributions of this piece of work.

For $\Omega = (0, 2T)$, $1 < \alpha < 2$, suppose $f \in C^\beta(\Omega)$, $\beta > 4 - \alpha$, $\|f\|_\beta^{(\alpha/2)} < \infty$

$$(1.1) \quad \begin{cases} (-\Delta)^{\frac{\alpha}{2}} u(x) = f(x), & x \in \Omega \\ u(x) = 0, & x \in \mathbb{R} \setminus \Omega \end{cases}$$

where

$$(1.2) \quad (-\Delta)^{\frac{\alpha}{2}} u(x) = -\frac{\partial^\alpha u}{\partial |x|^\alpha} = -\kappa_\alpha \frac{d^2}{dx^2} \int_\Omega \frac{|x-y|^{1-\alpha}}{\Gamma(2-\alpha)} u(y) dy$$

$$(1.3) \quad \kappa_\alpha = -\frac{1}{2 \cos(\alpha\pi/2)} > 0$$

and the solution $u \in C^{\alpha/2}(\Omega)$.

2. Regularity.

Remark 2.1. 1. $C^k(U)$ is the set of all k -times continuously differentiable functions on open set U .

2. $C^\beta(U)$ is the collection of function f which for any $V \subset\subset U$ $f|_V \in C^\beta(\bar{V})$.

THEOREM 2.2. If $f \in C^\beta(\Omega)$, $\beta > 2$ and $\|f\|_\beta^{(\alpha/2)} < \infty$, then for $l = 0, 1, 2$

$$(2.1) \quad |f^{(l)}(x)| \leq \|f\|_\beta^{(\alpha/2)} \begin{cases} x^{-l-\alpha/2}, & \text{if } 0 < x \leq T \\ (2T-x)^{-l-\alpha/2}, & \text{if } T \leq x < 2T \end{cases}$$

THEOREM 2.3 (Regularity up to the boundary [1]).

$$(2.2) \quad \|u\|_{\beta+\alpha}^{(-\alpha/2)} \leq C \left(\|u\|_{C^{\alpha/2}(\mathbb{R})} + \|f\|_\beta^{(\alpha/2)} \right)$$

*Submitted to the editors DATE.

Funding: This work was funded by the Fog Research Institute under contract no. FRI-454.

[†]Imagination Corp., Chicago, IL (ddoe@imag.com, <http://www.imag.com/~ddoe/>).

[‡]Department of Applied Mathematics, Fictional University, Boise, ID (ptfrank@fictional.edu, jesmith@fictional.edu).

29 **COROLLARY 2.4.** *Let u be a solution of (1.1) on Ω . Then, for any $x \in \Omega$ and*
 30 *$l = 0, 1, 2, 3, 4$*

$$31 \quad (2.3) \quad |u^{(l)}(x)| \leq C \begin{cases} x^{\alpha/2-l}, & \text{if } 0 < x \leq T \\ (2T-x)^{\alpha/2-l}, & \text{if } T \leq x < 2T \end{cases}$$

32 The paper is organized as follows. Our main results are in section 4, experimental
 33 results are in section 7, and the conclusions follow in section 8.

3. Numeric Format.

$$34 \quad (3.1) \quad x_i = \begin{cases} T \left(\frac{i}{N} \right)^r, & 0 \leq i \leq N \\ 2T - T \left(\frac{2N-i}{N} \right)^r, & N \leq i \leq 2N \end{cases}$$

35 where $r \geq 1$. And let

$$36 \quad (3.2) \quad h_j = x_j - x_{j-1}, \quad 1 \leq j \leq 2N$$

37 Let $\{\phi_j(x)\}_{j=1}^{2N-1}$ be standard hat functions, which are basis of the piecewise linear
 38 function space.

$$39 \quad (3.3) \quad \phi_j(x) = \begin{cases} \frac{1}{h_j}(x - x_{j-1}), & x_{j-1} \leq x \leq x_j \\ \frac{1}{h_{j+1}}(x_{j+1} - x), & x_j \leq x \leq x_{j+1} \\ 0, & \text{otherwise} \end{cases}$$

40 And then, we can approximate $u(x)$ with

$$41 \quad (3.4) \quad u_h(x) := \sum_{j=1}^{2N-1} u(x_j) \phi_j(x)$$

42 For convience, we denote

$$43 \quad (3.5) \quad I_h^{2-\alpha}(x_i) := \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u_h(y) dy$$

44 And now, we can approximate the operator (1.2) at x_i with

$$45 \quad (3.6) \quad \begin{aligned} D_h^\alpha u_h(x_i) &:= D_h^2 I_h^{2-\alpha}(x_i) \\ &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} I_h^{2-\alpha}(x_{i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) I_h^{2-\alpha}(x_i) + \frac{1}{h_{i+1}} I_h^{2-\alpha}(x_{i+1}) \right) \end{aligned}$$

46 Finally, we approximate the equation (1.1) with

$$47 \quad (3.7) \quad -\kappa_\alpha D_h^\alpha u_h(x_i) = f(x_i), \quad 1 \leq i \leq 2N-1$$

48 The discrete equation (3.7) can be written in matrix form

$$49 \quad (3.8) \quad AU = F$$

where U is unknown, $F = (f(x_1), \dots, f(x_{2N-1}))$. The matrix A is constructed as follows: Since

(3.9)

$$\begin{aligned}
 I_h^{2-\alpha}(x_i) &= \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u_h(y) dy \\
 &= \sum_{j=1}^{2N-1} \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u(x_j) \phi_j(y) dy \\
 &= \sum_{j=1}^{2N-1} u(x_j) \frac{1}{\Gamma(2-\alpha)} \int_{x_{j-1}}^{x_{j+1}} |x_i - y|^{1-\alpha} \phi_j(y) dy \\
 &= \sum_{j=1}^{2N-1} \frac{u(x_j)}{\Gamma(4-\alpha)} \left(\frac{|x_i - x_{j-1}|^{3-\alpha}}{h_j} - \frac{h_j + h_{j+1}}{h_j h_{j+1}} |x_i - x_j|^{3-\alpha} + \frac{|x_i - x_{j+1}|^{3-\alpha}}{h_{j+1}} \right) \\
 &=: \sum_{j=1}^{2N-1} \tilde{a}_{ij} u(x_j), \quad 0 \leq i \leq 2N
 \end{aligned}$$

Then, substitute in (3.6), we have

$$(3.10) \quad -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) = \sum_{j=1}^{2N-1} a_{ij} u(x_j)$$

where

$$(3.11) \quad a_{ij} = -\kappa_{\alpha} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \tilde{a}_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{a}_{i,j} + \frac{1}{h_{i+1}} \tilde{a}_{i+1,j} \right)$$

4. Main results. Here we state our main results; the proof is deferred to section 5 and section 6.

Let's denote $h = \frac{1}{N}$, we have

THEOREM 4.1 (Truncation Error). *If $f \in C^2(\Omega)$ and $\alpha \in (1, 2)$, and $u(x)$ is a solution of the equation (1.1), then there exists a constant $C_1, C_2 = C_1(T, \alpha, r, \|f\|_{C^2(\Omega)}), C_2(T, \alpha, r, \|f\|_{C^2(\Omega)})$, such that the truncation error of the discrete format satisfies*

$$\begin{aligned}
 |-\kappa_{\alpha} D_h^{\alpha} u_h(x_i) - f(x_i)| &\leq C_1 (h^{r\alpha/2+r} (x_i^{-1-\alpha} + (2T - x_i)^{-1-\alpha}) \\
 &\quad + h^2 (x_i^{-\alpha/2-2/r} + (2T - x_i)^{-\alpha/2-2/r})) \\
 &\quad + C_2 h^2 \begin{cases} |T - x_{i-1}|^{1-\alpha}, & 1 \leq i \leq N \\ |T - x_{i+1}|^{1-\alpha}, & N < i \leq 2N - 1 \end{cases}
 \end{aligned}$$

where $C_2 = 0$ if $r = 1$.

THEOREM 4.2 (Convergence). *The discrete equation (3.7) has solution U , and there exists a positive constant $C = C(T, \alpha, r, \|f\|_{C^2(\Omega)})$ such that the error between the numerical solution U with the exact solution $u(x_i)$ satisfies*

$$(4.2) \quad \max_{1 \leq i \leq 2N-1} |U_i - u(x_i)| \leq C h^{\min\{\frac{r\alpha}{2}, 2\}}$$

That means the numerical method has convergence order $\min\{\frac{r\alpha}{2}, 2\}$.

5. Proof of Theorem 4.1. For convience, let's denote

$$(5.1) \quad I^{2-\alpha}(x) = \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x-y|^{1-\alpha} u(y) dy$$

Then, the truncation error of the discrete format can be written as

$$(5.2) \quad \begin{aligned} -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) - f(x_i) &= -\kappa_{\alpha} (D_h^2 I_h^{2-\alpha}(x_i) - \frac{d^2}{dx^2} I^{2-\alpha}(x_i)) \\ &= -\kappa_{\alpha} D_h^2 (I_h^{2-\alpha} - I^{2-\alpha})(x_i) - \kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) \end{aligned}$$

5.1. Estimate of $-\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i)$.

THEOREM 5.1. *There exists a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$ such that*

$$(5.3) \quad \left| -\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) \right| \leq Ch^2 (x_i^{-\alpha/2-2/r} + (2T-x_i)^{-\alpha/2-2/r})$$

Proof. Since $f \in C^2(\Omega)$ and

$$(5.4) \quad \frac{d^2}{dx^2} (-\kappa_{\alpha} I^{2-\alpha}(x)) = f(x), \quad x \in \Omega,$$

we have $I^{2-\alpha} \in C^4(\Omega)$. Therefore, using equation (A.3) of Lemma A.1, for $1 \leq i \leq 2N-1$, we have

$$(5.5) \quad -\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) = \frac{h_{i+1} - h_i}{3} f'(x_i) + \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2))$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$. By Lemma B.2 and Theorem 2.2 we have 1.

$$(5.6) \quad \left| \frac{h_{i+1} - h_i}{3} f'(x_i) \right| \leq \frac{\|f\|_{\beta}^{(\alpha/2)}}{3} Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T-x_i)^{-\alpha/2-2/r}, & N < i \leq 2N-1 \end{cases}$$

2. See Proof 9, there is a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$ such that

$$(5.7) \quad \begin{aligned} &\left| \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \right| \\ &\leq Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T-x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N-1 \end{cases} \end{aligned}$$

Summarizes, we get the result. \square

5.2. Estimate of R_i . Now, we study the first part of (5.2)

$$(5.8) \quad D_h^2 (I^{2-\alpha} - I_h^{2-\alpha})(x_i) = D_h^2 \left(\int_0^{2T} (u(y) - u_h(y)) \frac{|y-x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \right)$$

For convience, let's denote

$$(5.9) \quad T_{ij} = \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) \frac{|y-x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy$$

92 And define

$$93 \quad (5.10) \quad R_i := D_h^2(I^{2-\alpha} - I_h^{2-\alpha})(x_i) \\ = \frac{2}{h_i + h_{i+1}} \sum_{j=1}^{2N} \left(\frac{1}{h_i} T_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_{i+1}} T_{i+1,j} \right)$$

94 We have some results about the estimate of R_i

95 **THEOREM 5.2.** *For $1 \leq i < N/2$, there exists a constant $C = C(T, \alpha, r, f)$ such*
 96 *that*

$$97 \quad (5.11) \quad R_i \leq C(h^{r\alpha/2+r} x_i^{-1-\alpha} + h^2 x_i^{-\alpha/2-2/r})$$

98

99 **THEOREM 5.3.** *For $N/2 \leq i \leq N$, there exists constant C, C_2 such that*

$$100 \quad (5.12) \quad R_i \leq C h^2 x_i^{-\alpha/2-2/r} + C_2 h^2 |T - x_{i-1}|^{1-\alpha}$$

101 *where $C_2 = 0$ if $r = 1$.*

102 And for $N < i \leq 2N - 1$, it is symmetric to the previous case.

103 To prove these results, we need some utils. Also for simplicity, we denote

$$104 \quad (5.13) \quad S_{ij} = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} T_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_{i+1}} T_{i+1,j} \right)$$

105 then

$$106 \quad (5.14) \quad R_i = \sum_{j=1}^{2N} S_{ij}$$

107 **5.3. Proof of Theorem 5.2.**

108 **LEMMA 5.4.** *There exists a constant $C = C(T, \alpha, r, f)$ such that for $1 \leq i < N/2$,*

$$109 \quad (5.15) \quad \sum_{j=\max\{2i+1, i+3\}}^{2N} S_{ij} \leq C h^2 x_i^{-\alpha/2-2/r}$$

110 *Proof.* For $\max\{2i+1, i+3\} \leq j \leq N$, by Lemma C.1 and Lemma C.2

$$111 \quad (5.16) \quad S_{ij} = \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) D_h^2 \left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)} \right) (x_i) dy \\ \leq C h^2 \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} dy \\ = C h^2 \int_{x_{j-1}}^{x_j} y^{-\alpha/2-2/r-1} dy$$

112 Therefore,

$$113 \quad (5.17) \quad \sum_{j=\max\{2i+1, i+3\}}^N S_{ij} \leq C h^2 \int_{x_{2i}}^{x_N} y^{-\alpha/2-2/r-1} dy \\ = \frac{C}{\alpha/2 + 2/r} h^2 (x_{2i}^{-\alpha/2-2/r} - T^{-\alpha/2-2/r}) \\ \leq \frac{C}{\alpha/2 + 2/r} 2^{r(-\alpha/2-2/r)} h^2 x_i^{-\alpha/2-2/r}$$

114 Otherwise, for $N + 1 \leq j \leq 2N - 1$, by equation (C.2) and Lemma C.2

$$\begin{aligned}
 115 \quad (5.18) \quad \sum_{j=N+1}^{2N-1} S_{ij} &\leq Ch^2 \int_{x_N}^{x_{2N-1}} (2T - y)^{\alpha/2-2/r} y^{-1-\alpha} dy \\
 &\leq CT^{-1-\alpha} \frac{1}{|\alpha/2 - 2/r + 1|} h^2
 \end{aligned}$$

116 For $i = 1, 2$.

117 LEMMA 5.5. *By Lemma C.5 and Lemma 5.4 we get*

$$\begin{aligned}
 118 \quad (5.19) \quad R_1 &= \sum_{j=1}^3 S_{1j} + \sum_{j=4}^{2N} S_{1j} \\
 &\leq Ch^2 x_1^{-\alpha/2-2/r}
 \end{aligned}$$

119

$$\begin{aligned}
 120 \quad (5.20) \quad R_2 &= \sum_{j=1}^4 S_{2j} + \sum_{j=5}^{2N} S_{2j} \\
 &\leq Ch^2 x_2^{-\alpha/2-2/r}
 \end{aligned}$$

121 For $3 \leq i < N/2$, we have a new separation of R_i , Let's denote $k = \lceil \frac{i}{2} \rceil$.

$$\begin{aligned}
 R_i &= \sum_{j=1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &= \sum_{j=1}^{k-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right) \\
 122 \quad (5.21) \quad &\quad + \sum_{j=k+1}^{2i-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\
 &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i-1}} (T_{i-1,2i} + T_{i-1,2i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2i} \right) \\
 &\quad + \sum_{j=2i+1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &= I_1 + I_2 + I_3 + I_4 + I_5
 \end{aligned}$$

123 where I_1 makes sense only if $i \geq 3$.

124 For convience, let's denote

$$125 \quad (5.22) \quad V_{ij} = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right)$$

126

127 LEMMA 5.6. *There exists a constant $C = C(T, \alpha, r, f)$ such that for $3 \leq i <$*
 128 *$N/2, k = \lceil \frac{i}{2} \rceil$*

$$129 \quad (5.23) \quad I_1 \leq C(h^{r\alpha/2+r} x_i^{-1-\alpha} + h^2 x_i^{-\alpha/2-2/r})$$

130 *Proof.* For $2 \leq j \leq k-1$, by Lemma C.1 and Lemma C.3

$$\begin{aligned}
 S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) D_h^2 \left(\frac{|\cdot - y|^{1-\alpha}}{\Gamma(2-\alpha)} \right) (x_i) dy \\
 (5.24) \quad &\leq Ch^2 \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)} dy \\
 &= Ch^2 x_i^{-1-\alpha} \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} dy
 \end{aligned}$$

132 And by Lemma A.3 , Lemma C.3

$$(5.25) \quad S_{i1} \leq Cx_1^{\alpha/2} x_i^{-1-\alpha} = Cx_1^{\alpha/2+1} x_i^{-1-\alpha} = CT^{\alpha/2+1} h^{r\alpha/2+r} x_i^{-1-\alpha}$$

134 Therefore,

$$\begin{aligned}
 I_1 &= \sum_{j=1}^{k-1} S_{ij} = S_{i1} + \sum_{j=2}^{k-1} S_{ij} \\
 (5.26) \quad &\leq Ch^{r\alpha/2+r} x_i^{-1-\alpha} + Ch^2 x_i^{-1-\alpha} \int_{x_1}^{x_{\lceil \frac{i}{2} \rceil - 1}} y^{\alpha/2-2/r} dy \\
 &\leq Ch^{r\alpha/2+r} x_i^{-1-\alpha} + Ch^2 x_i^{-1-\alpha} \int_{x_1}^{2^{-r} x_i} y^{\alpha/2-2/r} dy
 \end{aligned}$$

136 But

$$(5.27) \quad \int_{x_1}^{2^{-r} x_i} y^{\alpha/2-2/r} dy \leq \begin{cases} \frac{1}{\alpha/2-2/r+1} (2^{-r} x_i)^{\alpha/2-2/r+1}, & \alpha/2-2/r+1 > 0 \\ \ln(2^{-r} x_i) - \ln(x_1), & \alpha/2-2/r+1 = 0 \\ \frac{1}{|\alpha/2-2/r+1|} x_1^{\alpha/2-2/r+1}, & \alpha/2-2/r+1 < 0 \end{cases}$$

138 So we have

$$(5.28) \quad I_1 \leq \begin{cases} Ch^2 x_i^{-1-\alpha} + \frac{C}{\alpha/2-2/r+1} h^2 x_i^{-\alpha/2-2/r}, & \alpha/2-2/r+1 > 0 \\ Ch^2 x_i^{-1-\alpha} (1 + r \ln(\frac{i}{2})), & \alpha/2-2/r+1 = 0 \\ Ch^{r\alpha/2+r} x_i^{-1-\alpha}, & \alpha/2-2/r+1 < 0 \end{cases} \quad \square$$

140 **THEOREM 5.7.** *There exists a constant $C = C(T, \alpha, r, f)$ such that for $2 \leq i <$*
 141 *$N/2, k = \lceil \frac{i}{2} \rceil$,*

$$(5.29) \quad I_3 = \sum_{j=k+1}^{2i-1} V_{ij} \leq Ch^2 x_i^{-\alpha/2-2/r}$$

143 To estimate V_{ij} , we need some preparations.

LEMMA 5.8. Denote $y_j^\theta = \theta x_{j-1} + (1 - \theta)x_j$, $\theta \in [0, 1]$, by Lemma A.2

$$\begin{aligned}
 T_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\
 &= \int_{x_{j-1}}^{x_j} -\frac{\theta(1-\theta)}{2} h_j^2 u''(y_j^\theta) \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} \\
 &\quad + \frac{\theta(1-\theta)}{3!} h_j^3 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} (\theta^2 u'''(\eta_{j1}^\theta) - (1-\theta)^2 u'''(\eta_{j2}^\theta)) dy_j^\theta \\
 &= \int_0^1 -\frac{\theta(1-\theta)}{2} h_j^3 u''(y_j^\theta) \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} \\
 &\quad + \frac{\theta(1-\theta)}{3!} h_j^4 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} (\theta^2 u'''(\eta_{j1}^\theta) - (1-\theta)^2 u'''(\eta_{j2}^\theta)) d\theta
 \end{aligned}
 \tag{5.30}$$

where $\eta_{j1}^\theta \in [x_{j-1}, y_j^\theta]$, $\eta_{j2}^\theta \in [y_j^\theta, x_j]$.

Now Let's construct a series of functions to represent T_{ij} .

$$y_{j-i}(x) = (x^{1/r} + z_{j-i})^r, \quad z_{j-i} = T^{1/r} \frac{j-i}{N}$$

$$y_{j-i}^\theta(x) = \theta y_{j-1-i}(x) + (1-\theta)y_{j-i}(x)$$

$$h_{j-i}(x) = y_{j-i}(x) - y_{j-i-1}(x)$$

Now, we define

$$P_{j-i}^\theta(x) = (h_{j-i}(x))^3 u''(y_{j-i}^\theta(x)) \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

$$Q_{j-i}^\theta(x) = (h_{j-i}(x))^4 \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

And now we can rewrite T_{ij}

LEMMA 5.9. For $2 \leq i \leq N$, $2 \leq j \leq \min\{2i-1, N\}$,

$$\begin{aligned}
 T_{ij} &= \int_0^1 -\frac{\theta(1-\theta)}{2} P_{j-i}^\theta(x_i) d\theta \\
 &\quad + \int_0^1 \frac{\theta(1-\theta)}{3!} (\theta^2 Q_{j-i}^\theta(x_i) u'''(\eta_{j1}^\theta) - (1-\theta)^2 Q_{j-i}^\theta(x_i) u'''(\eta_{j2}^\theta)) d\theta
 \end{aligned}
 \tag{5.36}$$

Immediately, we can see that

LEMMA 5.10. For $2 \leq i < N/2$, $k = \lceil \frac{i}{2} \rceil$, $k+1 \leq j \leq \min\{2i-1, N\}$,

LEMMA 5.11. There exists a constant $C = C(T, \alpha, r, f)$ such that for $2 \leq i < N$, $k = \lceil \frac{i}{2} \rceil$, $k+1 \leq j \leq \min\{2i-1, N\}$,

$$V_{ij} \leq \text{Sorry}$$

Proof.

$$P_{ij}^\theta = h_j^3 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} u''(y_j^\theta)$$

167

$$Q_{ij}^\theta = h_j^4 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)}$$

169

$$\begin{aligned} V_{ij} &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\ &= \int_0^1 -\frac{\theta(1-\theta)}{2} \left(\frac{1}{h_{i+1}} P_{i+1,j+1}^\theta - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) P_{i,j}^\theta + \frac{1}{h_i} P_{i-1,j-1}^\theta \right) d\theta \quad \square \\ &\quad + \int_0^1 \frac{\theta(1-\theta)}{3!} \left(\frac{1}{h_{i+1}} Q_{i+1,j+1}^\theta - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) Q_{i,j}^\theta + \frac{1}{h_i} Q_{i-1,j-1}^\theta \right) d\theta \end{aligned}$$

170

6. Proof of Theorem 4.2.

7. Experimental results.

8. Conclusions. Some conclusions here.

Appendix A. Approximate of difference quotients.

LEMMA A.1. *If $g(x)$ is twice differentiable continuous function on open set Ω , there exists $\xi \in [x_{i-1}, x_{i+1}]$ such that*

$$(A.1) \quad D_h^2 g(x_i) := \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = g''(\xi), \quad \xi \in [x_{i-1}, x_{i+1}]$$

$$(A.2) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \int_{x_{i-1}}^{x_i} g''(y)(y - x_{i-1}) dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} g''(y)(x_{i+1} - y) dy \right)$$

And if $g(x) \in C^4(\Omega)$, then

$$(A.3) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = g''(x_i) + \frac{h_{i+1} - h_i}{3} g'''(x_i) + \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 g''''(\eta_1) + h_{i+1}^3 g''''(\eta_2))$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$.

Proof.

$$g(x_{i-1}) = g(x_i) - (x_i - x_{i-1})g'(x_i) + \frac{(x_i - x_{i-1})^2}{2} g''(\xi_1), \quad \xi_1 \in [x_{i-1}, x_i]$$

$$g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i)g'(x_i) + \frac{(x_{i+1} - x_i)^2}{2} g''(\xi_2), \quad \xi_2 \in [x_i, x_{i+1}]$$

Substitute them in the left side of (A.1), we have

$$\frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = \frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2)$$

Now, using intermediate value theorem, there exists $\xi \in [\xi_1, \xi_2]$ such that

$$\frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2) = g''(\xi)$$

For the second equation, similarly

$$g(x_{i-1}) = g(x_i) - (x_i - x_{i-1})g'(x_i) + \int_{x_{i-1}}^{x_i} g''(y)(y - x_{i-1}) dy$$

$$g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i)g'(x_i) + \int_{x_i}^{x_{i+1}} g''(y)(x_{i+1} - y)dy$$

And the last equation can be obtained by

$$g(x_{i-1}) = g(x_i) - h_i g'(x_i) + \frac{h_i^2}{2} g''(x_i) - \frac{h_i^3}{3!} g'''(x_i) + \frac{h_i^4}{4!} g''''(\eta_1)$$

$$g(x_{i+1}) = g(x_i) + h_{i+1} g'(x_i) + \frac{h_{i+1}^2}{2} g''(x_i) + \frac{h_{i+1}^3}{3!} g'''(x_i) + \frac{h_{i+1}^4}{4!} g''''(\eta_2)$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$. Expecially,

$$\begin{aligned} \frac{h_i^4}{4!} g''''(\eta_1) &= \int_{x_{i-1}}^{x_i} g''''(y) \frac{(y - x_{i-1})^3}{3!} dy \\ \frac{h_{i+1}^4}{4!} g''''(\eta_2) &= \int_{x_i}^{x_{i+1}} g''''(y) \frac{(x_{i+1} - y)^3}{3!} dy \end{aligned}$$

Substitute them to the left side of (A.3), we can get the result. \square

LEMMA A.2. If $y \in [x_{j-1}, x_j]$, denote $y = \theta x_{j-1} + (1 - \theta)x_j$, $\theta \in [0, 1]$,

$$u(y_j^\theta) - u_h(y_j^\theta) = -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [x_{j-1}, x_j]$$

(A.6)

$$u(y_j^\theta) - u_h(y_j^\theta) = -\frac{\theta(1-\theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1-\theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1-\theta)^2 u'''(\eta_2))$$

where $\eta_1 \in [x_{j-1}, y_j^\theta]$, $\eta_2 \in [y_j^\theta, x_j]$.

Proof. By Taylor expansion, we have

$$u(x_{j-1}) = u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(\xi_1), \quad \xi_1 \in [x_{j-1}, y_j^\theta]$$

$$u(x_j) = u(y_j^\theta) + (1-\theta) h_j u'(y_j^\theta) + \frac{(1-\theta)^2 h_j^2}{2!} u''(\xi_2), \quad \xi_2 \in [y_j^\theta, x_j]$$

Thus

$$\begin{aligned} u(y_j^\theta) - u_h(y_j^\theta) &= u(y_j^\theta) - (1-\theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1-\theta)}{2} h_j^2 (\theta u''(\xi_1) + (1-\theta)u''(\xi_2)) \\ &= -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [\xi_1, \xi_2] \end{aligned}$$

The second equation is similar,

$$u(x_{j-1}) = u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(y_j^\theta) - \frac{\theta^3 h_j^3}{3!} u'''(\eta_1)$$

$$u(x_j) = u(y_j^\theta) + (1-\theta) h_j u'(y_j^\theta) + \frac{(1-\theta)^2 h_j^2}{2!} u''(y_j^\theta) + \frac{(1-\theta)^3 h_j^3}{3!} u'''(\eta_2)$$

where $\eta_1 \in [x_{j-1}, y_j^\theta]$, $\eta_2 \in [y_j^\theta, x_j]$. Thus \square

$$u(y_j^\theta) - u_h(y_j^\theta) = u(y_j^\theta) - (1-\theta)u(x_{j-1}) - \theta u(x_j)$$

$$= -\frac{\theta(1-\theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1-\theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1-\theta)^2 u'''(\eta_2))$$

LEMMA A.3. For $x \in [x_{j-1}, x_j]$

$$|u(x) - u_h(x)| = \left| \frac{x_j - x}{h_j} \int_{x_{j-1}}^x u'(y) dy - \frac{x - x_{j-1}}{h_j} \int_x^{x_j} u'(y) dy \right|$$

$$\leq \int_{x_{j-1}}^{x_j} |u'(y)| dy$$

If $x \in [0, x_1]$, with Corollary 2.4, we have

$$|u(x) - u_h(x)| \leq \int_0^{x_1} |u'(y)| dy \leq \int_0^{x_1} C y^{\alpha/2-1} dy \leq C \frac{2}{\alpha} x_1^{\alpha/2}$$

Similarly, if $x \in [x_{2N-1}, 1]$, we have

$$|u(x) - u_h(x)| \leq C \frac{2}{\alpha} (2T - x_{2N-1})^{\alpha/2} = C \frac{2}{\alpha} x_1^{\alpha/2}$$

Appendix B. Inequality.

LEMMA B.1.

$$h_i \leq r T^{1/r} h \begin{cases} x_i^{1-1/r}, & 1 \leq i \leq N \\ (2T - x_{i-1})^{1-1/r}, & N < i \leq 2N-1 \end{cases}$$

Proof. For $1 \leq i \leq N$,

$$h_i = T \left(\left(\frac{i}{N} \right)^r - \left(\frac{i-1}{N} \right)^r \right)$$

$$\leq r T \frac{1}{N} \left(\frac{i}{N} \right)^{r-1} = r T^{1/r} h x_i^{1-1/r}$$

For $N < i \leq 2N-1$,

$$h_i = T \left(\left(\frac{2N-i+1}{N} \right)^r - \left(\frac{2N-i}{N} \right)^r \right)$$

$$\leq r T \frac{1}{N} \left(\frac{2N-i+1}{N} \right)^{r-1} = r T^{1/r} h (2T - x_{i-1})^{1-1/r}$$

□

LEMMA B.2. There is a constant $C = 2^{|r-2|} r(r-1) T^{2/r}$ such that for all $i \in \{1, 2, \dots, 2N-1\}$

$$|h_{i+1} - h_i| \leq C h^2 \begin{cases} x_i^{1-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{1-2/r}, & N < i \leq 2N-1 \end{cases}$$

Proof.

$$h_{i+1} - h_i = \begin{cases} T \left(\left(\frac{i+1}{N} \right)^r - 2 \left(\frac{i}{N} \right)^r + \left(\frac{i-1}{N} \right)^r \right), & 1 \leq i \leq N-1 \\ 0, & i = N \\ -T \left(\left(\frac{2N-i-1}{N} \right)^r - 2 \left(\frac{2N-i}{N} \right)^r + \left(\frac{2N-i+1}{N} \right)^r \right), & N+1 \leq i \leq 2N-1 \end{cases}$$

230 For $i = 1$,

$$231 \quad h_2 - h_1 = T(2^r - 2) \left(\frac{1}{N} \right)^r = (2^r - 2)T^{2/r} h^2 x_1^{1-2/r}$$

232 For $2 \leq i \leq N - 1$,

$$233 \quad h_{i+1} - h_i = r(r-1)T N^{-2} \eta^{r-2}, \quad \eta \in \left[\frac{i-1}{N}, \frac{i+1}{N} \right]$$

234 If $r \in [1, 2]$,

$$\begin{aligned} 235 \quad h_{i+1} - h_i &= r(r-1)T N^{-2} \eta^{r-2} \leq r(r-1)T h^2 \left(\frac{i-1}{N} \right)^{r-2} \\ &\leq r(r-1)T h^2 2^{2-r} \left(\frac{i}{N} \right)^{r-2} \\ &= 2^{2-r} r(r-1)T^{2/r} h^2 x_i^{1-2/r} \end{aligned}$$

236 else if $r > 2$,

$$\begin{aligned} h_{i+1} - h_i &= r(r-1)T N^{-2} \eta^{r-2} \leq r(r-1)T h^2 \left(\frac{i+1}{N} \right)^{r-2} \\ 237 \quad &\leq r(r-1)T h^2 2^{r-2} \left(\frac{i}{N} \right)^{r-2} \\ &= 2^{r-2} r(r-1)T^{2/r} h^2 x_i^{1-2/r} \end{aligned}$$

238 Since

$$239 \quad 2^r - 2 \leq 2^{|r-2|} r(r-1), \quad r \geq 1$$

240 we have

$$241 \quad h_{i+1} - h_i \leq 2^{|r-2|} r(r-1)T^{2/r} h^2 x_i^{1-2/r}, \quad 1 \leq i \leq N-1$$

242 For $i = N$, $h_{N+1} - h_N = 0$. For $N < i \leq 2N - 1$, it's central symmetric to the first
243 half of the proof, which is

$$244 \quad h_i - h_{i+1} \leq 2^{|r-2|} r(r-1)T^{2/r} h^2 (2T - x_i)^{1-2/r}$$

245 Summarizes the inequalities, we can get

$$246 \quad (\text{B.3}) \quad |h_{i+1} - h_i| \leq 2^{|r-2|} r(r-1)T^{2/r} h^2 \begin{cases} x_i^{1-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{1-2/r}, & N < i \leq 2N-1 \end{cases} \quad \square$$

247 **Appendix C. Proofs of some technical details.**

248 *Additional proof of Theorem 5.1.* For $2 \leq i \leq N-1$,

$$\begin{aligned} & \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \\ 249 & \leq C \frac{2}{h_i + h_{i+1}} (h_i^3 x_{i-1}^{-2-\alpha/2} + h_{i+1}^3 x_i^{-2-\alpha/2}) \\ & \leq 2C (h_i^2 x_{i-1}^{-2-\alpha/2} + h_{i+1}^2 x_i^{-2-\alpha/2}) \end{aligned}$$

250 Since Lemma B.1, we have

$$\begin{aligned} 251 & h_i \leq r T^{1/r} h x_i^{1-1/r}, \quad 1 \leq i \leq N \\ 252 & h_{i+1} \leq r T^{1/r} h x_{i+1}^{1-1/r}, \quad 1 \leq i \leq N-1 \end{aligned}$$

253 and

$$\begin{aligned} 254 & x_{i-1}^{-2-\alpha/2} \leq 2^{-r(-2-\alpha/2)} x_i^{-2-\alpha/2} \quad 2 \leq i \leq N-1 \\ 255 & x_{i+1}^{1-1/r} \leq 2^{r-1} x_i^{1-1/r} \quad 1 \leq i \leq N-1 \end{aligned}$$

256 So there is a constant $C = C(T, \alpha, r, \|f\|_\beta^{\alpha/2})$ such that

$$257 \quad \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \leq C h^2 x_i^{-\alpha/2-2/r}, \quad 2 \leq i \leq N-1$$

258 For $i = 1$, by (A.4)

$$\begin{aligned} & \frac{1}{4!} \frac{2}{h_1 + h_2} (h_1^3 f''(\eta_1) + h_2^3 f''(\eta_2)) \\ 259 & = \frac{2}{h_1 + h_2} \left(\frac{1}{h_1} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy + \frac{1}{4!} h_2^3 f''(\eta_2) \right) \end{aligned}$$

260 We have proved above that

$$261 \quad \frac{2}{h_1 + h_2} h_2^3 f''(\eta_2) \leq C h^2 x_1^{-\alpha/2-2/r}$$

262 and we can get

$$\begin{aligned} 263 & \int_0^{x_1} f''(y) \frac{y^3}{3!} dy \leq C \frac{1}{3!} \int_0^{x_1} y^{1-\alpha/2} dy \\ & = C \frac{1}{3!(2-\alpha/2)} x_1^{2-\alpha/2} \end{aligned}$$

264 so

$$265 \quad \frac{2}{h_1 + h_2} \frac{1}{h_1} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy = \frac{C 2^{1-r}}{3!(2-\alpha/2)} x_1^{-\alpha/2} = \frac{C 2^{1-r}}{3!(2-\alpha/2)} T^{2/r} h^2 x_1^{-\alpha/2-2/r}$$

266 And for $i = N$, we have

$$\begin{aligned} & \frac{2}{h_N + h_{N+1}} (h_N^3 f''(\eta_1) + h_{N+1}^3 f''(\eta_2)) \\ 267 & = h_N^2 (f''(\eta_1) + f''(\eta_2)) \\ & \leq r^2 T^{2/r} h^2 x_N^{2-2/r} 2C x_{N-1}^{-2-\alpha/2} \\ & \leq 2r^2 T^{2/r} C 2^{-r(-2-\alpha/2)} h^2 x_N^{-\alpha/2-2/r} \end{aligned}$$

268 Finally, $N + 1 \leq i \leq 2N - 1$ is symmetric to the first half of the proof, so we can
 269 conclude that □

$$270 \quad \frac{2}{h_i + h_{i+1}}(h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \leq Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T - x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N - 1 \end{cases}$$

271 LEMMA C.1. *There is a constant $C = C(T, \alpha, r, f)$ for $2 \leq j \leq N$, if $y \in$
 272 $[x_{j-1}, x_j]$,*

$$273 \quad (C.1) \quad |u(y) - u_h(y)| \leq Ch^2 y^{\alpha/2-2/r}$$

274 *Proof.* For $2 \leq j \leq N$, we have

$$275 \quad x_j \leq 2^r y, \quad x_{j-1} \geq 2^{-r} y$$

276 And by Lemma A.2, Lemma B.1 and Corollary 2.4, we have

$$\begin{aligned} u(y) - u_h(y) &= -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [x_{j-1}, x_j] \\ &\leq \frac{C}{4} h^2 x_j^{2-2/r} x_{j-1}^{\alpha/2-2} \\ &\leq \frac{C}{4} h^2 2^{2r-2} y^{2-2/r} 2^{-r(\alpha/2-2)} y^{\alpha/2-2} \\ &= C 2^{-r\alpha/2+4r-2} h^2 y^{\alpha/2-2/r} \end{aligned}$$

278 symmetricly, for $N < j \leq 2N - 1$, we have

$$279 \quad (C.2) \quad |u(y) - u_h(y)| \leq Ch^2 (2T - y)^{\alpha/2-2/r} \quad \square$$

280 LEMMA C.2. *There is a constant $C = C(\alpha, r)$ such that for all $1 \leq i < N/2$,
 281 $\max\{2i + 1, i + 3\} \leq j \leq 2N$ and $y \in [x_{j-1}, x_j]$, we have*

$$282 \quad (C.3) \quad D_h^2 \left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)} \right)(x_i) \leq C \frac{y^{-1-\alpha}}{\Gamma(-\alpha)}$$

283 *Proof.* Since $y \geq x_{j-1} > x_{i+1}$, by Lemma A.1, if $j - 1 > i + 1$ □

$$\begin{aligned} D_h^2 \left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)} \right)(x_i) &= \frac{|y - \xi|^{-1-\alpha}}{\Gamma(-\alpha)}, \quad \xi \in [x_{i-1}, x_{i+1}] \\ &\leq \frac{(y - x_{i+1})^{-1-\alpha}}{\Gamma(-\alpha)} \\ &\leq \left(1 - \left(\frac{2}{3}\right)^r\right)^{-1-\alpha} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} \end{aligned}$$

285 LEMMA C.3. *There is a constant $C = C(\alpha, r)$ such that for all $3 \leq i < N/2, k =$
 286 $\lceil \frac{i}{2} \rceil, 1 \leq j \leq k - 1$ and $y \in [x_{j-1}, x_j]$, we have*

$$287 \quad (C.4) \quad D_h^2 \left(\frac{|\cdot - y|^{1-\alpha}}{\Gamma(2-\alpha)} \right)(x_i) \leq C \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)}$$

Proof. Since $y \leq x_j < x_{i-1}$, by Lemma A.1,

$$\begin{aligned} D_h^2\left(\frac{|\cdot - y|^{1-\alpha}}{\Gamma(2-\alpha)}\right)(x_i) &= \frac{|\xi - y|^{-1-\alpha}}{\Gamma(-\alpha)}, \quad \xi \in [x_{i-1}, x_{i+1}] \\ &\leq \frac{(x_{i-1} - x_j)^{-1-\alpha}}{\Gamma(-\alpha)} \leq \frac{(x_{i-1} - x_{k-1})^{-1-\alpha}}{\Gamma(-\alpha)} \\ &\leq \left(\left(\frac{2}{3}\right)^r - \left(\frac{1}{2}\right)^r\right)^{-1-\alpha} \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)} \end{aligned}$$

□

LEMMA C.4. While $0 \leq i < N/2$, By Lemma A.3

$$\begin{aligned} |T_{i1}| &\leq C \int_0^{x_1} x_1^{\alpha/2} \frac{|x_i - y|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\ &= C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2} |x_i^{2-\alpha} - |x_i - x_1|^{2-\alpha}| \\ &\leq C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2+2-\alpha} = C \frac{1}{\Gamma(3-\alpha)} x_1^{2-\alpha/2} \quad 0 < 2-\alpha < 1 \end{aligned}$$

For $2 \leq j \leq N$, by Lemma A.2 and Corollary 2.4

$$\begin{aligned} |T_{ij}| &\leq \frac{C}{4} \int_{x_{j-1}}^{x_j} h_j^2 x_{j-1}^{\alpha/2-2} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\ &\leq \frac{C}{4\Gamma(3-\alpha)} h_j^2 x_{j-1}^{\alpha/2-2} ||x_j - x_i|^{2-\alpha} - |x_{j-1} - x_i|^{2-\alpha}| \end{aligned}$$

LEMMA C.5. There exists a constant $C = C(T, \alpha, r, f)$ such that

$$\sum_{j=1}^3 S_{1j} \leq Ch^2 x_1^{-\alpha/2-2/r}$$

$$\sum_{j=1}^4 S_{2j} \leq Ch^2 x_2^{-\alpha/2-2/r}$$

Proof.

$$S_{1j} = \frac{2}{x_2} \left(\frac{1}{x_1} T_{0j} - \left(\frac{1}{x_1} + \frac{1}{h_2} \right) T_{1j} + \frac{1}{h_2} T_{2j} \right)$$

So, by Lemma C.4

$$S_{11} \leq \frac{2}{x_2 x_1} 4 \frac{C}{\Gamma(3-\alpha)} x_1^{2-\alpha/2} \leq C x_1^{-\alpha/2}$$

$$S_{12} \leq \frac{2}{x_2 x_1} \frac{C}{4\Gamma(3-\alpha)} h_2^2 x_1^{\alpha/2-2} (x_2^{2-\alpha} + 2h_2^{2-\alpha} + h_2^{2-\alpha}) \leq C x_1^{-\alpha/2}$$

$$S_{13} \leq \frac{2}{x_2 x_1} \frac{C}{4\Gamma(3-\alpha)} h_3^2 x_2^{\alpha/2-2} (x_3^{2-\alpha} + 2x_3^{2-\alpha} + h_3^{2-\alpha}) \leq C x_1^{-\alpha/2}$$

307 But

308
$$x_1^{-\alpha/2} = T^{2/r} h^2 x_1^{-\alpha/2-2/r}$$

309 For $i = 2$, Sorry

□

310 **Acknowledgments.** We would like to acknowledge the assistance of volunteers
 311 in putting together this example manuscript and supplement.

312

REFERENCES

- 313 [1] X. ROS-OTON AND J. SERRA, *The dirichlet problem for the fractional laplacian: Regular-*
 314 *ity up to the boundary*, Journal de Mathématiques Pures et Appliquées, 101 (2014),
 315 pp. 275–302, <https://doi.org/https://doi.org/10.1016/j.matpur.2013.06.003>, <https://www.sciencedirect.com/science/article/pii/S0021782413000895>.
 316