

A SECOND ORDER NUMERICAL METHODS FOR REISZ-FRACTIONAL ELLIPTIC EQUATION ON GRADED MESH*

JIANXING HAN[†] AND MINGHUA CHEN[‡]

Abstract. This is an example SIAM L^AT_EX article. This can be used as a template for new articles. Abstracts must be able to stand alone and so cannot contain citations to the paper's references, equations, etc. An abstract must consist of a single paragraph and be concise. Because of online formatting, abstracts must appear as plain as possible. Any equations should be inline.

Key words. example, L^AT_EX

MSC codes. ??????????????????

1. Introduction. For $\Omega = (0, 2T)$, $1 < \alpha < 2$,

$$(1.1) \quad \begin{cases} (-\Delta)^{\frac{\alpha}{2}} u(x) = f(x), & x \in \Omega \\ u(x) = 0, & x \in \mathbb{R} \setminus \Omega \end{cases}$$

where

$$(1.2) \quad (-\Delta)^{\frac{\alpha}{2}} u(x) = -\frac{\partial^\alpha u}{\partial |x|^\alpha} = -\kappa_\alpha \frac{d^2}{dx^2} \int_\Omega \frac{|x-y|^{1-\alpha}}{\Gamma(2-\alpha)} u(y) dy$$

$$(1.3) \quad \kappa_\alpha = -\frac{1}{2 \cos(\alpha\pi/2)} > 0$$

2. Preliminaries: Numeric scheme and main results.

2.1. Numeric Format.

$$(2.1) \quad x_i = \begin{cases} T \left(\frac{i}{N} \right)^r, & 0 \leq i \leq N \\ 2T - T \left(\frac{2N-i}{N} \right)^r, & N \leq i \leq 2N \end{cases}$$

where $r \geq 1$. And let

$$(2.2) \quad h_j = x_j - x_{j-1}, \quad 1 \leq j \leq 2N$$

Let $\{\phi_j(x)\}_{j=1}^{2N-1}$ be standard hat functions, which are basis of the piecewise linear function space.

$$(2.3) \quad \phi_j(x) = \begin{cases} \frac{1}{h_j}(x - x_{j-1}), & x_{j-1} \leq x \leq x_j \\ \frac{1}{h_{j+1}}(x_{j+1} - x), & x_j \leq x \leq x_{j+1} \\ 0, & \text{otherwise} \end{cases}$$

And then, define the piecewise linear interpolant of the true solution u to be

$$(2.4) \quad \Pi_h u(x) := \sum_{j=1}^{2N-1} u(x_j) \phi_j(x)$$

*Submitted to the editors DATE.

[†]School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, PR China (hanjx2023@mail.lzu.edu.cn).

[‡]School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, PR China (chen@mail.lzu.edu.cn).

For convience, we denote

$$(2.5) \quad I^{2-\alpha}u(x) := \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x-y|^{1-\alpha} u(y) dy$$

and

$$(2.6) \quad D_h^2 u(x_i) := \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} u(x_{i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) u(x_i) + \frac{1}{h_{i+1}} u(x_{i+1}) \right)$$

Now, we discretise (1.1) by replacing $u(x)$ by a continuous piecewise linear function

$$(2.7) \quad u_h(x) := \sum_{j=1}^{2N-1} u_j \phi_j(x)$$

whose nodal values u_j are to be determined by collocation at each mesh point x_i for $i = 1, 2, \dots, 2N-1$:

$$(2.8) \quad -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) := -\kappa_{\alpha} D_h^2 I^{2-\alpha} u_h(x_i) = f(x_i) =: f_i$$

Here,

$$(2.9) \quad -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) = \sum_{j=1}^{2N-1} -\kappa_{\alpha} D_h^2 I^{2-\alpha} \phi_j(x_i) u_j = \sum_{j=1}^{2N-1} a_{ij} u_j$$

where

$$(2.10) \quad a_{ij} = -\kappa_{\alpha} D_h^2 I^{2-\alpha} \phi_j(x_i) \quad \text{for } i, j = 1, 2, \dots, 2N-1$$

We have replaced $(-\Delta)^{\alpha/2} u(x_i) = f(x_i)$ in (1.1) by $-\kappa_{\alpha} D_h^{\alpha} u_h(x_i) = f(x_i)$ in (2.8), with truncation error

$$(2.11) \quad \tau_i := -\kappa_{\alpha} \left(D_h^{\alpha} \Pi_h u(x_i) - \frac{d^2}{dx^2} I^{2-\alpha} u(x_i) \right) \quad \text{for } i = 1, 2, \dots, 2N-1$$

where $-\kappa_{\alpha} D_h^{\alpha} \Pi_h u(x_i) = \sum_{j=1}^{2N-1} -\kappa_{\alpha} D_h^{\alpha} \phi_j(x_i) u(x_j) = \sum_{j=1}^{2N-1} a_{ij} u(x_j)$.

The discrete equation (2.8) can be written in matrix form

$$(2.12) \quad AU = F$$

where $A = (a_{ij}) \in \mathbb{R}^{(2N-1) \times (2N-1)}$, $U = (u_1, \dots, u_{2N-1})^T$ is unknown and $F = (f_1, \dots, f_{2N-1})^T$.

We can deduce a_{ij} ,

$$(2.13) \quad \begin{aligned} a_{ij} &= -\kappa_{\alpha} D_h^2 I^{2-\alpha} \phi_j(x_i) \\ &= -\kappa_{\alpha} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \tilde{a}_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{a}_{i,j} + \frac{1}{h_{i+1}} \tilde{a}_{i+1,j} \right) \end{aligned}$$

where

$$(2.14) \quad \begin{aligned} \tilde{a}_{ij} &= I^{2-\alpha} \phi_j(x_i) \\ &= \frac{1}{\Gamma(4-\alpha)} \left(\frac{|x_i - x_{j-1}|^{3-\alpha}}{h_j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) |x_i - x_j|^{3-\alpha} + \frac{|x_i - x_{j+1}|^{3-\alpha}}{h_{j+1}} \right) \end{aligned}$$

We shall finally introduce some notation.

For convenience, we use the notation \simeq . That $x_1 \simeq y_1$, means that $c_1 x_1 \leq y_1 \leq C_1 x_1$ for some constants c_1 and C_1 that are independent of N .

Meanwhile, let

$$(2.15) \quad K_y(x) := \frac{|y-x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

We define the difference quotients

$$(2.16) \quad D_h g(x_i) := \frac{g(x_{i+1}) - g(x_i)}{h_{i+1}}, \quad D_{\bar{h}} g(x_i) := \frac{g(x_i) - g(x_{i-1})}{h_i}$$

Thus

$$D_h g(x_i) = D_{\bar{h}} g(x_{i+1})$$

$$D_h^2 g(x_i) = \frac{2}{h_i + h_{i+1}} (D_h g(x_i) - D_{\bar{h}} g(x_i)) = \frac{2}{h_i + h_{i+1}} (D_h g(x_i) - D_{\bar{h}} g(x_{i-1}))$$

And for $j = 1, 2, \dots, 2N$, we define

$$(2.17) \quad y_j^\theta = (1-\theta)x_{j-1} + \theta x_{j+1}, \quad \theta \in (0, 1)$$

2.2. Regularity of the true solution. For any $\beta > 0$, we use the standard notation $C^\beta(\bar{\Omega})$, $C^\beta(\mathbb{R})$, etc., for Hölder spaces and their norms and seminorms. When no confusion is possible, we use the notation $C^\beta(\Omega)$ to refer to $C^{k,\beta'}(\Omega)$, where k is the greatest integer such that $k < \beta$ and where $\beta' = \beta - k$. The Hölder spaces $C^{k,\beta'}(\Omega)$ are defined as the subspaces of $C^k(\Omega)$ consisting of functions whose k -th order partial derivatives are locally Hölder continuous[1] with exponent β' in Ω , where $C^k(\Omega)$ is the set of all k -times continuously differentiable functions on open set Ω .

DEFINITION 2.1 (delta dependent norm [2]). ...

LEMMA 2.2. Let $f \in C^\beta(\Omega)$, $\beta > 2$ be such that $\|f\|_\beta^{(\alpha/2)} < \infty$, then for $l = 0, 1, 2$

$$(2.18) \quad |f^{(l)}(x)| \leq \|f\|_\beta^{(\alpha/2)} \begin{cases} x^{-l-\alpha/2}, & \text{if } 0 < x \leq T \\ (2T-x)^{-l-\alpha/2}, & \text{if } T \leq x < 2T \end{cases}$$

THEOREM 2.3 (Regularity up to the boundary [2]). Let Ω be a bounded domain, and $\beta > 0$ be such that neither β nor $\beta + \alpha$ is an integer. Let $f \in C^\beta(\Omega)$ be such that $\|f\|_\beta^{(\alpha/2)} < \infty$, and $u \in C^{\alpha/2}(\mathbb{R}^n)$ be a solution of (1.1). Then, $u \in C^{\beta+\alpha}(\Omega)$ and

$$(2.19) \quad \|u\|_{\beta+\alpha}^{(-\alpha/2)} \leq C \left(\|u\|_{C^{\alpha/2}(\mathbb{R})} + \|f\|_\beta^{(\alpha/2)} \right)$$

where C is a constant depending only on Ω , α , and β .

COROLLARY 2.4. Let u be a solution of (1.1) where $f \in L^\infty(\Omega)$ and $\|f\|_\beta^{(\alpha/2)} < \infty$. Then, for any $x \in \Omega$ and $l = 0, 1, 2, 3, 4$

$$(2.20) \quad |u^{(l)}(x)| \leq \|u\|_{\beta+\alpha}^{(-\alpha/2)} \begin{cases} x^{\alpha/2-l}, & \text{if } 0 < x \leq T \\ (2T-x)^{\alpha/2-l}, & \text{if } T \leq x < 2T \end{cases}$$

And in this paper bellow, without special instructions, we allways assume that

$$(2.21) \quad f \in L^\infty(\Omega) \cap C^\beta(\Omega) \quad \text{and} \quad \|f\|_\beta^{(\alpha/2)} < \infty, \quad \text{with } \alpha + \beta > 4$$

2.3. Main results. Here we state our main results; the proof is deferred to section 3 and section 4.

Let's denote $h = \frac{1}{N}$, we have

THEOREM 2.5 (Local Truncation Error). *If $u(x)$ is a solution of the equation (1.1) where f satisfy the regular condition (2.21), then there exists $C_1(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)}, \|f\|_{\beta}^{(\alpha/2)})$ and $C_2(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$, such that the truncation error (2.11) satisfies*

$$\begin{aligned} |\tau_i| &:= |-\kappa_{\alpha} D_h^{\alpha} \Pi_h u(x_i) - f(x_i)| \\ &\leq C_1 h^{\min\{\frac{r\alpha}{2}, 2\}} \begin{cases} x_i^{-\alpha}, & 1 \leq i \leq N \\ (2T - x_i)^{-\alpha}, & N < i \leq 2N - 1 \end{cases} \\ &\quad + C_2(r-1)h^2 \begin{cases} |T - x_{i-1}|^{1-\alpha}, & 1 \leq i \leq N \\ |T - x_{i+1}|^{1-\alpha}, & N < i \leq 2N - 1 \end{cases} \end{aligned} \quad (2.22)$$

THEOREM 2.6 (Global Error). *The discrete equation (2.8) has solution and there exists a positive constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)}, \|f\|_{\beta}^{(\alpha/2)})$ such that the error between the numerical solution U with the exact solution $u(x_i)$ satisfies*

$$(2.23) \quad \max_{1 \leq i \leq 2N-1} |u_i - u(x_i)| \leq Ch^{\min\{\frac{r\alpha}{2}, 2\}}$$

That means the numerical method has convergence order $\min\{\frac{r\alpha}{2}, 2\}$.

Remark 2.7. ...

3. Local Truncation Error.

3.1. Proof of Theorem 2.5. The truncation error of the discrete format can be written as

$$\begin{aligned} -\kappa_{\alpha} D_h^{\alpha} \Pi_h u(x_i) - f(x_i) &= -\kappa_{\alpha} (D_h^2 I^{2-\alpha} \Pi_h u(x_i) - \frac{d^2}{dx^2} I^{2-\alpha} u(x_i)) \\ &= -\kappa_{\alpha} D_h^2 I^{2-\alpha} (\Pi_h u - u)(x_i) - \kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha} u(x_i) \end{aligned}$$

THEOREM 3.1. *There exists a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$ such that*

$$(3.2) \quad \left| -\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha} u(x_i) \right| \leq Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T - x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N - 1 \end{cases}$$

Proof. Since $f \in C^2(\Omega)$ and

$$(3.3) \quad \frac{d^2}{dx^2} (-\kappa_{\alpha} I^{2-\alpha} u(x)) = f(x), \quad x \in \Omega,$$

we have $I^{2-\alpha} u \in C^4(\Omega)$. Therefore, using equation (A.2) of Lemma A.1, for $1 \leq i \leq$

111 $2N - 1$, we have
 (3.4)

$$112 \quad -\kappa_\alpha(D_h^2 - \frac{d^2}{dx^2})I^{2-\alpha}u(x_i) = \frac{h_{i+1} - h_i}{3}f'(x_i) \\ + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \int_{x_{i-1}}^{x_i} f''(y) \frac{(y - x_{i-1})^3}{3!} dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} f''(y) \frac{(y - x_{i+1})^3}{3!} dy \right)$$

113 By Lemma B.2, Lemma 2.2 and Lemma B.3, we get the result. \square

114 And now define

$$115 \quad (3.5) \quad R_i := D_h^2 I^{2-\alpha}(u - \Pi_h u)(x_i)$$

116 We have some results about the estimate of R_i

117 **THEOREM 3.2.** *For $1 \leq i < N/2$, there exists $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that*

$$118 \quad (3.6) \quad R_i \leq \begin{cases} Ch^2 x_i^{-\alpha/2-2/r}, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2(x_i^{-1-\alpha} \ln(i) + \ln(N)), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r} x_i^{-1-\alpha}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

119
 120 **THEOREM 3.3.** *For $N/2 \leq i \leq N$, there exists constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$*
 121 *such that*

$$122 \quad (3.7) \quad R_i \leq C(r-1)h^2|T - x_{i-1}|^{1-\alpha} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

123 And for $N < i \leq 2N - 1$, it is symmetric to the previous case.

124 Combine Theorem 3.1, Theorem 3.2 and Theorem 3.3, the proof of Theorem 2.5
 125 completed.

126 We prove Theorem 3.2 and Theorem 3.3 in next subsections below.

127 **3.2. Mesh Transport Functions.**

128 **DEFINITION 3.4** (Mesh Transport Functions).

$$129 \quad (3.8) \quad y_{i,j}(x) = \begin{cases} (x^{1/r} + Z_{j-i})^r & i < N, j < N \\ \frac{x^{1/r} - Z_i}{Z_1} h_N + x_N & i < N, j = N \\ 2T - (Z_{2N-(j-i)} - x^{1/r})^r & i < N, j > N \\ \left(\frac{Z_1}{h_N} (x - x_N) + Z_j \right)^r & i = N, j < N \\ x, & i = N, j = N \end{cases}$$

130 *where*

$$131 \quad (3.9) \quad Z_j := T^{1/r} \frac{j}{N}$$

132 *And*

$$133 \quad (3.10) \quad h_{i,j}(x) = y_{i,j}(x) - y_{i,j-1}(x)$$

$$135 \quad (3.11) \quad y_{i,j}^\theta(x) = (1 - \theta)y_{i,j-1}(x) + \theta y_{i,j}(x), \quad \theta \in (0, 1)$$

We give some properties of mesh transport functions.

LEMMA 3.5. *Obviously,*

$$(3.12) \quad y_{i,j}(x_{i-1}) = x_{j-1}, \quad y_{i,j}(x_i) = x_j, \quad y_{i,j}(x_{i+1}) = x_{j+1}$$

$$(3.13) \quad h_{i,j}(x_{i-1}) = h_{j-1}, \quad h_{i,j}(x_i) = h_j, \quad h_{i,j}(x_{i+1}) = h_{j+1}$$

$$(3.14) \quad y_{i,j}^\theta(x_{i-1}) = y_{j-1}^\theta, \quad y_{i,j}^\theta(x_i) = y_j^\theta, \quad y_{i,j}^\theta(x_{i+1}) = y_{j+1}^\theta$$

LEMMA 3.6. *For $1 \leq i \leq 2N-1, 2 \leq j \leq 2N-1$,*

$$(3.15) \quad h_{i,j}(\xi) \simeq h_j, \quad \text{for } \xi \in (x_{i-1}, x_{i+1})$$

For $1 \leq i, j \leq 2N-1$,

$$(3.16) \quad |y_{i,j}(\xi) - \xi| \simeq |x_j - x_i| \quad \text{for } \xi \in (x_{i-1}, x_{i+1})$$

3.3. Proof of Theorem 3.2. For convenience, let's denote

$$(3.17) \quad T_{ij} = \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy, \quad i = 0, \dots, 2N, \quad j = 1, \dots, 2N$$

Also for simplicity, we denote

DEFINITION 3.7.

$$(3.18) \quad S_{ij} = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} T_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_{i+1}} T_{i+1,j} \right)$$

then

$$(3.19) \quad R_i = \sum_{j=1}^{2N} S_{ij}$$

LEMMA 3.8. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $1 \leq i < N/2$,*

$$(3.20) \quad \sum_{j=\max\{2i+1, i+3\}}^N S_{ij} \leq Ch^2 x_i^{-\alpha/2-2/r}$$

Proof. For $\max\{2i+1, i+3\} \leq j \leq N$, by Lemma B.4 and Lemma B.5

$$(3.21) \quad \begin{aligned} S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) D_h^2 K_y(x_i) dy \\ &\leq Ch^2 \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} dy \\ &= Ch^2 \int_{x_{j-1}}^{x_j} y^{-\alpha/2-2/r-1} dy \end{aligned}$$

Therefore,

$$(3.22) \quad \begin{aligned} \sum_{j=\max\{2i+1, i+3\}}^N S_{ij} &\leq Ch^2 \int_{x_{2i}}^{x_N} y^{-\alpha/2-2/r-1} dy \\ &= \frac{C}{\alpha/2 + 2/r} h^2 (x_{2i}^{-\alpha/2-2/r} - T^{-\alpha/2-2/r}) \\ &\leq \frac{C}{\alpha/2 + 2/r} 2^{r(-\alpha/2-2/r)} h^2 x_i^{-\alpha/2-2/r} \end{aligned} \quad \square$$

LEMMA 3.9. *Thert exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $1 \leq i < N/2$,*

$$(3.23) \quad \sum_{j=N+1}^{2N} S_{ij} \leq \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

Proof. For $1 \leq i < N/2, N+1 \leq j \leq 2N-1$, by equation (B.12) and Lemma B.5

$$\begin{aligned} S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) D_h^2 K_y(x_i) dy \\ &\leq \int_{x_{j-1}}^{x_j} Ch^2 (2T - y)^{\alpha/2-2/r} y^{-1-\alpha} dy \\ &\leq Ch^2 T^{-1-\alpha} \int_{x_{j-1}}^{x_j} (2T - y)^{\alpha/2-2/r} dy \end{aligned}$$

$$\begin{aligned} \sum_{j=N+1}^{2N-1} S_{ij} &\leq CT^{-1-\alpha} h^2 \int_{x_N}^{x_{2N-1}} (2T - y)^{\alpha/2-2/r} dy \\ &\leq CT^{-1-\alpha} h^2 \begin{cases} \frac{1}{\alpha/2-2/r+1} T^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 > 0 \\ \ln(T) - \ln(h_{2N}), & \alpha/2 - 2/r + 1 = 0 \\ \frac{1}{|\alpha/2-2/r+1|} h_{2N}^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \\ &= \begin{cases} \frac{C}{\alpha/2-2/r+1} T^{-\alpha/2-2/r} h^2, & \alpha/2 - 2/r + 1 > 0 \\ CrT^{-1-\alpha} h^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ \frac{C}{|\alpha/2-2/r+1|} T^{-\alpha/2-2/r} h^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \end{aligned}$$

And by Lemma A.3

$$S_{i,2N} \leq CT^{-1-\alpha} h_{2N}^{\alpha/2+1} = CT^{-\alpha/2} h^{r\alpha/2+r}$$

And when $\alpha/2 - 2/r + 1 \geq 0$,

$$h^{r\alpha/2+r} \leq h^2$$

Summarizes, we get the result. □

For $i = 1, 2$.


LEMMA 3.10. *By Lemma B.8 , Lemma 3.8 and Lemma 3.9 we get*

$$\begin{aligned} R_1 &= \sum_{j=1}^3 S_{1j} + \sum_{j=4}^{2N} S_{1j} \\ &\leq Ch^2 x_1^{-\alpha/2-2/r} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \end{aligned}$$

175

$$\begin{aligned}
R_2 &= \sum_{j=1}^4 S_{2j} + \sum_{j=5}^{2N} S_{2j} \\
(3.26) \quad &\leq Ch^2 x_2^{-\alpha/2-2/r} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}
\end{aligned}$$

177 For $3 \leq i < N/2$, we have a new separation of R_i , Let's denote $k = \lceil \frac{i}{2} \rceil$.



178

$$\begin{aligned}
R_i &= \sum_{j=1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
&= \sum_{j=1}^{k-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
&\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right) \\
(3.27) \quad &\quad + \sum_{j=k+1}^{2i-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\
&\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1,2i} + T_{i-1,2i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2i} \right) \\
&\quad + \sum_{j=2i+1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
&= I_1 + I_2 + I_3 + I_4 + I_5
\end{aligned}$$

180

181 **LEMMA 3.11.** *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for*

182 $3 \leq i \leq N, k = \lceil \frac{i}{2} \rceil$

$$183 \quad (3.28) \quad |I_1| = \left| \sum_{j=1}^{k-1} S_{ij} \right| \leq \begin{cases} Ch^2 x_i^{-\alpha/2-2/r}, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 x_i^{-1-\alpha} \ln(i), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r} x_i^{-1-\alpha}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

184 *Proof.* by Lemma A.3 , Lemma B.6

$$185 \quad (3.29) \quad S_{i1} \leq C x_1^{\alpha/2} x_1 x_i^{-1-\alpha} = C x_1^{\alpha/2+1} x_i^{-1-\alpha} = C T^{\alpha/2+1} h^{r\alpha/2+r} x_i^{-1-\alpha}$$

186 For $2 \leq j \leq k-1$, by Lemma B.4 and Lemma B.6

$$\begin{aligned} S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) D_h^2 K_y(x_i) dy \\ 187 \quad (3.30) \quad &\leq Ch^2 \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)} dy \\ &= Ch^2 x_i^{-1-\alpha} \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} dy \end{aligned}$$

188 Therefore,

$$\begin{aligned} I_1 &= \sum_{j=1}^{k-1} S_{ij} = S_{i1} + \sum_{j=2}^{k-1} S_{ij} \\ 189 \quad (3.31) \quad &\leq Ch^{r\alpha/2+r} x_i^{-1-\alpha} + Ch^2 x_i^{-1-\alpha} \int_{x_1}^{x_{\lceil \frac{i}{2} \rceil - 1}} y^{\alpha/2-2/r} dy \\ &\leq Ch^{r\alpha/2+r} x_i^{-1-\alpha} + Ch^2 x_i^{-1-\alpha} \int_{x_1}^{2^{-r} x_i} y^{\alpha/2-2/r} dy \end{aligned}$$

190 But

$$191 \quad (3.32) \quad \int_{x_1}^{2^{-r} x_i} y^{\alpha/2-2/r} dy \leq \begin{cases} \frac{1}{\alpha/2-2/r+1} (2^{-r} x_i)^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 > 0 \\ \ln(2^{-r} x_i) - \ln(x_1), & \alpha/2 - 2/r + 1 = 0 \\ \frac{1}{|\alpha/2-2/r+1|} x_1^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

192 So we have

$$193 \quad (3.33) \quad I_1 \leq \begin{cases} \frac{C}{\alpha/2-2/r+1} h^2 x_i^{-\alpha/2-2/r}, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 x_i^{-1-\alpha} \ln(i), & \alpha/2 - 2/r + 1 = 0 \\ \frac{C}{|\alpha/2-2/r+1|} h^{r\alpha/2+r} x_i^{-1-\alpha}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \quad \square$$

194 **DEFINITION 3.12.** For convience, let's denote

$$195 \quad (3.34) \quad V_{ij} = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right)$$

196

197 **THEOREM 3.13.** There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for

198 $3 \leq i < N/2, k = \lceil \frac{i}{2} \rceil$,

$$199 \quad (3.35) \quad I_3 = \sum_{j=k+1}^{2i-1} V_{ij} \leq Ch^2 x_i^{-\alpha/2-2/r}$$

To estimate V_{ij} , we need some preparations.

LEMMA 3.14. For $y \in (x_{j-1}, x_j)$, we can rewrite

$$(3.36) \quad y = x_{j-1} + \theta h_j = (1 - \theta)x_{j-1} + \theta x_j =: y_j^\theta, \quad \theta \in (0, 1)$$

by Lemma A.2,

$$(3.37) \quad \begin{aligned} T_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\ &= \int_0^1 (u(y_j^\theta) - \Pi_h u(y_j^\theta)) \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} h_j d\theta \\ &= \int_0^1 -\frac{\theta(1-\theta)}{2} h_j^3 u''(y_j^\theta) \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} \\ &\quad + \frac{\theta(1-\theta)}{3!} h_j^4 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} (\theta^2 u'''(\eta_{j1}^\theta) - (1-\theta)^2 u'''(\eta_{j2}^\theta)) d\theta \end{aligned}$$

where $\eta_{j1}^\theta \in (x_{j-1}, y_j^\theta)$, $\eta_{j2}^\theta \in (y_j^\theta, x_j)$.

Now Let's construct a series of functions to represent T_{ij} .

DEFINITION 3.15.

$$(3.38) \quad y_{j-i}(x) = (x^{1/r} + Z_{j-i})^r, \quad Z_{j-i} = T^{1/r} \frac{j-i}{N}$$

Particularly, for $i, j \leq N-1$,

$$(3.39) \quad y_{j-i}(x_{i-1}) = x_{j-1}, \quad y_{j-i}(x_i) = x_j, \quad y_{j-i}(x_{i+1}) = x_{j+1}$$

$$(3.40) \quad y_{j-i}'(x) = y_{j-i}(x)^{1-1/r} x^{1/r-1}$$

$$(3.41) \quad y_{j-i}''(x) = \frac{1-r}{r} y_{j-i}(x)^{1-2/r} x^{1/r-2} Z_{j-i}$$

$$(3.42) \quad y_{j-i}^\theta(x) = (1-\theta)y_{j-1-i}(x) + \theta y_{j-i}(x)$$

$$(3.43) \quad h_{j-i}(x) = y_{j-i}(x) - y_{j-i-1}(x)$$

Now, we define

$$(3.44) \quad P_{j-i}^\theta(x) = (h_{j-i}(x))^3 u''(y_{j-i}^\theta(x)) \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

$$(3.45) \quad Q_{j-i}^\theta(x) = (h_{j-i}(x))^4 \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

And now we can rewrite T_{ij}

LEMMA 3.16. For $2 \leq i \leq N, 2 \leq j \leq N$,

$$(3.46) \quad \begin{aligned} T_{ij} &= \int_0^1 -\frac{\theta(1-\theta)}{2} P_{j-i}^\theta(x_i) d\theta \\ &+ \int_0^1 \frac{\theta(1-\theta)}{3!} Q_{j-i}^\theta(x_i) [\theta^2 u'''(\eta_{j,1}^\theta) - (1-\theta)^2 u'''(\eta_{j,2}^\theta)] d\theta \end{aligned}$$

Immediately, we can see from (3.34) that

LEMMA 3.17. For $3 \leq i, j \leq N-1$,
(3.47)

$$\begin{aligned} V_{ij} &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\ &= \int_0^1 -\frac{\theta(1-\theta)}{2} D_h^2 P_{j-i}^\theta(x_i) d\theta \\ &+ \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1}) u'''(\eta_{j+1,1}^\theta) - Q_{j-i}^\theta(x_i) u'''(\eta_{j,1}^\theta)}{h_{i+1}} \right) d\theta \\ &- \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i) u'''(\eta_{j,1}^\theta) - Q_{j-i}^\theta(x_{i-1}) u'''(\eta_{j-1,1}^\theta)}{h_i} \right) d\theta \\ &- \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1}) u'''(\eta_{j+1,2}^\theta) - Q_{j-i}^\theta(x_i) u'''(\eta_{j,2}^\theta)}{h_{i+1}} \right) d\theta \\ &+ \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i) u'''(\eta_{j,2}^\theta) - Q_{j-i}^\theta(x_{i-1}) u'''(\eta_{j-1,2}^\theta)}{h_i} \right) d\theta \end{aligned}$$

To estimate V_{ij} , we first estimate $D_h^2 P_{j-i}^\theta(x_i)$, but By Lemma A.1,

$$(3.48) \quad D_h^2 P_{j-i}^\theta(x_i) = P_{j-i}^{\theta''}(\xi), \quad \xi \in (x_{i-1}, x_{i+1})$$

By Leibniz formula, we calculate and estimate the derivations of $h_{j-i}^3(x)$, $u''(y_{j-i}^\theta(x))$

and $\frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$ separately.

Firstly, we have

LEMMA 3.18. There exists a constant $C = C(T, r)$ such that For $3 \leq i \leq N-1, \lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\}$, $\xi \in (x_{i-1}, x_{i+1})$,

$$(3.49) \quad h_{j-i}^3(\xi) \leq C h^2 x_i^{2-2/r} h_j$$

$$(3.50) \quad (h_{j-i}^3(\xi))' \leq C(r-1) h^2 x_i^{1-2/r} h_j$$

$$(3.51) \quad (h_{j-i}^3(\xi))'' \leq C(r-1) h^2 x_i^{-2/r} h_j$$

The proof of this theorem see Lemma B.9 and Lemma B.10

Second,

LEMMA 3.19. There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $3 \leq i \leq N-1, \lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\}$, $\xi \in (x_{i-1}, x_{i+1})$,

$$(3.52) \quad u''(y_{j-i}^\theta(\xi)) \leq C x_i^{\alpha/2-2}$$

$$(3.53) \quad (u''(y_{j-i}^\theta(\xi)))' \leq C x_i^{\alpha/2-3}$$

$$(3.54) \quad (u''(y_{j-i}^\theta(\xi)))'' \leq C x_i^{\alpha/2-4}$$

The proof of this theorem see Proof 30

And Finally, we have

LEMMA 3.20. *There exists a constant $C = C(T, \alpha, r)$ such that For $3 \leq i \leq N - 1, \lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\}, \xi \in (x_{i-1}, x_{i+1}),$*

$$(3.55) \quad |y_{j-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_j^\theta - x_i|^{1-\alpha}$$

$$(3.56) \quad |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'| \leq C|y_j^\theta - x_i|^{1-\alpha} x_i^{-1}$$

$$(3.57) \quad |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})''| \leq C|y_j^\theta - x_i|^{1-\alpha} x_i^{-2}$$

where $y_j^\theta = \theta x_{j-1} + (1 - \theta)x_j$

The proof of this theorem see Proof 31

LEMMA 3.21. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $3 \leq i \leq N - 1, \lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i - 1, N - 1\},$*

$$(3.58) \quad D_h^2 P_{j-i}^\theta(x_i) \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j$$

where $y_j^\theta = \theta x_{j-1} + (1 - \theta)x_j$

Proof. Since Lemma A.1

$$(3.59) \quad D_h^2 P_{j-i}^\theta(x_i) = P_{j-i}^{\theta''}(\xi), \quad \xi \in (x_{i-1}, x_{i+1})$$

From (3.44), using Leibniz formula and Lemma 3.18, Lemma 3.19 and Lemma 3.20 \square

LEMMA 3.22. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $3 \leq i \leq N - 1.$
For $\lceil \frac{i}{2} \rceil \leq j \leq \min\{2i - 1, N - 1\},$*

$$(3.60) \quad \begin{aligned} & \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \right) \\ & \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j \end{aligned}$$

And for $\lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i, N\},$

$$(3.61) \quad \begin{aligned} & \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta) - Q_{j-i}^\theta(x_{i-1})u'''(\eta_{j-1}^\theta)}{h_i} \right) \\ & \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j \end{aligned}$$

where $\eta_j^\theta \in (x_{j-1}, x_j).$

proof see Proof 32

272 **LEMMA 3.23.** *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for*
 273 $3 \leq i \leq N-1, \lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i-1, N-1\},$

$$\begin{aligned} V_{ij} &\leq Ch^2 \int_0^1 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j d\theta \\ &= Ch^2 \int_{x_{j-1}}^{x_j} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} dy \end{aligned}$$

274 (3.62)

275 *Proof.* Since Lemma 3.17, by Lemma 3.21 and Lemma 3.22, we get the result
 276 immediately. \square

277 Now we can prove Theorem 3.13 using Lemma 3.23, $k = \lceil \frac{i}{2} \rceil$

$$\begin{aligned} I_3 &= \sum_{k+1}^{2i-1} V_{ij} \leq Ch^2 \int_{x_k}^{x_{2i-1}} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} dy \\ &= Ch^2 \left(\frac{|x_k - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} + \frac{|x_{2i-1} - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} \right) x_i^{\alpha/2-2-2/r} \\ &\leq Ch^2 x_i^{2-\alpha} x_i^{\alpha/2-2-2/r} = Ch^2 x_i^{-\alpha/2-2/r} \end{aligned}$$

278 (3.63)

279 Now we study I_2, I_4 .

280 **LEMMA 3.24.** *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for*
 281 $3 \leq i \leq N-1, k = \lceil \frac{i}{2} \rceil,$
 282 (3.64)

$$I_2 = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right) \leq Ch^2 x_i^{-\alpha/2-2/r}$$

283 And for $3 \leq i < N/2,$

$$I_4 = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1,2i} + T_{i-1,2i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2i} \right) \leq Ch^2 x_i^{-\alpha/2-2/r}$$

284 (3.65)

285 *Proof.* In fact,

$$\begin{aligned} &\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \\ &= \frac{1}{h_{i+1}} (T_{i+1,k} - T_{i,k}) + \frac{1}{h_{i+1}} (T_{i+1,k+1} - T_{i,k}) + \left(\frac{1}{h_{i+1}} - \frac{1}{h_i} \right) T_{i,k} \end{aligned}$$

286 (3.66)

287 While, by Lemma A.2 and Lemma B.1

$$\begin{aligned} \frac{1}{h_{i+1}} (T_{i+1,k} - T_{i,k}) &= \int_{x_{k-1}}^{x_k} (u(y) - \Pi_h u(y)) \frac{|x_{i+1} - y|^{1-\alpha} - |x_i - y|^{1-\alpha}}{h_{i+1} \Gamma(2-\alpha)} dy \\ &\leq h_k^2 \max_{\eta \in (x_{k-1}, x_k)} |u''(\eta)| \int_{x_{k-1}}^{x_k} \frac{|\xi - y|^{-\alpha}}{\Gamma(1-\alpha)} dy, \quad \xi \in (x_i, x_{i+1}) \\ &\leq Ch^2 x_k^{2-2/r} x_{k-1}^{\alpha/2-2} h_k |x_i - x_k|^{-\alpha} \\ &\leq Ch^2 x_i^{-\alpha/2-2/r} h_k \end{aligned}$$

288 (3.67)

289 Thus,

$$290 \quad (3.68) \quad \frac{2}{h_i + h_{i+1}} \frac{1}{h_{i+1}} |T_{i+1,k} - T_{i,k}| \leq Ch^2 x_i^{-\alpha/2-2/r}$$

291 From Lemma 3.16

$$292 \quad (3.69) \quad \begin{aligned} \frac{1}{h_{i+1}} (T_{i+1,k+1} - T_{i,k}) &= \int_0^1 -\frac{\theta(1-\theta)}{2} \frac{P_{k-i}^\theta(x_{i+1}) - P_{k-i}^\theta(x_i)}{h_{i+1}} d\theta \\ &+ \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{Q_{k-i}^\theta(x_{i+1})u'''(\eta_{k+1,1}^\theta) - Q_{k-i}^\theta(x_i)u'''(\eta_{k,1}^\theta)}{h_{i+1}} d\theta \\ &- \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{Q_{k-i}^\theta(x_{i+1})u'''(\eta_{k+1,2}^\theta) - Q_{k-i}^\theta(x_i)u'''(\eta_{k,2}^\theta)}{h_{i+1}} d\theta \end{aligned}$$

293 and

$$294 \quad (3.70) \quad D_h P_{k-i}^\theta(x_i) := \frac{P_{k-i}^\theta(x_{i+1}) - P_{k-i}^\theta(x_i)}{h_{i+1}} = P_{k-i}^{\theta'}(\xi), \quad \xi \in (x_i, x_{i+1})$$

295 Similar with Lemma 3.21, from Lemma 3.16, using Leibniz formula, by Lemma B.9,
296 Lemma 3.19 and Lemma 3.20 we get

$$297 \quad (3.71) \quad |D_h P_{k-i}^\theta(x_i)| \leq Ch^2 x_i^{-\alpha/2-2/r} h_k$$

298 And with Lemma 3.22, we can get

$$299 \quad (3.72) \quad \frac{2}{h_i + h_{i+1}} \frac{1}{h_{i+1}} |T_{i+1,k+1} - T_{i,k}| \leq Ch^2 x_i^{-\alpha/2-2/r}$$

300 For the third term, by Lemma B.1, Lemma B.2 and Lemma A.2

$$301 \quad (3.73) \quad \begin{aligned} \frac{2}{h_i + h_{i+1}} \frac{h_{i+1} - h_i}{h_i h_{i+1}} T_{i,k} &\leq h_i^{-3} h^2 x_i^{1-2/r} h_k Ch_k^2 x_{k-1}^{\alpha/2-2} |x_k - x_i|^{1-\alpha} \\ &\leq Ch^2 x_i^{-\alpha/2-2/r} \end{aligned}$$

302 Summarizes, we have

$$303 \quad (3.74) \quad I_2 \leq Ch^2 x_i^{-\alpha/2-2/r}$$

304 The case for I_4 is similar. □

305 Now combine Lemma 3.10, Lemma 3.11, Lemma 3.24, Theorem 3.13, Lemma 3.8
306 and Lemma 3.9, we get Theorem 3.2.

3.4. Proof of Theorem 3.3. For $N/2 \leq i < N, k = \lceil \frac{i}{2} \rceil$, we have

$$\begin{aligned}
 (3.75) \quad R_i &= \sum_{j=1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &= \sum_{j=1}^{k-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right) \\
 &\quad + \sum_{j=k+1}^{N-1} + \sum_{j=N}^{N+1} + \sum_{j=N+2}^{2N-\lceil \frac{N}{2} \rceil} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\
 &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1,2N-\lceil \frac{N}{2} \rceil+1} + T_{i-1,2N-\lceil \frac{N}{2} \rceil}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2N-\lceil \frac{N}{2} \rceil+1} \right) \\
 &\quad + \sum_{j=2N-\lceil \frac{N}{2} \rceil+2}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &= I_1 + I_2 + I_3^1 + I_3^2 + I_3^3 + I_4 + I_5
 \end{aligned}$$

We have estimate I_1 in Lemma 3.11 and I_2 in Lemma 3.24. We can control I_3^1 similar with Theorem 3.13 by Lemma 3.23 where $2i - 1 \geq N - 1$

LEMMA 3.25. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $N/2 \leq i < N, k = \lceil \frac{i}{2} \rceil$,*

$$\begin{aligned}
 (3.76) \quad I_3^1 &= \sum_{j=k+1}^{N-1} V_{ij} \leq Ch^2 \int_{x_k}^{x_{N-1}} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} dy \\
 &= Ch^2 \left(\frac{|x_k - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} + \frac{|x_{N-1} - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} \right) x_i^{\alpha/2-2-2/r} \\
 &\leq Ch^2 x_i^{2-\alpha} x_i^{\alpha/2-2-2/r} = Ch^2 x_i^{-\alpha/2-2/r}
 \end{aligned}$$

Let's study I_3^3 before I_3^2 .

$$(3.77) \quad I_3^3 = \sum_{j=N+2}^{2N-\lceil \frac{N}{2} \rceil} V_{ij}$$

Similarly, Let's define a new series of functions

DEFINITION 3.26. *For $i \leq N - 1, j \geq N + 1$, with no confusion, we also denote in this section*

$$(3.78) \quad y_{j-i}(x) = 2T - (Z_{2N-j+i} - x^{1/r})^r, \quad Z_{2N-j+i} = T^{1/r} \frac{2N-j+i}{N}$$

Particularly

$$y_{j-i}(x_{i-1}) = x_{j-1}, \quad y_{j-i}(x_i) = x_j, \quad y_{j-i}(x_{i+1}) = x_{j+1}$$

322 $y \rightarrow z?$

323 (3.79) $y_{j-i}'(x) = (2T - y_{j-i}(x))^{1-1/r} x^{1/r-1}$

324 (3.80) $y_{j-i}''(x) = \frac{1-r}{r} (2T - y_{j-i}(x))^{1-2/r} x^{1/r-2} Z_{2N-j+i}$

325 (3.81)

326

327 (3.82) $y_{j-i}^\theta(x) = (1 - \theta)y_{j-i-1}(x) + \theta y_{j-i}(x)$

328

329 (3.83) $h_{j-i}(x) = y_{j-i}(x) - y_{j-i-1}(x)$

330

331 (3.84) $P_{j-i}^\theta(x) = (h_{j-i}(x))^3 u''(y_{j-i}^\theta(x)) \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$

332

333 (3.85) $Q_{j-i}^\theta(x) = (h_{j-i}(x))^4 \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$

334 Now we have the same formula Lemma 3.17 for $i \leq N-1, j \geq N+2$,

335 Similarly, we first estimate

336 (3.86) $D_h^2 P_{j-i}^\theta(\xi) = P_{j-i}^{\theta''}(\xi), \quad \xi \in (x_{i-1}, x_{i+1})$

337 Combine Definition 3.26, Lemma B.11, Lemma B.12 and Lemma B.13, using
338 Leibniz formula, we have

339 LEMMA 3.27. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For*
340 *$N/2 \leq i \leq N-1, N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1$, we have*

341 (3.87)
$$\begin{aligned} |D_h^2 P_{j-i}^\theta(\xi)| &\leq Ch_j h^2 \left(|y_j^\theta - x_i|^{1-\alpha} \right. \\ &\quad + |y_j^\theta - x_i|^{-\alpha} (|2T - x_i - y_j^\theta| + h_N) \\ &\quad + |y_j^\theta - x_i|^{-1-\alpha} (|2T - x_i - y_j^\theta| + h_N)^2 \\ &\quad \left. + (r-1) |y_j^\theta - x_i|^{-\alpha} \right) \end{aligned}$$

342 And

343 LEMMA 3.28. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For*
344 *$N/2 \leq i \leq N-1, N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil, \xi \in (x_{i-1}, x_{i+1})$, we have*

345 (3.88)
$$\begin{aligned} \frac{2}{h_i + h_{i+1}} \left| \frac{Q_{j-i}^\theta(x_{i+1}) u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i) u'''(\eta_j^\theta)}{h_{i+1}} \right| \\ \leq Ch^2 h_j (|y_j^\theta - x_i|^{1-\alpha} + |y_j^\theta - x_i|^{-\alpha} (|2T - x_i - y_j^\theta| + h_N)) \end{aligned}$$

346 and

347 (3.89)
$$\begin{aligned} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i) u'''(\eta_j^\theta) - Q_{j-i}^\theta(x_{i-1}) u'''(\eta_{j-1}^\theta)}{h_{i+1}} \right) \\ \leq Ch^2 h_j (|y_j^\theta - x_i|^{1-\alpha} + |y_j^\theta - x_i|^{-\alpha} (|2T - x_i - y_j^\theta| + h_N)) \end{aligned}$$

Proof. From Definition 3.26, by Lemma B.11 and Lemma B.13, for $\xi \in (x_i, x_{i+1})$, by Leibniz formula, we have

$$(3.90) \quad |Q_{j-i}^\theta(\xi)| \leq Ch^2 h_j^2 ((r-1)|y_j^\theta - x_i|^{1-\alpha} + |y_j^\theta - x_i|^{-\alpha}(|2T - x_i - y_j^\theta| + h_N))$$

$$(3.91) \quad |Q_{j-i}^\theta(\xi)| \leq Ch^2 h_j^2 |y_j^\theta - x_i|^{1-\alpha}$$

So use the skill in Proof 32 with Lemma B.12

$$(3.92) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \right) \leq Ch^2 h_j (|y_j^\theta - x_i|^{1-\alpha} + |y_j^\theta - x_i|^{-\alpha}(|2T - x_i - y_j^\theta| + h_N)) \quad \square$$

Combine Lemma 3.27, Lemma 3.28 and formula Lemma 3.17 for $i \leq N-1, j \geq N+2$, we have

LEMMA 3.29. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq i \leq N-1, N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1$*

$$(3.93) \quad V_{ij} \leq Ch^2 \int_{x_{j-1}}^{x_j} \left(|y - x_i|^{1-\alpha} + |y - x_i|^{-\alpha}(|2T - x_i - y| + h_N) + |y - x_i|^{-1-\alpha}(|2T - x_i - y| + h_N)^2 + (r-1)|y - x_i|^{-\alpha} \right) dy$$

We can estimate I_3^3 Now.

LEMMA 3.30. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq i \leq N-1$, we have*

$$(3.94) \quad I_3^3 = \sum_{j=N+2}^{2N - \lceil \frac{N}{2} \rceil} V_{ij} \leq Ch^2 + C(r-1)h^2 |T - x_{i-1}|^{1-\alpha}$$

Proof.

$$(3.95) \quad I_3^3 = \sum_{j=N+2}^{2N - \lceil \frac{N}{2} \rceil} V_{ij} \leq Ch^2 \int_{x_{N+1}}^{x_{2N - \lceil \frac{N}{2} \rceil}} \left(|y - x_i|^{1-\alpha} + |y - x_i|^{-\alpha}(|2T - x_i - y| + h_N) + |y - x_i|^{-1-\alpha}(|2T - x_i - y| + h_N)^2 + (r-1)|y - x_i|^{-\alpha} \right) dy$$

Since

$$(3.96) \quad |2T - x_i - y| + h_N \leq y - x_i$$

$$(3.97) \quad \begin{aligned} I_3^3 &\leq Ch^2 \int_{x_{N+1}}^{x_{2N - \lceil \frac{N}{2} \rceil}} |y - x_i|^{1-\alpha} + (r-1)|y - x_i|^{-\alpha} \\ &\leq Ch^2 (T^{2-\alpha} + (r-1)|x_{N+1} - x_i|^{1-\alpha}) \\ &\leq Ch^2 + C(r-1)h^2 |T - x_{i-1}|^{1-\alpha} \end{aligned} \quad \square$$

For I_3^2 , we have

THEOREM 3.31. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that, for $N/2 \leq i \leq N-1$*

$$(3.98) \quad \begin{aligned} V_{iN} &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1, N+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i, N} + \frac{1}{h_i} T_{i-1, N-1} \right) \\ &\leq Ch^2 + C(r-1)h^2 |T - x_{i-1}|^{1-\alpha} \end{aligned}$$

Proof. We use the similar skill in the last section, but more complicated. for $j = N$, Let

$$(3.99) \quad {}_L y_{N-1-i}(x) = (x^{1/r} + Z_{N-1-i})^r, \quad Z_{N-1-i} = T^{1/r} \frac{N-1-i}{N}$$

$$(3.100) \quad {}_0 y_{N-i}(x) = \frac{x^{1/r} - Z_i}{Z_1} h_N + T, \quad Z_i = T^{1/r} \frac{i}{N}, x_N = T$$

and

$$(3.101) \quad {}_R y_{N+1-i}(x) = 2T - (Z_{N-1+i} - x^{1/r})^r, \quad Z_{N-1+i} = T^{1/r} \frac{N-1+i}{N}$$

Thus,

$${}_L y_{N-1-i}(x_{i-1}) = x_{N-2}, \quad {}_L y_{N-1-i}(x_i) = x_{N-1}, \quad {}_L y_{N-1-i}(x_{i+1}) = x_N$$

$${}_0 y_{N-i}(x_{i-1}) = x_{N-1}, \quad {}_0 y_{N-i}(x_i) = x_N, \quad {}_0 y_{N-i}(x_{i+1}) = x_{N+1}$$

$${}_R y_{N+1-i}(x_{i-1}) = x_N, \quad {}_R y_{N+1-i}(x_i) = x_{N+1}, \quad {}_R y_{N+1-i}(x_{i+1}) = x_{N+2}$$

Then, define

$$(3.102) \quad {}_L y_{N-i}^\theta(x) = \theta {}_L y_{N-1-i}(x) + (1-\theta) {}_0 y_{N-i}(x)$$

$$(3.103) \quad {}_R y_{N+1-i}^\theta(x) = \theta {}_0 y_{N-i}(x) + (1-\theta) {}_R y_{N+1-i}(x)$$

$$(3.104) \quad {}_L h_{N-i}(x) = {}_0 y_{N-i}(x) - {}_L y_{N-1-i}(x)$$

$$(3.105) \quad {}_R h_{N+1-i}(x) = {}_R y_{N+1-i}(x) - {}_0 y_{N-i}(x)$$

We have

$$(3.106) \quad {}_L y_{N-1-i}'(x) = {}_L y_{N-1-i}^{1-1/r}(x) x^{1/r-1}$$

$$(3.107) \quad {}_L y_{N-1-i}''(x) = \frac{1-r}{r} {}_L y_{N-1-i}^{1-2/r}(x) x^{1/r-2} Z_{N-1-i}$$

$$(3.108) \quad {}_0 y_{N-i}'(x) = \frac{1}{r} \frac{h_N}{Z_1} x^{1/r-1}$$

$$(3.109) \quad {}_0 y_{N-i}''(x) = \frac{1-r}{r^2} \frac{h_N}{Z_1} x^{1/r-2}$$

$$(3.110) \quad {}_R y_{N+1-i}'(x) = (2T - {}_R y_{N+1-i}(x))^{1-1/r} x^{1/r-1}$$

$$(3.111) \quad {}_R y_{N+1-i}''(x) = \frac{1-r}{r} (2T - {}_R y_{N+1-i}(x))^{1-2/r} x^{1/r-2} Z_{N-1+i}$$

397

$$398 \quad (3.112) \quad {}_L P_{N-i}^\theta(x) = ({}_L h_{N-i}(x))^3 \frac{|{}_L y_{N-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)} u''({}_L y_{N-i}^\theta(x))$$

$$399 \quad (3.113) \quad {}_R P_{N+1-i}^\theta(x) = ({}_R h_{N+1-i}(x))^3 \frac{|{}_R y_{N+1-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)} u''({}_R y_{N+1-i}^\theta(x))$$

$$400 \quad (3.114) \quad {}_L Q_{N-i}^\theta(x) = ({}_L h_{N-i}(x))^4 \frac{|{}_L y_{N-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

$$401 \quad (3.115) \quad {}_R Q_{N+1-i}^\theta(x) = ({}_R h_{N+1-i}(x))^4 \frac{|{}_R y_{N+1-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

 402 Similar with Lemma 3.16, we can get for $l = -1, 0, 1$,

$$403 \quad (3.116) \quad \begin{aligned} T_{i+l, N+l} &= \int_0^1 -\frac{\theta(1-\theta)}{2} {}_L P_{N-i}^\theta(x_{i+l}) d\theta \\ &+ \int_0^1 \frac{\theta(1-\theta)}{3!} {}_L Q_{N-i}^\theta(x_{i+l}) (\theta^2 u'''(\eta_{N+l,1}^\theta) - (1-\theta)^2 u'''(\eta_{N+l,2}^\theta)) d\theta \end{aligned}$$

404

$$(3.117) \quad \begin{aligned} T_{i+l, N+1+l} &= \int_0^1 -\frac{\theta(1-\theta)}{2} {}_R P_{N+1-i}^\theta(x_{i+l}) d\theta \\ &+ \int_0^1 \frac{\theta(1-\theta)}{3!} {}_R Q_{N+1-i}^\theta(x_{i+l}) (\theta^2 u'''(\eta_{N+1+l,1}^\theta) - (1-\theta)^2 u'''(\eta_{N+1+l,2}^\theta)) d\theta \end{aligned}$$

406 So we have

$$(3.118) \quad \begin{aligned} V_{i,N} &= \int_0^1 -\frac{\theta(1-\theta)}{2} D_{hL}^2 {}_L P_{N-i}^\theta(x_i) d\theta \\ &+ \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_{i+1}) u'''(\eta_{N+1,1}^\theta) - {}_L Q_{N-i}^\theta(x_i) u'''(\eta_{N,1}^\theta)}{h_{i+1}} \right) d\theta \\ &- \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_i) u'''(\eta_{N,1}^\theta) - {}_L Q_{N-i}^\theta(x_{i-1}) u'''(\eta_{N-1,1}^\theta)}{h_i} \right) d\theta \\ &- \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_{i+1}) u'''(\eta_{N+1,2}^\theta) - {}_L Q_{N-i}^\theta(x_i) u'''(\eta_{N,2}^\theta)}{h_{i+1}} \right) d\theta \\ &+ \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_i) u'''(\eta_{N,2}^\theta) - {}_L Q_{N-i}^\theta(x_{i-1}) u'''(\eta_{N-1,2}^\theta)}{h_i} \right) d\theta \end{aligned}$$

 408 $N+1$ is similar.

 409 We estimate $D_{hL}^2 {}_L P_{N-i}^\theta(x_i) = {}_L P_{N-i}^{\theta''}(\xi), \xi \in (x_{i-1}, x_{i+1})$,

410

LEMMA 3.32.

$$411 \quad (3.119) \quad {}_L h_{N-i}^3(\xi) \leq C h_N^3 \leq C h^3$$

$$412 \quad (3.120) \quad {}_R h_{N+1-i}^3(\xi) \leq C h_N^3 \leq C h^3$$

$$(3.121) \quad ({}_L h_{N-i}^3(\xi))' \leq C(r-1)h_N^2 h \leq C(r-1)h^3$$

$$(3.122) \quad ({}_R h_{N+1-i}^3(\xi))' \leq C(r-1)h_N^2 h \leq C(r-1)h^3$$

$$(3.123) \quad ({}_L h_{N-i}^3(\xi))'' \leq C(r-1)h^2$$

$$(3.124) \quad ({}_R h_{N+1-i}^3(\xi))'' \leq C(r-1)h^2$$

Proof.

$$(3.125) \quad {}_L h_{N-i}(\xi) \leq 2(C?)h_N, \quad {}_R h_{N+1-i}(\xi) \leq 2h_N$$

418

$$(3.126) \quad \begin{aligned} ({}_L h_{N-i}^l(\xi))' &= {}_L h_{N-i}^{l-1}(\xi)({}_0 y_{N-i}'(\xi) - {}_L y_{N-1-i}'(\xi)) \\ &= {}_L h_{N-i}^{l-1}(\xi)\xi^{1/r-1}\left(\frac{1}{r}\frac{h_N}{Z_1} - {}_L y_{N-1-i}^{1-1/r}(\xi)\right) \end{aligned}$$

420 while

(3.127)

$$\begin{aligned} \left|\frac{1}{r}\frac{h_N}{Z_1} - {}_L y_{N-1-i}^{1-1/r}(\xi)\right| &= \left|\frac{1}{r}\frac{x_N - (x_N^{1/r} - Z_1)^r}{Z_1} - \eta^{1-1/r}\right| \quad \eta \in [x_{N-2}, x_N] \\ &= T^{1-1/r} \left| \left(\frac{N-t}{N}\right)^{r-1} - \left(\frac{N-s}{N}\right)^{r-1} \right| \quad t \in [0, 1], s \in [0, 2] \\ &\leq T^{1-1/r} \left| 1 - \left(\frac{N-2}{N}\right)^{r-1} \right| \leq CT^{1-1/r}(r-1)\frac{2}{N} \end{aligned}$$

422 Thus,

$$(3.128) \quad ({}_L h_{N-i}^l(\xi))' \leq C(r-1)h_N^{l-1}x_i^{1/r-1}h$$

424 And

(3.129)

$$\begin{aligned} ({}_L h_{N-i}^3(\xi))'' &= 3{}_L h_{N-i}^2(\xi){}_L h_{N-i}''(\xi) + 6{}_L h_{N-i}(\xi)({}_L h_{N-i}'(\xi))^2 \\ &\leq Ch_N^2 \frac{1-r}{r} x_i^{1/r-2} \left(\frac{1}{r}\frac{h_N}{Z_1} - {}_L y_{N-1-i}^{1-2/r}(\xi)Z_{N-1-i}\right) + Ch_N(r-1)^2 h^2 x_i^{2/r-2} \end{aligned}$$

$$(3.130) \quad \left|\frac{h_N}{rZ_1} - {}_L y_{N-1-i}^{1-2/r}(\xi)Z_{N-1-i}\right| \leq T^{1-1/r} + Cx_N^{1-2/r}x_N^{1/r} = CT^{1-1/r}$$

427 So

$$\begin{aligned} (3.130) \quad ({}_L h_{N-i}^3(\xi))'' &\leq Ch_N^2 \frac{1-r}{r} x_i^{1/r-2} + C(r-1)^2 h_N x_i^{2/r-2} h^2 \\ &\leq C(r-1)h_N^2 \end{aligned}$$

429 ${}_R h_{N+1-i}^3(\xi)$ is similar. □

LEMMA 3.33.

$$(3.131) \quad u''({}_L y_{N-i}^\theta(\xi)) \leq Cx_{N-2}^{-\alpha/2-2} \leq C$$

$$(3.132) \quad (u''({}_L y_{N-i}^\theta(\xi)))' \leq C$$

$$(3.133) \quad (u''({}_L y_{N-i}^\theta(\xi)))'' \leq C$$

Proof.

$$\begin{aligned}
 (u''(Ly_{N-i}^\theta(\xi)))' &= u'''(Ly_{N-i}^\theta(\xi))Ly_{N-i}^{\theta'}(\xi) \\
 &\leq C(\theta Ly_{N-1-i}'(\xi) + (1-\theta)_0y_{N-i}'(\xi)) \\
 &\leq Cx_i^{1/r-1}(\theta Ly_{N-1-i}^{1-1/r}(\xi) + (1-\theta)\frac{h_N}{rZ_1}) \\
 &\leq Cx_i^{1/r-1}x_N^{1-1/r}
 \end{aligned}
 \tag{3.134}$$

And

$$\begin{aligned}
 (u''(Ly_{N-i}^\theta(\xi)))'' &= u''''(Ly_{N-i}^\theta(\xi))(Ly_{N-i}^{\theta'}(\xi))^2 + u'''(Ly_{N-i}^\theta(\xi))Ly_{N-i}^{\theta''}(\xi) \\
 &\leq Cx_i^{2/r-2}x_N^{2-2/r} + C\frac{r-1}{r}x_i^{1/r-2}(\theta x_N^{1-2/r}Z_{N-1-i} + (1-\theta)\frac{h_N}{rZ_1}) \\
 &\leq Cx_i^{2/r-2} + C(r-1)x_i^{1/r-2}T^{1-1/r}
 \end{aligned}
 \tag{3.135}$$

□

LEMMA 3.34.

$$|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_N^\theta - x_i|^{1-\alpha} \tag{3.136}$$

$$(|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha})' \leq C|y_N^\theta - x_i|^{1-\alpha} \tag{3.137}$$

$$(|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha})'' \leq C(r-1)|y_N^\theta - x_i|^{-\alpha} + |y_N^\theta - x_i|^{1-\alpha} \tag{3.138}$$

Proof.

$$\begin{aligned}
 (Ly_{N-i}^\theta(\xi) - \xi)' &= (\theta(Ly_{N-1-i}(\xi) - \xi) + (1-\theta)(_0y_{N-i}(\xi) - \xi))' \\
 &= \theta(Ly_{N-1-i}'(\xi) - 1) + (1-\theta)(_0y_{N-i}'(\xi) - 1) \\
 &= \theta\xi^{1/r-1}(Ly_{N-1-i}^{1-1/r}(\xi) - \xi^{1-1/r}) + (1-\theta)\xi^{1/r-1}(\frac{h_N}{rZ_1} - \xi^{1-1/r})
 \end{aligned}
 \tag{3.139}$$

$$\begin{aligned}
 (Ly_{N-i}^\theta(\xi) - \xi)'' &= \theta(Ly_{N-1-i}''(\xi)) + (1-\theta)(_0y_{N-i}''(\xi)) \\
 &= \frac{1-r}{r}\xi^{1/r-2}(\theta Ly_{N-1-i}^{1-2/r}(\xi)Z_{N-1-i} + (1-\theta)\frac{h_N}{rZ_1}) \leq 0
 \end{aligned}
 \tag{3.140}$$

And

$$|(Ly_{N-i}^\theta(\xi) - \xi)''| \leq C(r-1)\xi^{1/r-2}T^{1-1/r} \tag{3.141}$$

We have known

$$C|x_{N-1} - x_i| \leq |Ly_{N-1-i}(\xi) - \xi| \leq C|x_{N-1} - x_i| \tag{3.142}$$

If $\xi \leq x_{N-1}$, then $(_0y_{N-i}(\xi) - \xi)' \geq 0$, so

$$C|x_N - x_i| \leq |x_{N-1} - x_{i-1}| \leq |Ly_{N-i}^\theta(\xi) - \xi| \leq |x_{N+1} - x_{i+1}| \leq C|x_N - x_i| \tag{3.143}$$

If $i = N-1$ and $\xi \in [x_{N-1}, x_N]$, then $_0y_{N-i}(\xi) - \xi$ is concave, bigger than its two neighboring points, which are equal to h_N , so

$$h_N = |x_N - x_{N-1}| \leq |_0y_{N-i}(\xi) - \xi| \leq |x_{N+1} - x_{N-1}| = 2h_N \tag{3.144}$$

So we have

$$(3.145) \quad |Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_N^\theta - x_i|^{1-\alpha}$$

While

$$(3.146) \quad Ly_{N-1-i}^{1-1/r}(\xi) - \xi^{1-1/r} \leq (Ly_{N-1-i}(\xi) - \xi)\xi^{-1/r}$$

and

$$(3.147) \quad \left| \frac{h_N}{rZ_1} - \xi^{1-1/r} \right| \leq \max\left\{ \left| \frac{h_N}{rZ_1} - x_{i-1}^{1-1/r} \right|, \left| \frac{h_N}{rZ_1} - x_{i+1}^{1-1/r} \right| \right\}$$

$$\leq \max \begin{cases} T^{1-1/r} - x_{i-1}^{1-1/r} \leq |x_N - x_{i-1}|T^{-1/r} \leq C|x_N - x_i| \\ |x_{i+1}^{1-1/r} - x_{N-1}^{1-1/r}| \leq |x_{i+1} - x_{N-1}|x_{N-1}^{-1/r} \leq C|x_N - x_i| \end{cases}$$

So we have

$$(3.148) \quad (Ly_{N-i}^\theta(\xi) - \xi)' \leq C|y_N^\theta - x_i|$$

$$(3.149) \quad (|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha})' = |Ly_{N-i}^\theta(\xi) - \xi|^{-\alpha} (Ly_{N-i}^\theta(\xi) - \xi)' \leq |y_N^\theta - x_i|^{1-\alpha}$$

Finally,

$$(3.150) \quad \begin{aligned} (|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha})'' &= (1-\alpha)|Ly_{N-i}^\theta(\xi) - \xi|^{-\alpha} (Ly_{N-i}^\theta(\xi) - \xi)'' \\ &\quad + \alpha(\alpha-1)|Ly_{N-i}^\theta(\xi) - \xi|^{-1-\alpha} ((Ly_{N-i}^\theta(\xi) - \xi)')^2 \\ &\leq C(r-1)|y_N^\theta - x_i|^{-\alpha} + C|y_N^\theta - x_i|^{1-\alpha} \end{aligned} \quad \square$$

By the three lemmas above, for $N/2 \leq i \leq N-1$, we have

LEMMA 3.35.

$$(3.151) \quad \begin{aligned} D_{hL}^2 P_{N-i}^\theta(x_i) &= {}_L P_{N-i}^{\theta''}(\xi) \quad \xi \in (x_{i-1}, x_{i+1}) \\ &\leq Ch^3|y_N^\theta - x_i|^{1-\alpha} + C(r-1)(h^3|y_N^\theta - x_i|^{-\alpha} + h^2|y_N^\theta - x_i|^{1-\alpha}) \end{aligned}$$

while $\theta h_N = y_N^\theta - x_{N-1} \leq y_N^\theta - x_i$, we have

$$(3.152) \quad \theta D_{hL}^2 P_{N-i}^\theta(x_i) \leq Ch^3|y_N^\theta - x_i|^{1-\alpha} + C(r-1)(h^2|y_N^\theta - x_i|^{1-\alpha})$$

And

LEMMA 3.36.

$$(3.153) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_{i+1})u'''(\eta_{N+1}^\theta) - {}_L Q_{N-i}^\theta(x_i)u'''(\eta_N^\theta)}{h_{i+1}} \right) \leq Ch^3|y_N^\theta - x_i|^{1-\alpha}$$

And immediately with Lemma 3.17, For $N/2 \leq i \leq N-1$

$$(3.154) \quad \begin{aligned} V_{iN} &\leq C \int_{x_{N-1}}^{x_N} h^2|y - x_i|^{1-\alpha} + C(r-1)h|y - x_i|^{1-\alpha} dy \\ &\leq Ch^2h_N|T - x_i|^{1-\alpha} + C(r-1)h^2|x_N - x_i|^{1-\alpha} \\ &\leq Ch^2 + C(r-1)h^2|T - x_{i-1}|^{1-\alpha} \end{aligned}$$

Similarly with $j = N+1$. \square

I_4, I_5 is easy. Similar with Lemma 3.24 and Lemma 3.9, we have

THEOREM 3.37. *There is a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For*
 $N/2 \leq i \leq N,$
(3.155)

$$I_4 = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1, 2N - \lceil \frac{N}{2} \rceil + 1} + T_{i-1, 2N - \lceil \frac{N}{2} \rceil}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i, 2N - \lceil \frac{N}{2} \rceil + 1} \right) \\ \leq Ch^2$$

Proof. Similar with Lemma 3.24. In fact, let $m = 2N - \lceil \frac{N}{2} \rceil + 1$

$$(3.156) \quad \begin{aligned} & \frac{1}{h_i} (T_{i-1, l} + T_{i-1, l-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i, l} \\ &= \frac{1}{h_i} (T_{i-1, l} - T_{i, l}) + \frac{1}{h_i} (T_{i-1, l-1} - T_{i, l}) + \left(\frac{1}{h_i} - \frac{1}{h_{i+1}} \right) T_{i, l} \end{aligned}$$

While, by Lemma A.2

$$(3.157) \quad \begin{aligned} \frac{1}{h_i} (T_{i-1, l} - T_{i, l}) &= \int_{x_{l-1}}^{x_l} (u(y) - \Pi_h u(y)) \frac{|x_{i-1} - y|^{1-\alpha} - |x_i - y|^{1-\alpha}}{h_i \Gamma(2-\alpha)} dy \\ &\leq C \int_{x_{l-1}}^{x_l} h_l^2 u''(\eta) \frac{|\xi - y|^{-\alpha}}{\Gamma(1-\alpha)} dy, \quad \xi \in (x_{i-1}, x_i) \\ &\leq Ch_l^3 (2T - x_{l-1})^{\alpha/2-2} T^{-\alpha} \\ &\leq Ch_l^3 \end{aligned}$$

Thus,

$$(3.158) \quad \frac{2}{h_i + h_{i+1}} \frac{1}{h_i} (T_{i-1, l} - T_{i, l}) \leq Ch_l^2$$

For

(3.159)

$$\frac{1}{h_i} (T_{i-1, l-1} - T_{i, l}) = \int_0^1 -\frac{\theta(1-\theta)}{2} \frac{h_{l-1}^3 |y_{l-1}^\theta - x_{i-1}|^{1-\alpha} u''(\eta_{l-1}^\theta) - h_l^3 |y_l^\theta - x_i|^{1-\alpha} u''(\eta_l^\theta)}{h_i} d\theta$$

And Similar with Lemma 3.22, we can get

$$(3.160) \quad \frac{h_{l-1}^3 |y_{l-1}^\theta - x_{i-1}|^{1-\alpha} u''(\eta_{l-1}^\theta) - h_l^3 |y_l^\theta - x_i|^{1-\alpha} u''(\eta_l^\theta)}{(h_i + h_{i+1}) h_i} \leq Ch_l^2 |y_l^\theta - x_i|^{1-\alpha}$$

So

$$(3.161) \quad \frac{2}{h_i + h_{i+1}} \frac{1}{h_i} (T_{i-1, l-1} - T_{i, l}) \leq Ch^2$$

For the third term, by Lemma B.1, Lemma B.2 and Lemma A.2

$$(3.162) \quad \begin{aligned} \frac{2}{h_i + h_{i+1}} \frac{h_{i+1} - h_i}{h_i h_{i+1}} T_{i, l} &\leq h_i^{-3} h^2 x_i^{1-2/r} h_l C h_l^2 x_{l-1}^{\alpha/2-2} |x_l - x_i|^{1-\alpha} \\ &\leq Ch^2 \end{aligned}$$

Summarizes, we have

$$(3.163) \quad I_4 \leq Ch^2$$

□

And

LEMMA 3.38. *There is a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq i \leq N$,*

$$I_5 = \sum_{j=2N-\lceil \frac{N}{2} \rceil+2}^{2N} S_{ij} \leq \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

Proof. For $i \leq N, j \geq 2N - \lceil \frac{N}{2} \rceil + 2$, we have

$$\begin{aligned} S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) D_h^2 K_y(x_i) dy \\ &\leq \int_{x_{j-1}}^{x_j} Ch^2 (2T - y)^{\alpha/2-2/r} |y - x_{i+1}|^{-1-\alpha} dy \\ &\leq Ch^2 T^{-1-\alpha} \int_{x_{j-1}}^{x_j} (2T - y)^{\alpha/2-2/r} dy \end{aligned}$$

$$\begin{aligned} \sum_{j=2N-\lceil \frac{N}{2} \rceil+2}^{2N-1} S_{ij} &\leq CT^{-1-\alpha} h^2 \int_{(2-2^{-r})T}^{x_{2N-1}} (2T - y)^{\alpha/2-2/r} dy \\ &\leq CT^{-1-\alpha} h^2 \begin{cases} \frac{1}{\alpha/2-2/r+1} T^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 > 0 \\ \ln(2^{-r}T) - \ln(h_{2N}), & \alpha/2 - 2/r + 1 = 0 \\ \frac{1}{|\alpha/2-2/r+1|} h_{2N}^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \\ &= \begin{cases} \frac{C}{\alpha/2-2/r+1} T^{-\alpha/2-2/r} h^2, & \alpha/2 - 2/r + 1 > 0 \\ CrT^{-1-\alpha} h^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ \frac{C}{|\alpha/2-2/r+1|} T^{-\alpha/2-2/r} h^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \end{aligned}$$

Now we can conclude a part of the theorem Theorem 3.3 at the beginning of this section.

By Lemma 3.11 Lemma 3.24 Lemma 3.25 Theorem 3.31 Lemma 3.30 Theorem 3.37 Lemma 3.38, we have

THEOREM 3.39. *there exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $N/2 \leq i \leq N - 1$,*

$$\begin{aligned} R_i &= I_1 + I_2 + I_3^1 + I_3^2 + I_3^3 + I_4 + I_5 \\ &\leq C(r-1)h^2 |T - x_{i-1}|^{1-\alpha} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \end{aligned}$$

And what we left is the case $i = N$. Fortunately, we can use the same department of R_i above, and it is symmetric. Most of the item has been esitimated by Lemma 3.11 and Theorem 3.37, we just need to consider I_3, I_4 .

THEOREM 3.40. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that*

$$(3.167) \quad I_3 = \sum_{j=\lceil \frac{N}{2} \rceil + 1}^{N-1} V_{Nj} \leq Ch^2 + C(r-1)h^2|T - x_{N-1}|^{1-\alpha}$$

Proof. **DEFINITION 3.41.** *For $N/2 \leq j < N$, Let's define*

$$(3.168) \quad y_j(x) = \left(\frac{Z_1}{h_N}(x - x_N) + Z_j \right)^r, \quad Z_j = T^{1/r} \frac{j}{N}$$

We can see that is the inverse of the function ${}_0y_{N-i}(x)$ defined in Theorem 3.31.

$$(3.169) \quad y'_j(x) = y_j^{1-1/r}(x) \frac{rZ_1}{h_N}$$

$$(3.170) \quad y''_j(x) = y_j^{1-2/r}(x) \frac{r(r-1)Z_1}{h_N}$$

With the scheme we used several times, we can get

LEMMA 3.42. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq j < N$, $\xi \in [x_{N-1}, x_{N+1}]$,*

$$(3.171) \quad h_j(\xi)^3 \leq Ch^3$$

$$(3.172) \quad (h_j^3(\xi))' \leq C(r-1)h^3$$

$$(3.173) \quad (h_j^3(\xi))'' \leq C(r-1)h^3$$

$$(3.174) \quad u''(y_j^\theta(\xi)) \leq C$$

$$(3.175) \quad (u''(y_j^\theta(\xi)))' \leq C$$

$$(3.176) \quad (u''(y_j^\theta(\xi)))'' \leq C$$

$$(3.177) \quad |\xi - y_j^\theta(\xi)|^{1-\alpha} \leq C|x_N - y_j^\theta|^{1-\alpha}$$

$$(3.178) \quad (|\xi - y_j^\theta(\xi)|^{1-\alpha})' \leq C|x_N - y_j^\theta|^{1-\alpha}$$

$$(3.179) \quad (|\xi - y_j^\theta(\xi)|^{1-\alpha})'' \leq C|x_N - y_j^\theta|^{1-\alpha} + C(r-1)|x_N - y_j^\theta|^{-\alpha}$$

LEMMA 3.43. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq j < N$,*

$$(3.180) \quad V_{Nj} \leq Ch^2 \int_{x_{j-1}}^{x_j} |x_N - y|^{1-\alpha} + (r-1)|x_N - y|^{-\alpha} dy$$

Therefore,

$$(3.181) \quad \begin{aligned} I_3 &\leq Ch^2 \int_{\lceil \frac{N}{2} \rceil}^{N-1} |x_N - y|^{1-\alpha} + (r-1)|x_N - y|^{-\alpha} dy \\ &\leq Ch^2(|T - x_{N-1}|^{2-\alpha} + (r-1)|T - x_{N-1}|^{1-\alpha}) \end{aligned}$$

□

For $j = N$,

LEMMA 3.44.

(3.182)

$$V_{N,N} = \frac{1}{h_N^2} (T_{N-1,N-1} - 2T_{N,N} + T_{N+1,N+1}) \leq Ch^2 + C(r-1)h^2|T - x_{N-1}|^{1-\alpha}$$

Proof.

(3.183)

□

$$\begin{aligned} V_{N,N} = & \int_0^1 -\frac{\theta(1-\theta)^{2-\alpha}}{2} \frac{1}{h_N^2} (h_{N-1}^{4-\alpha} u''(y_{N-1}^\theta) - 2h_N^{4-\alpha} u''(y_N^\theta) + h_{N+1}^{4-\alpha} u''(y_{N+1}^\theta)) d\theta \\ & + \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{1}{h_N} \left(\frac{Q_{N \rightarrow N}^\theta(x_{N+1}) u'''(\eta_{N+1,1}^\theta) - Q_{N \rightarrow N}^\theta(x_i) u'''(\eta_{N,1}^\theta)}{h_N} \right) d\theta \\ & - \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{1}{h_N} \left(\frac{Q_{N \rightarrow N}^\theta(x_N) u'''(\eta_{N,1}^\theta) - Q_{N \rightarrow N}^\theta(x_{N-1}) u'''(\eta_{N-1,1}^\theta)}{h_N} \right) d\theta \\ & - \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{1}{h_N} \left(\frac{Q_{N \rightarrow N}^\theta(x_{N+1}) u'''(\eta_{N+1,2}^\theta) - Q_{N \rightarrow N}^\theta(x_N) u'''(\eta_{N,2}^\theta)}{h_N} \right) d\theta \\ & + \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{1}{h_N} \left(\frac{Q_{N \rightarrow N}^\theta(x_N) u'''(\eta_{N,2}^\theta) - Q_{N \rightarrow N}^\theta(x_{N-1}) u'''(\eta_{N-1,2}^\theta)}{h_N} \right) d\theta \end{aligned}$$

So combine Lemma 3.11, Theorem 3.37, Theorem 3.40, Lemma 3.44 We have

LEMMA 3.45.

$$R_N \leq C(r-1)h^2|T - x_{N-1}|^{1-\alpha} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

and with Theorem 3.39 we prove the Theorem 3.3

4. Convergence analysis.

4.1. Properties of some Matrices. Review subsection 2.1, we have got (2.10).

DEFINITION 4.1. We call one matrix an M matrix, which means its entries are positive on major diagonal and nonpositive on others, and strictly diagonally dominant in rows.

Now we have

LEMMA 4.2. Matrix A defined by (2.12) where (2.13) is an M matrix. And there exists a constant $C_A = C(T, \alpha, r)$ such that

$$(4.1) \quad S_i := \sum_{j=1}^{2N-1} a_{ij} \geq C_A(x_i^{-\alpha} + (2T - x_i)^{-\alpha})$$

Proof. From (2.14), we have

$$(4.2) \quad \sum_{j=1}^{2N-1} \tilde{a}_{ij} = \frac{1}{\Gamma(4-\alpha)} \left(\frac{|x_i - x_0|^{3-\alpha} - |x_i - x_1|^{3-\alpha}}{h_1} + \frac{|x_{2N} - x_i|^{3-\alpha} - |x_{2N-1} - x_i|^{3-\alpha}}{h_{2N}} \right)$$

Let

$$(4.3) \quad g(x) = g_0(x) + g_{2N}(x)$$

where

$$g_0(x) := \frac{-\kappa_\alpha}{\Gamma(4-\alpha)} \frac{|x - x_0|^{3-\alpha} - |x - x_1|^{3-\alpha}}{h_1}$$

$$g_{2N}(x) := \frac{-\kappa_\alpha}{\Gamma(4-\alpha)} \frac{|x_{2N} - x|^{3-\alpha} - |x_{2N-1} - x|^{3-\alpha}}{h_{2N}}$$

Thus

$$-\kappa_\alpha \sum_{j=1}^{2N-1} \tilde{a}_{ij} = g(x_i)$$

Then

$$(4.4) \quad S_i := \sum_{j=1}^{2N-1} a_{ij}$$

$$= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right)$$

$$= D_h^2 g_0(x_i) + D_h^2 g_{2N}(x_i)$$

When $i = 1$

$$(4.5) \quad D_h^2 g_0(x_1) = \frac{2}{h_1 + h_2} \left(\frac{1}{h_2} g_0(x_2) - \left(\frac{1}{h_1} + \frac{1}{h_2} \right) g_0(x_1) + \frac{1}{h_1} g_0(x_0) \right)$$

$$= \frac{2\kappa_\alpha}{\Gamma(4-\alpha)} \frac{h_1^{3-\alpha} + h_2^{3-\alpha} + 2h_1^{2-\alpha}h_2 - (h_1 + h_2)^{3-\alpha}}{(h_1 + h_2)h_1h_2}$$

$$= \frac{2\kappa_\alpha}{\Gamma(4-\alpha)} \frac{h_1^{3-\alpha} + h_2^{3-\alpha} + 2h_1^{2-\alpha}h_2 - (h_1 + h_2)^{3-\alpha}}{(h_1 + h_2)h_1^{1-\alpha}h_2} h_1^{-\alpha}$$

$$= \frac{2\kappa_\alpha}{\Gamma(4-\alpha)} \frac{1 + (2^r - 1)^{3-\alpha} + 2(2^r - 1) - (2^r)^{3-\alpha}}{2^r(2^r - 1)} h_1^{-\alpha}$$

but

$$(4.6) \quad 1 + (2^r - 1)^{3-\alpha} + 2(2^r - 1) - (2^r)^{3-\alpha} > 0$$

While for $i \geq 2$

$$(4.7) \quad \begin{aligned} D_h^2 g_0(x_i) &= g_0''(\xi), \quad \xi \in (x_{i-1}, x_{i+1}) \\ &= -\kappa_\alpha \frac{|\xi - x_0|^{1-\alpha} - |\xi - x_1|^{1-\alpha}}{\Gamma(2-\alpha)h_1} \\ &= \frac{\kappa_\alpha}{-\Gamma(1-\alpha)} |\xi - \eta|^{-\alpha}, \quad \eta \in [x_0, x_1] \\ &\geq \frac{\kappa_\alpha}{-\Gamma(1-\alpha)} x_{i+1}^{-\alpha} \geq \frac{\kappa_\alpha}{-\Gamma(1-\alpha)} 2^{-r\alpha} x_i^{-\alpha} \end{aligned}$$

So

$$(4.8) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g_0(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g_0(x_i) + \frac{1}{h_i} g_0(x_{i-1}) \right) \geq C x_i^{-\alpha}$$

symmetricly,

$$(4.9) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g_{2N}(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g_{2N}(x_i) + \frac{1}{h_i} g_{2N}(x_{i-1}) \right) \geq C(\alpha, r)(2T - x_i)^{-\alpha} \quad \square$$

Let

$$(4.10) \quad g(x) = \begin{cases} x, & 0 < x \leq T \\ 2T - x, & T < x < 2T \end{cases}$$

And define

$$(4.11) \quad G = \text{diag}(g(x_1), \dots, g(x_{2N-1}))$$

Then

LEMMA 4.3. *The matrix $B := AG$, the major diagonal is positive, and nonpositive on others. And there is a constant $C_{AG}, C = C(\alpha, r)$ such that*

$$(4.12) \quad M_i := \sum_{j=1}^{2N-1} b_{ij} \geq -C_{AG}(x_i^{1-\alpha} + (2T - x_i)^{1-\alpha}) + C \begin{cases} |T - x_{i-1}|^{1-\alpha}, & i \leq N \\ |x_{i+1} - T|^{1-\alpha}, & i \geq N \end{cases}$$

Proof.

$$b_{ij} = a_{ij}g(x_j) = -\kappa_\alpha \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} \tilde{a}_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{a}_{i,j} + \frac{1}{h_i} \tilde{a}_{i-1,j} \right) g(x_j)$$

Since

$$(4.13) \quad g(x) \equiv \Pi_h g(x)$$

by ??, we have

$$\begin{aligned}
 \tilde{M}_i &:= \sum_{j=1}^{2N-1} \tilde{b}_{ij} = \sum_{j=1}^{2N-1} \tilde{a}_{ij} g(x_j) \\
 &= \int_0^{2T} \frac{|x_i - y|^{1-\alpha}}{\Gamma(2-\alpha)} \Pi_h g(y) dy = \int_0^{2T} \frac{|x_i - y|^{1-\alpha}}{\Gamma(2-\alpha)} g(y) dy \\
 &= \frac{-2}{\Gamma(4-\alpha)} |T - x_i|^{3-\alpha} + \frac{1}{\Gamma(4-\alpha)} (x_i^{3-\alpha} + (2T - x_i)^{3-\alpha}) \\
 &:= w(x_i) = p(x_i) + q(x_i)
 \end{aligned}
 \tag{4.14}$$

Thus,

$$\begin{aligned}
 M_i &:= \sum_{j=1}^{2N-1} b_{ij} = \sum_{j=1}^{2N-1} a_{ij} g(x_j) \\
 &= -\kappa_\alpha \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} \tilde{M}_{i+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{M}_i + \frac{1}{h_i} \tilde{M}_{i-1} \right) \\
 &= D_h^2(-\kappa_\alpha p)(x_i) - \kappa_\alpha D_h^2 q(x_i)
 \end{aligned}
 \tag{4.15}$$

for $1 \leq i \leq N-1$, by Lemma A.1

$$\begin{aligned}
 D_h^2(-\kappa_\alpha p)(x_i) &:= -\kappa_\alpha \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} p(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) p(x_i) + \frac{1}{h_i} p(x_{i-1}) \right) \\
 &= \frac{2\kappa_\alpha}{\Gamma(2-\alpha)} |T - \xi|^{1-\alpha} \quad \xi \in (x_{i-1}, x_{i+1}) \\
 &\geq \frac{2\kappa_\alpha}{\Gamma(2-\alpha)} |T - x_{i-1}|^{1-\alpha}
 \end{aligned}
 \tag{4.16}$$

$$\begin{aligned}
 D_h^2(-\kappa_\alpha p)(x_N) &:= -\kappa_\alpha \frac{2}{h_N + h_{N+1}} \left(\frac{1}{h_{N+1}} p(x_{N+1}) - \left(\frac{1}{h_N} + \frac{1}{h_{N+1}} \right) p(x_N) + \frac{1}{h_N} p(x_{N-1}) \right) \\
 &= \frac{4\kappa_\alpha}{\Gamma(4-\alpha) h_N^2} h_N^{3-\alpha} \\
 &= \frac{4\kappa_\alpha}{\Gamma(4-\alpha)} (T - x_{N-1})^{1-\alpha}
 \end{aligned}
 \tag{4.17}$$

Symmetricly for $i \geq N$, we get

$$D_h^2(-\kappa_\alpha p)(x_i) \geq \frac{2\kappa_\alpha}{\Gamma(2-\alpha)} \begin{cases} |T - x_{i-1}|^{1-\alpha}, & i \leq N \\ |x_{i+1} - T|^{1-\alpha}, & i \geq N \end{cases}
 \tag{4.18}$$

Similarly, we can get

$$\begin{aligned}
 D_h^2 q(x_i) &:= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} q(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) q(x_i) + \frac{1}{h_i} q(x_{i-1}) \right) \\
 &\leq \frac{2^{r(\alpha-1)+1}}{\Gamma(2-\alpha)} (x_i^{1-\alpha} + (2T - x_i)^{1-\alpha}), \quad i = 1, \dots, 2N-1
 \end{aligned}
 \tag{4.19}$$

So, we get the result. \square

Notice that

$$(4.20) \quad x_i^{-\alpha} \geq (2T)^{-1} x_i^{1-\alpha}$$

We can get

THEOREM 4.4. *There exists a real $\lambda = \lambda(T, \alpha, r) > 0$ and $C = C(T, \alpha, r) > 0$ such that $B := A(\lambda I + G)$ is an M matrix. And*

$$(4.21) \quad M_i := \sum_{j=1}^{2N-1} b_{ij} \geq C(x_i^{-\alpha} + (2T - x_i)^{-\alpha}) + C \begin{cases} |T - x_{i-1}|^{1-\alpha}, & i \leq N \\ |x_{i+1} - T|^{1-\alpha}, & i \geq N \end{cases}$$

Proof. By Lemma 4.2 with C_A and Lemma 4.3 with C_{AG} , it's sufficient to take $\lambda = (C + 2TC_{AG})/C_A$, then

$$(4.22) \quad M_i \geq C \left((x_i^{-\alpha} + (1 - x_i)^{-\alpha}) + \begin{cases} |T - x_{i-1}|^{1-\alpha}, & i \leq N \\ |x_{i+1} - T|^{1-\alpha}, & i \geq N \end{cases} \right) \quad \square$$

4.2. Proof of Theorem 2.6. For equation

$$(4.23) \quad AU = F \Leftrightarrow A(\lambda I + G)(\lambda I + G)^{-1}U = F \quad \text{i.e.} \quad B(\lambda I + G)^{-1}U = F$$

which means

$$(4.24) \quad \sum_{j=1}^{2N-1} b_{ij} \frac{\epsilon_j}{\lambda + g(x_j)} = -\tau_i$$

where $\epsilon_i = u(x_i) - u_i$.

And if

$$(4.25) \quad \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| = \max_{1 \leq i \leq 2N-1} \left| \frac{\epsilon_i}{\lambda + g(x_i)} \right|$$

Then, since $B = A(\lambda I + G)$ is an M matrix, it is Strictly diagonally dominant. Thus,

$$\begin{aligned} |\tau_{i_0}| &= \left| \sum_{j=1}^{2N-1} b_{i_0,j} \frac{\epsilon_j}{\lambda + g(x_j)} \right| \\ &\geq b_{i_0,i_0} \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| - \sum_{j \neq i_0} |b_{i_0,j}| \left| \frac{\epsilon_j}{\lambda + g(x_j)} \right| \\ &\geq b_{i_0,i_0} \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| - \sum_{j \neq i_0} |b_{i_0,j}| \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| \\ &= \sum_{j=1}^{2N-1} b_{i_0,j} \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| \\ &= M_{i_0} \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| \end{aligned} \quad (4.26)$$

By Theorem 2.5 and Theorem 4.4,

We know that there exists constants $C_1(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)}, \|f\|_{\beta}^{(\alpha/2)})$, and $C_2(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that

$$(4.27) \quad \left| \frac{\epsilon_i}{\lambda + g(x_i)} \right| \leq \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| \leq C_1 h^{\min\{\frac{r\alpha}{2}, 2\}} + C_2(r-1)h^2$$

620 as $\lambda + g(x_i) \leq \lambda + T$

621 So, we can get

$$622 \quad (4.28) \quad |\epsilon_i| \leq C(\lambda + T)h^{\min\{\frac{\alpha}{2}, 2\}}$$

623 The convergency has been proved.

624 Remarks:

5. Experimental results.

5.1. $f \equiv 1$.

5.2. $f = x^\gamma, \gamma < 0$. Appendix A. Approximate of difference quotients.

LEMMA A.1. If $g(x) \in C^2(\Omega)$, there exists $\xi \in (x_{i-1}, x_{i+1})$ such that

$$(A.1) \quad D_h^2 g(x_i) = g''(\xi), \quad \xi \in (x_{i-1}, x_{i+1})$$

And if $g(x) \in C^4(\Omega)$, then

$$(A.2) \quad D_h^2 g(x_i) = g''(x_i) + \frac{h_{i+1} - h_i}{3} g'''(x_i) + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \int_{x_{i-1}}^{x_i} g'''(y) \frac{(y - x_{i-1})^3}{3!} dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} g'''(y) \frac{(x_{i+1} - y)^3}{3!} dy \right)$$

Proof.

$$g(x_{i-1}) = g(x_i) - (x_i - x_{i-1})g'(x_i) + \frac{(x_i - x_{i-1})^2}{2} g''(\xi_1), \quad \xi_1 \in (x_{i-1}, x_i)$$

$$g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i)g'(x_i) + \frac{(x_{i+1} - x_i)^2}{2} g''(\xi_2), \quad \xi_2 \in (x_i, x_{i+1})$$

Substitute them in the left side of (A.1), we have

$$\begin{aligned} D_h^2 g(x_i) &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (g(x_{i+1}) - g(x_i) + \frac{1}{h_i} (g(x_{i-1}) - g(x_i))) \right) \\ &= \frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2) \end{aligned}$$

Now, using [intermediate value theorem](#), there exists $\xi \in [\xi_1, \xi_2]$ such that

$$\frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2) = g''(\xi)$$

And the last equation can be obtained by

$$g(x_{i-1}) = g(x_i) - h_i g'(x_i) + \frac{h_i^2}{2} g''(x_i) - \frac{h_i^3}{3!} g'''(x_i) + \int_{x_{i-1}}^{x_i} g'''(y) \frac{(y - x_{i-1})^3}{3!} dy$$

$$g(x_{i+1}) = g(x_i) + h_{i+1} g'(x_i) + \frac{h_{i+1}^2}{2} g''(x_i) + \frac{h_{i+1}^3}{3!} g'''(x_i) + \int_{x_i}^{x_{i+1}} g'''(y) \frac{(x_{i+1} - y)^3}{3!} dy$$

Especially,

$$(A.3) \quad \begin{aligned} \int_{x_{i-1}}^{x_i} g'''(y) \frac{(y - x_{i-1})^3}{3!} dy &= \frac{h_i^4}{4!} g'''(\eta_1) \\ \int_{x_i}^{x_{i+1}} g'''(y) \frac{(x_{i+1} - y)^3}{3!} dy &= \frac{h_{i+1}^4}{4!} g'''(\eta_2) \end{aligned}$$

where $\eta_1 \in (x_{i-1}, x_i), \eta_2 \in (x_i, x_{i+1})$. □

LEMMA A.2. Denote $y_j^\theta = (1 - \theta)x_{j-1} + \theta x_j, \theta \in (0, 1)$,

$$(A.4) \quad u(y_j^\theta) - \Pi_h u(y_j^\theta) = -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in (x_{j-1}, x_j)$$

$$(A.5) \quad u(y_j^\theta) - \Pi_h u(y_j^\theta) = -\frac{\theta(1-\theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1-\theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1-\theta)^2 u'''(\eta_2))$$

where $\eta_1 \in (x_{j-1}, y_j^\theta), \eta_2 \in (y_j^\theta, x_j)$.

Proof. By Taylor expansion, we have

$$u(x_{j-1}) = u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(\xi_1), \quad \xi_1 \in (x_{j-1}, y_j^\theta)$$

$$u(x_j) = u(y_j^\theta) + (1-\theta) h_j u'(y_j^\theta) + \frac{(1-\theta)^2 h_j^2}{2!} u''(\xi_2), \quad \xi_2 \in (y_j^\theta, x_j)$$

Thus

$$\begin{aligned} u(y_j^\theta) - \Pi_h u(y_j^\theta) &= u(y_j^\theta) - (1-\theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1-\theta)}{2} h_j^2 (\theta u''(\xi_1) + (1-\theta)u''(\xi_2)) \\ &= -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [\xi_1, \xi_2] \end{aligned}$$

The second equation is similar,

$$u(x_{j-1}) = u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(y_j^\theta) - \frac{\theta^3 h_j^3}{3!} u'''(\eta_1)$$

$$u(x_j) = u(y_j^\theta) + (1-\theta) h_j u'(y_j^\theta) + \frac{(1-\theta)^2 h_j^2}{2!} u''(y_j^\theta) + \frac{(1-\theta)^3 h_j^3}{3!} u'''(\eta_2)$$

where $\eta_1 \in (x_{j-1}, y_j^\theta), \eta_2 \in (y_j^\theta, x_j)$. Thus

$$\begin{aligned} u(y_j^\theta) - \Pi_h u(y_j^\theta) &= u(y_j^\theta) - (1-\theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1-\theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1-\theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1-\theta)^2 u'''(\eta_2)) \end{aligned}$$

LEMMA A.3. For $x \in [x_{j-1}, x_j]$

$$(A.6) \quad \begin{aligned} |u(x) - \Pi_h u(x)| &= \left| \frac{x_j - x}{h_j} \int_{x_{j-1}}^x u'(y) dy - \frac{x - x_{j-1}}{h_j} \int_x^{x_j} u'(y) dy \right| \\ &\leq \int_{x_{j-1}}^{x_j} |u'(y)| dy \end{aligned}$$

If $x \in [0, x_1]$, with Corollary 2.4, we have

$$(A.7) \quad |u(x) - \Pi_h u(x)| \leq \int_0^{x_1} |u'(y)| dy \leq \int_0^{x_1} C y^{\alpha/2-1} dy \leq C \frac{2}{\alpha} x_1^{\alpha/2}$$

Similarly, if $x \in [x_{2N-1}, 1]$, we have

$$(A.8) \quad |u(x) - \Pi_h u(x)| \leq C \frac{2}{\alpha} (2T - x_{2N-1})^{\alpha/2} = C \frac{2}{\alpha} x_1^{\alpha/2}$$

LEMMA A.4.

$$(A.9) \quad b^{1-\theta}|a^\theta - b^\theta| \leq |a - b| \quad (\text{also } a^{1-\theta}|a^\theta - b^\theta| \leq |a - b|), \quad a, b \geq 0, \theta \in [0, 1]$$

Appendix B. Proofs of some technical details. Review that $h = \frac{1}{N}$ and the definition of \simeq in subsection 2.1

LEMMA B.1.

$$(B.1) \quad h_i \simeq \begin{cases} hx_i^{1-1/r}, & 1 \leq i \leq N \\ h(2T - x_{i-1})^{1-1/r}, & N < i \leq 2N \end{cases}$$

Since $i^r - (i-1)^r \simeq i^{r-1}$, for $i \geq 1$.

And

$$(B.2) \quad h_i \simeq h_{i+1}, \quad x_i \simeq x_{i+1}, \quad \text{for } 1 \leq i \leq 2N-1$$

LEMMA B.2. There is a constant C such that for $i = 1, 2, \dots, 2N-1$

$$(B.3) \quad |h_{i+1} - h_i| \leq Ch^2 \begin{cases} x_i^{1-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{1-2/r}, & N < i \leq 2N-1 \end{cases}$$

Proof. By (2.2),

$$(B.4) \quad h_{i+1} - h_i = \begin{cases} T \left(\left(\frac{i+1}{N} \right)^r - 2 \left(\frac{i}{N} \right)^r + \left(\frac{i-1}{N} \right)^r \right), & 1 \leq i \leq N-1 \\ 0, & i = N \\ -T \left(\left(\frac{2N-i-1}{N} \right)^r - 2 \left(\frac{2N-i}{N} \right)^r + \left(\frac{2N-i+1}{N} \right)^r \right), & N+1 \leq i \leq 2N-1 \end{cases}$$

Since

$$(B.5) \quad (i+1)^r - 2i^r + (i-1)^r \simeq r(r-1)i^{r-2}, \quad \text{for } i \geq 1$$

We get the result. \square

LEMMA B.3. there is a constant $C = C(T, \alpha, r, \|f\|_\beta^{(\alpha/2)})$ such that

$$(B.6) \quad \begin{aligned} & \frac{2}{h_i + h_{i+1}} \left| \frac{1}{h_i} \int_{x_{i-1}}^{x_i} f''(y) \frac{(y - x_{i-1})^3}{3!} dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} f''(y) \frac{(y - x_{i+1})^3}{3!} dy \right| \\ & \leq Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T - x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N-1 \end{cases} \end{aligned}$$

Proof. By Lemma 2.2, we have for $1 \leq i \leq N$

$$(B.7) \quad \left| \int_{x_{i-1}}^{x_i} f''(y) \frac{(y - x_{i-1})^3}{3!} dy \right| \leq \frac{\|f\|_\beta^{(\alpha/2)}}{3!} \int_{x_{i-1}}^{x_i} y^{-\alpha/2-2} (y - x_{i-1})^3 dy$$

For $i = 1$,

$$\int_{x_{i-1}}^{x_i} y^{-\alpha/2-2} (y - x_{i-1})^3 dy = \int_0^{x_1} y^{1-\alpha/2} dy = \frac{1}{2-\alpha/2} x_1^{2-\alpha/2} = \frac{1}{2-\alpha/2} x_1^{-\alpha/2-2} h_1^4$$

And for $2 \leq i \leq N$, since $x_i \simeq x_{i-1} \leq y \leq x_i$, we have

$$\int_{x_{i-1}}^{x_i} y^{-\alpha/2-2}(y-x_{i-1})^3 dy \simeq \int_{x_{i-1}}^{x_i} x_i^{-\alpha/2-2}(y-x_{i-1})^3 dy = \frac{1}{4!} x_i^{-\alpha/2-2} h_i^4$$

So for $1 \leq i \leq N$, we have

$$(B.8) \quad \left| \int_{x_{i-1}}^{x_i} f''(y) \frac{(y-x_{i-1})^3}{3!} dy \right| \leq C x_i^{-\alpha/2-2} h_i^4$$

and similarly,

$$(B.9) \quad \left| \int_{x_i}^{x_{i+1}} f''(y) \frac{(x_{i+1}-y)^3}{3!} dy \right| \leq C x_i^{-\alpha/2-2} h_{i+1}^4$$

Thus for $1 \leq i \leq N$, with Lemma B.1 we have

$$(B.10) \quad \begin{aligned} & \frac{2}{h_i + h_{i+1}} \left| \frac{1}{h_i} \int_{x_{i-1}}^{x_i} f''(y) \frac{(y-x_{i-1})^3}{3!} dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} f''(y) \frac{(y-x_{i+1})^3}{3!} dy \right| \\ & \leq C x_i^{-\alpha/2-2} \frac{2}{h_i + h_{i+1}} (h_i^3 + h_{i+1}^3) \simeq x_i^{-\alpha/2-2} h_i^2 \simeq x_i^{-\alpha/2-2} h^2 x_i^{2-2/r} \\ & = C h^2 x_i^{-\alpha/2-2/r} \end{aligned}$$

It's symmetric for $N < i \leq 2N - 1$. \square

LEMMA B.4. *By a standard error estimate for linear interpolation, and Corollary 2.4, There is a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ for $2 \leq j \leq N$,*

$$(B.11) \quad |u(y) - \Pi_h u(y)| \leq C h^2 y^{\alpha/2-2/r}, \quad \text{for } y \in [x_{j-1}, x_j]$$

symmetricly, for $N < j \leq 2N - 1$, we have

$$(B.12) \quad |u(y) - \Pi_h u(y)| \leq C h^2 (2T - y)^{\alpha/2-2/r}$$

LEMMA B.5. *There is a constant $C = C(\alpha, r)$ such that for all $1 \leq i < N/2$, $\max\{2i+1, i+3\} \leq j \leq 2N$, we have*

$$(B.13) \quad D_h^2 K_y(x_i) \leq C \frac{y^{-1-\alpha}}{\Gamma(-\alpha)}, \quad y \in [x_{j-1}, x_j]$$

Proof. Since $y \geq x_{j-1} > x_{i+1}$, by Lemma A.1, if $j-1 > i+1$

$$\begin{aligned} D_h^2 K_y(x_i) &= K_y''(\xi) = \frac{|y-\xi|^{-1-\alpha}}{\Gamma(-\alpha)}, \quad \xi \in (x_{i-1}, x_{i+1}) \\ &\leq \frac{(y-x_{i+1})^{-1-\alpha}}{\Gamma(-\alpha)} \\ &\leq (1 - (\frac{2}{3})^r)^{-1-\alpha} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} \end{aligned}$$

LEMMA B.6. *There is a constant $C = C(\alpha, r)$ such that for all $3 \leq i \leq N, k = \lceil \frac{i}{2} \rceil$, $1 \leq j \leq k-1$ and $y \in [x_{j-1}, x_j]$, we have*

$$(B.14) \quad D_h^2 K_y(x_i) \leq C \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)}$$

Proof. Since $y \leq x_j < x_{i-1}$, by Lemma A.1,

$$\begin{aligned} D_h^2 K_y(x_i) &= \frac{|\xi - y|^{-1-\alpha}}{\Gamma(-\alpha)}, \quad \xi \in (x_{i-1}, x_{i+1}) \\ &\leq \frac{(x_{i-1} - x_j)^{-1-\alpha}}{\Gamma(-\alpha)} \leq \frac{(x_{i-1} - x_{k-1})^{-1-\alpha}}{\Gamma(-\alpha)} \\ &\leq \left(\left(\frac{2}{3}\right)^r - \left(\frac{1}{2}\right)^r\right)^{-1-\alpha} \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)} \end{aligned}$$

□

LEMMA B.7. While $0 \leq i < N/2$, By Lemma A.3

$$\begin{aligned} |T_{i1}| &\leq C \int_0^{x_1} x_1^{\alpha/2} \frac{|x_i - y|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\ &= C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2} |x_i^{2-\alpha} - |x_i - x_1|^{2-\alpha}| \\ &\leq C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2+2-\alpha} = C \frac{1}{\Gamma(3-\alpha)} x_1^{2-\alpha/2} \quad 0 < 2-\alpha < 1 \end{aligned}$$

For $2 \leq j \leq N$, by Lemma A.2 and Corollary 2.4

$$\begin{aligned} |T_{ij}| &\leq \frac{C}{4} \int_{x_{j-1}}^{x_j} h_j^2 x_{j-1}^{\alpha/2-2} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\ &\leq \frac{C}{4\Gamma(3-\alpha)} h_j^2 x_{j-1}^{\alpha/2-2} ||x_j - x_i|^{2-\alpha} - |x_{j-1} - x_i|^{2-\alpha}| \end{aligned}$$

LEMMA B.8. There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that

$$\sum_{j=1}^3 S_{1j} \leq C h^2 x_1^{-\alpha/2-2/r}$$

$$\sum_{j=1}^4 S_{2j} \leq C h^2 x_2^{-\alpha/2-2/r}$$

Proof.

$$S_{1j} = \frac{2}{x_2} \left(\frac{1}{x_1} T_{0j} - \left(\frac{1}{x_1} + \frac{1}{h_2} \right) T_{1j} + \frac{1}{h_2} T_{2j} \right)$$

So, by Lemma B.7

$$S_{11} \leq \frac{2}{x_2 x_1} 4 \frac{C}{\Gamma(3-\alpha)} x_1^{2-\alpha/2} \leq C x_1^{-\alpha/2}$$

$$S_{12} \leq \frac{2}{x_2 x_1} \frac{C}{4\Gamma(3-\alpha)} h_2^2 x_1^{\alpha/2-2} (x_2^{2-\alpha} + 2h_2^{2-\alpha} + h_2^{2-\alpha}) \leq C x_1^{-\alpha/2}$$

$$S_{13} \leq \frac{2}{x_2 x_1} \frac{C}{4\Gamma(3-\alpha)} h_3^2 x_2^{\alpha/2-2} (x_3^{2-\alpha} + 2x_3^{2-\alpha} + h_3^{2-\alpha}) \leq C x_1^{-\alpha/2}$$

727 But

$$728 \quad x_1^{-\alpha/2} = T^{2/r} h^2 x_1^{-\alpha/2-2/r}$$

729 $i = 2$ is similar. □

730

731 **LEMMA B.9.** *There exists a constant $C = C(T, r, l)$ such that For $3 \leq i \leq N -$*
 732 *$1, \lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\},$*
 733 *when $\xi \in (x_{i-1}, x_{i+1}),$*

$$734 \quad (\text{B.19}) \quad (h_{j-i}^3(\xi))' \leq (r-1)Ch^2x_i^{1-2/r}h_j$$

735

$$736 \quad (\text{B.20}) \quad (h_{j-i}^4(\xi))' \leq (r-1)Ch^2x_i^{1-2/r}h_j^2$$

737 *Proof.* From (3.38)

$$738 \quad (\text{B.21}) \quad y'_{j-i}(x) = y_{j-i}^{1-1/r}(x)x^{1/r-1}$$

$$739 \quad (\text{B.22}) \quad y''_{j-i}(x) = \frac{1-r}{r}y_{j-i}^{1-2/r}(x)x^{1/r-2}Z_{j-i}$$

740 For $\xi \in (x_{i-1}, x_{i+1})$ and $2 \leq k \leq j \leq \min\{2i-1, N-1\}$, using Lemma B.1

$$741 \quad \xi \simeq x_i \simeq x_j$$

742

$$743 \quad h_{j-i}(\xi) \simeq h_j \simeq hx_j^{1-1/r} \simeq hx_i^{1-1/r}$$

$$744 \quad (\text{B.23}) \quad \begin{aligned} h'_{j-i}(\xi) &= y'_{j-i}(\xi) - y'_{j-i-1}(\xi) \\ &= \xi^{1/r-1}(y_{j-i}^{1-1/r}(\xi) - y_{j-i-1}^{1-1/r}(\xi)) \end{aligned}$$

745 Since

$$746 \quad (\text{B.24}) \quad \begin{aligned} y_{j-i}^{1-1/r}(\xi) - y_{j-i-1}^{1-1/r}(\xi) &\leq x_{j+1}^{1-1/r} - x_{j-2}^{1-1/r} \\ &= T^{1-1/r}N^{1-r}((j+1)^{r-1} - (j-2)^{r-1}) \\ &\leq C(r-1)j^{r-2}N^{1-r} \\ &= C(r-1)hx_j^{1-2/r} \end{aligned}$$

747 Therefore,

$$748 \quad (\text{B.25}) \quad h'_{j-i}(\xi) \leq Cx_i^{1/r-1}(r-1)hx_j^{1-2/r} \simeq (r-1)hx_i^{-1/r}$$

749 for $l = 3, 4$

$$750 \quad (\text{B.26}) \quad \begin{aligned} (h_{j-i}^l(\xi))' &= lh_{j-i}^{l-1}(\xi)h'_{j-i}(\xi) \\ &\leq Ch_{j-i}^{l-1}(\xi)(r-1)hx_i^{-1/r} \\ &\simeq Ch_j^{l-2}hx_j^{1-1/r}(r-1)hx_i^{-1/r} \\ &\simeq C(r-1)h^2x_i^{1-2/r}h_j^{l-2} \end{aligned}$$

Meanwhile, we can get

$$(B.27) \quad h_{j-i}^3(\xi) \simeq h_j^3 \leq Ch^2 x_i^{2-2/r} h_j$$

$$(B.28) \quad h_{j-i}^4(\xi) \simeq h_j^4 \leq Ch^2 x_i^{2-2/r} h_j^2 \quad \square$$

LEMMA B.10. *There exists a constant $C = C(T, r, l)$ such that For $3 \leq i \leq N-1$, $\lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\}$, when $\xi \in (x_{i-1}, x_{i+1})$,*

$$(B.29) \quad (h_{j-i}^3(\xi))'' \leq C(r-1)h^2 x_i^{-2/r} h_j$$

Proof.

$$(B.30) \quad (h_{j-i}^3(\xi))'' = 6h_{j-i}(\xi)(h'_{j-i}(\xi))^2 + 3h_{j-i}^2(\xi)h''_{j-i}(\xi)$$

By (B.25)

$$(B.31) \quad h_{j-i}(\xi)(h'_{j-i}(\xi))^2 \leq Ch_j(r-1)^2 h^2 x_i^{-2/r}$$

For the second partial

$$(B.32) \quad \begin{aligned} h''_{j-i}(\xi) &= y''_{j-i}(\xi) - y''_{j-i-1}(\xi) \\ &= \frac{1-r}{r} \xi^{1/r-2} (y_{j-i}^{1-2/r}(\xi) Z_{j-i} - y_{j-i-1}^{1-2/r}(\xi) Z_{j-i-1}) \\ &= \frac{1-r}{r} \xi^{1/r-2} ((y_{j-i}^{1-2/r}(\xi) - y_{j-i-1}^{1-2/r}(\xi)) Z_{j-i} + y_{j-i-1}^{1-2/r}(\xi) Z_1) \end{aligned}$$

but

$$(B.33) \quad \begin{aligned} |y_{j-i}^{1-2/r}(\xi) - y_{j-i-1}^{1-2/r}(\xi)| &\leq |x_{j+1}^{1-2/r} - x_{j-2}^{1-2/r}| \\ &= T^{1-2/r} N^{2-r} |(j+1)^{r-2} - (j-2)^{r-2}| \\ &\leq C|r-2|N^{2-r} j^{r-3} \\ &= C|r-2|h x_j^{1-3/r} \end{aligned}$$

So we can get

$$(B.34) \quad \begin{aligned} |h''_{j-i}(\xi)| &\leq C(r-1)x_i^{1/r-2} (|r-2|h x_i^{1-3/r} x_i^{1/r} + x_i^{1-2/r} h) \\ &\leq C(r-1)h x_i^{-1-1/r} \end{aligned}$$

Summarizes, we have

$$(B.35) \quad (h_{j-i}^3(\xi))'' \leq C(r-1)h^2 x_i^{-2/r} h_j \quad \square$$

proof of Lemma 3.19. From (3.38)

$$(B.36) \quad y'_{j-i}(x) = y_{j-i}^{1-1/r}(x) x^{1/r-1}$$

$$(B.37) \quad y''_{j-i}(x) = \frac{1-r}{r} y_{j-i}^{1-2/r}(x) x^{1/r-2} Z_{j-i}$$

773 Since

$$774 \quad y_{j-i}^\theta(\xi) \simeq x_j \simeq x_i$$

775 We have known

$$776 \quad (\text{B.38}) \quad u''(y_{j-i}^\theta(\xi)) \leq C(y_{j-i}^\theta(\xi))^{\alpha/2-2} \simeq x_j^{\alpha/2-2} \simeq x_i^{\alpha/2-2}$$

777

$$\begin{aligned} (u''(y_{j-i}^\theta(\xi)))' &= u'''(y_{j-i}^\theta(\xi))(y_{j-i}^\theta(\xi))' \\ &\leq Cx_i^{\alpha/2-3}\xi^{1/r-1}y_{j-i}^{1-1/r}(\xi) \\ &\simeq x_i^{\alpha/2-3}x_i^{1/r-1}x_i^{1-1/r} = Cx_i^{\alpha/2-3} \end{aligned}$$

779

$$\begin{aligned} (u''(y_{j-i}^\theta(\xi)))'' &= u''''(y_{j-i}^\theta(\xi))(y_{j-i}^\theta(\xi))'^2 + u'''(y_{j-i}^\theta(\xi))y_{j-i}^{\theta''}(\xi) \\ &\leq Cx_i^{\alpha/2-4} + Cx_i^{\alpha/2-3}\frac{r-1}{r}x_i^{1-2/r}x_i^{1/r-2}Z_{|j-i|+1} \\ &\leq Cx_i^{\alpha/2-4} + C\frac{r-1}{r}x_i^{\alpha/2-3}x_i^{-1-1/r}x_i^{1/r} \\ &= Cx_i^{\alpha/2-4} \end{aligned} \quad \square$$

Proof of Lemma 3.20.

$$\begin{aligned} 781 \quad (\text{B.41}) \quad |y_{j-i}^\theta(\xi) - \xi| &= |\theta(y_{j-i-1}(\xi) - \xi) + (1-\theta)(y_{j-i}(\xi) - \xi)| \\ &= \theta|y_{j-i-1}(\xi) - \xi| + (1-\theta)|y_{j-i}(\xi) - \xi| \end{aligned}$$

782 where $y_{j-i-1}(\xi) - \xi$ and $y_{j-i}(\xi) - \xi$ have the same sign (≥ 0 or ≤ 0), independent
783 with ξ .

784 Since $|y_{j-i}(\xi) - \xi| = \text{sign}(j-i)(y_{j-i}(\xi) - \xi)$ is increasing with ξ ,

$$\begin{aligned} 785 \quad (\text{B.42}) \quad \left(\frac{i-1}{i}\right)^r |x_j - x_i| &\leq |x_{j-1} - x_{i-1}| \leq |y_{j-i}(\xi) - \xi| \leq |x_{j+1} - x_{i+1}| \leq \left(\frac{i+1}{i}\right)^r |x_j - x_i| \end{aligned}$$

786 we have

$$787 \quad (\text{B.43}) \quad |y_{j-i}(\xi) - \xi| \simeq |x_j - x_i|$$

788 Similarly, $|y_{j-1-i}(\xi) - \xi| \simeq |x_{j-1} - x_i|$. Thus, with (B.41), (B.43) and (2.17) we get

$$789 \quad (\text{B.44}) \quad |y_{j-i}^\theta(\xi) - \xi| \simeq |y_j^\theta - x_i|$$

790 Next, since $|y_{j-i}^\theta(\xi) - \xi| = \text{sign}(j-i-1+\theta)(y_{j-i}^\theta(\xi) - \xi)$, so we can derivate it.

$$791 \quad (\text{B.45}) \quad (|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})' = (\alpha-1)|y_{j-i}^\theta(\xi) - \xi|^{-\alpha}|(y_{j-i}^\theta(\xi))' - 1|$$

792 While, similar with (B.41), we have

$$793 \quad (\text{B.46}) \quad |(y_{j-i}^\theta(\xi))' - 1| = (1-\theta)|y_{j-i-1}'(\xi) - 1| + \theta|y_{j-i}'(\xi) - 1|$$

By Lemma A.4 and (B.43), we have

$$\begin{aligned} |y'_{j-i}(\xi) - 1| &= \xi^{1/r-1} |y_{j-i}^{1-1/r}(\xi) - \xi^{1-1/r}| \\ &\leq \xi^{-1} |y_{j-i}(\xi) - \xi| \\ &\simeq x_i^{-1} |x_j - x_i| \end{aligned} \quad (B.47)$$

So similar with (B.44), we can get

$$|(y_{j-i}^\theta(\xi))' - 1| \leq C x_i^{-1} |y_j^\theta - x_i| \quad (B.48)$$

Combine with (B.44), we get

$$|(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'| \leq C |y_j^\theta - x_i|^{-\alpha} x_i^{-1} |y_j^\theta - x_i| = C |y_j^\theta - x_i|^{1-\alpha} x_i^{-1} \quad (B.49)$$

Finally, we have

$$\begin{aligned} (|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'' &= \alpha(\alpha-1) |y_{j-i}^\theta(\xi) - \xi|^{-\alpha-1} ((y_{j-i}^\theta(\xi))' - 1)^2 \\ &\quad + \text{sign}(j-i-1+\theta)(1-\alpha) |y_{j-i}^\theta(\xi) - \xi|^{-\alpha} (y_{j-i}^\theta(\xi))'' \end{aligned} \quad (B.50)$$

For

$$(y_{j-i}^\theta(\xi))'' = (1-\theta)y_{j-i-1}''(\xi) + \theta y_{j-i}''(\xi) \quad (B.51)$$

and

$$\begin{aligned} y_{j-i}''(\xi) &= \frac{1-r}{r} y_{j-i}^{1-2/r}(x) x^{1/r-2} Z_{j-i} \\ &\simeq \frac{1-r}{r} x_j^{1-2/r} x_i^{1/r-2} Z_{j-i} \end{aligned} \quad (B.52)$$

while by Lemma A.4

$$|Z_{j-i}| = |x_j^{1/r} - x_i^{1/r}| \leq |x_j - x_i| x_i^{1/r-1} \quad (B.53)$$

we have

$$|y_{j-i}''(\xi)| \leq C(r-1) x_i^{-2} |x_j - x_i| \quad (B.54)$$

Therefore

$$|(y_{j-i}^\theta(\xi))''| \leq C(r-1) x_i^{-2} |y_j^\theta - x_i| \quad (B.55)$$

Then, combine with (B.48),

$$|(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})''| \leq C |y_j^\theta - x_i|^{1-\alpha} x_i^{-2} \quad (B.56)$$

□

proof of Lemma 3.22. For $\lceil \frac{i}{2} \rceil \leq j \leq \min\{2i-1, N-1\}$

$$\begin{aligned} &\frac{Q_{j-i}^\theta(x_{i+1}) u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i) u'''(\eta_j^\theta)}{h_{i+1}} \\ &= \frac{Q_{j-i}^\theta(x_{i+1}) - Q_{j-i}^\theta(x_i)}{h_{i+1}} u'''(\eta_{j+1}^\theta) + Q_{j-i}^\theta(x_i) \frac{u'''(\eta_{j+1}^\theta) - u'''(\eta_j^\theta)}{h_{i+1}} \end{aligned} \quad (B.57)$$

816 Since mean value theorem

$$817 \quad (B.58) \quad \frac{Q_{j-i}^\theta(x_{i+1}) - Q_{j-i}^\theta(x_i)}{h_{i+1}} = Q_{j-i}^{\theta'}(\xi), \quad \xi \in (x_i, x_{i+1})$$

818 From (3.45) and Leibniz rule, by Lemma B.9 and Lemma 3.20, we have

$$819 \quad (B.59) \quad |Q_{j-i}^{\theta'}(\xi)| \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{1-2/r} h_j^2$$

820 And by Definition 3.15 and Lemma B.1

$$821 \quad (B.60) \quad Q_{j-i}^\theta(x_i) = h_j^4 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} \simeq Ch^2 x_i^{2-2/r} \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} h_j^2$$

822 With $\eta_j^\theta \in (x_{j-1}, x_j)$

$$823 \quad u'''(\eta_{j+1}^\theta) \leq C(\eta_{j+1}^\theta)^{\alpha/2-3} \simeq x_j^{\alpha/2-3} \simeq x_i^{\alpha/2-3}$$

824 and

$$\begin{aligned} \frac{u'''(\eta_{j+1}^\theta) - u'''(\eta_j^\theta)}{h_{i+1}} &= u''''(\eta) \frac{\eta_{j+1}^\theta - \eta_j^\theta}{h_{i+1}} \\ 825 \quad &\leq C\eta^{\alpha/2-4} \frac{x_{j+1} - x_{j-1}}{h_{i+1}} = C\eta^{\alpha/2-4} \frac{h_{j+1} + h_j}{h_{i+1}} \\ &\simeq x_j^{\alpha/2-4} \simeq x_i^{\alpha/2-4} \end{aligned}$$

826 So we have

$$\begin{aligned} &\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \\ 827 \quad (B.61) \quad &\leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{1-2/r} h_j^2 x_i^{\alpha/2-3} + Ch^2 x_i^{2-2/r} \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} h_j^2 x_{j-1}^{\alpha/2-4} \\ &= Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j^2 \end{aligned}$$

828 while $h_j \simeq h_i$, substitute into the inequality above, we get the goal

$$\begin{aligned} &\frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \right) \\ 829 \quad (B.62) \quad &\leq \frac{1}{h_i} Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j h_i \\ &= Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j \end{aligned}$$

830 While, the later is similar. □

831

832 LEMMA B.11. *There exists a constant $C = C(T, r)$ such that For $N/2 \leq i \leq$
 833 $N - 1$, $N + 2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1$, $l = 3, 4$, $\xi \in (x_{i-1}, x_{i+1})$, we have*

$$834 \quad (B.63) \quad h_{j-i}^l(\xi) \leq Ch_j^l \leq Ch^2 h_j^{l-2}$$

$$835 \quad (B.64) \quad (h_{j-i-1}^l(\xi))' \leq C(r-1)h^2 h_j^{l-2}$$

$$836 \quad (B.65) \quad (h_{j-i}^3(\xi))'' \leq C(r-1)h^2 h_j$$

Proof.

$$837 \quad (B.66) \quad \begin{aligned} (h_{j-i}(\xi))' &= y_{j-i}'(\xi) - y_{j-i-1}'(\xi) \\ &= \xi^{1/r-1}((2T - y_{j-i}(\xi))^{1-1/r} - (2T - y_{j-i-1}(\xi))^{1-1/r}) \leq 0 \end{aligned}$$

838 Thus,

$$839 \quad (B.67) \quad Ch_j \leq h_{j+1} \leq h_{j-i}(\xi) \leq h_{j-i}(x_{i-1}) = h_{j-1} \leq Ch_j$$

840 So as $4^{-r}T \leq 2T - x_j \leq T$, $2^{-r}T \leq x_i \leq T$, we have

$$841 \quad (B.68) \quad h_{j-i}^l(\xi) \leq Ch_j^l \leq Ch^2(2T - x_j)^{2-2/r} h_j^{l-2} \leq Ch^2 h_j^{l-2}$$

842 Since

$$843 \quad (B.69) \quad \begin{aligned} &|(2T - y_{j-i}(\xi))^{1-1/r} - (2T - y_{j-i-1}(\xi))^{1-1/r}| \\ &= |(Z_{2N-(j-i)} - \xi^{1/r})^{r-1} - (Z_{2N-(j-1-i)} - \xi^{1/r})^{r-1}| \\ &= (r-1)Z_1(Z_{2N-(j-i-\gamma)} - \xi^{1/r})^{r-2} \quad \gamma \in [0, 1] \\ &\leq C(r-1)h(2T - x_j)^{1-2/r} \end{aligned}$$

844 we have

$$845 \quad (B.70) \quad |(h_{j-i}(\xi))'| \leq C(r-1)h(2T - x_j)^{1-2/r} x_i^{1/r-1}$$

846 And

$$847 \quad (B.71) \quad \begin{aligned} (h_{j-i}^l(\xi))' &= l h_{j-i}^{l-1}(\xi) h_{j-i}'(\xi) \\ &\leq C(r-1)h_j^{l-1} h(2T - x_j)^{1-2/r} x_i^{1/r-1} \\ &\leq C(r-1)h^2 h_j^{l-2} (2T - x_j)^{2-3/r} x_i^{1-1/r} \\ &\leq C(r-1)h^2 h_j^{l-2} \end{aligned}$$

(B.72)

$$\begin{aligned} (h_{j-i}^3(\xi))'' &= 6h_{j-i}(\xi)(y_{j-i}'(\xi) - y_{j-i-1}'(\xi))^2 + 3h_{j-i}^2(\xi)(y_{j-i}''(\xi) - y_{j-i-1}''(\xi)) \\ &\leq C(r-1)h_j h^2 + Ch_j^2 \frac{1-r}{r} \xi^{1/r-2} ((2T - y_{j-i}(\xi))^{1-2/r} Z_{2N-(j-i)} - (2T - y_{j-i-1}(\xi))^{1-2/r} Z_{2N-(j-1-i)}) \\ &\leq C(r-1)h_j h^2 + C(r-1)h_j^2 (C(r-2)h(2T - x_j)^{1-3/r} Z_{2N-(j-i)} + Z_1(2T - x_{j-1})^{1-2/r}) \\ &\leq C(r-1)h_j h^2 + C(r-1)h_j^2 h = Ch^2 h_j \end{aligned}$$

849

850 **LEMMA B.12.** *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For*
 851 *$N/2 \leq i \leq N-1$, $N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1$, $\xi \in (x_{i-1}, x_{i+1})$, we have*

$$852 \quad (B.73) \quad u''(y_{j-i}^\theta(\xi)) \leq C$$

$$853 \quad (B.74) \quad (u''(y_{j-i}^\theta(\xi)))' \leq C$$

$$854 \quad (B.75) \quad (u''(y_{j-i}^\theta(\xi)))'' \leq C$$

Proof.

$$855 \quad (B.76) \quad x_{j-2} \leq y_{j-i}^\theta(\xi) \leq x_{j+1} \Rightarrow 4^{-r}T \leq 2T - y_{j-i}^\theta(\xi) \leq T$$

856 Thus, for $l = 2, 3, 4$,

$$857 \quad (B.77) \quad u^{(l)}(y_{j-i}^\theta(\xi)) \leq C(2T - y_{j-i}^\theta(\xi))^{\alpha/2-l} \leq C$$

858 and

$$\begin{aligned} (y_{j-i}^\theta(\xi))' &= \theta y_{j-1-i}'(\xi) + (1-\theta)y_{j-i-1}'(\xi) \\ 859 \quad (B.78) \quad &= \xi^{1/r-1}(\theta(2T - y_{j-1-i}(\xi))^{1-1/r} + (1-\theta)(2T - y_{j-i-1}(\xi))^{1-1/r}) \\ &\leq C(2T - x_{j-2})^{1-1/r} \leq C \end{aligned}$$

860 With

$$861 \quad (B.79) \quad Z_{2N-j-i} \leq 2T^{1/r}$$

$$\begin{aligned} 862 \quad (B.80) \quad &(y_{j-i}^\theta(\xi))'' = \theta y_{j-1-i}''(\xi) + (1-\theta)y_{j-i-1}''(\xi) \\ 863 \quad &= \frac{1-r}{r} \xi^{1/r-2}(\theta(2T - y_{j-i-1}(\xi))^{1-2/r} Z_{2N-(j-i-1)} + (1-\theta)(2T - y_{j-i}(\xi))^{1-2/r} Z_{2N-(j-i)}) \\ &\leq C(r-1) \end{aligned}$$

864 Therefore,

$$\begin{aligned} 865 \quad (B.81) \quad &(u''(y_{j-i}^\theta(\xi)))' = u'''(y_{j-i}^\theta(\xi))(y_{j-i}^\theta(\xi))' \\ &\leq C \end{aligned}$$

866

$$\begin{aligned} 867 \quad (B.82) \quad &(u''(y_{j-i}^\theta(\xi)))'' = u'''(y_{j-i}^\theta(\xi))(y_{j-i}^\theta(\xi))'^2 + u''''(y_{j-i}^\theta(\xi))y_{j-i}^{\theta''}(\xi) \\ &\leq C + C(r-1) = C \end{aligned} \quad \square$$

868

869 **LEMMA B.13.** *There exists a constant $C = C(T, \alpha, r)$ such that For $N/2 \leq i \leq$
 870 $N-1$, $N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1$, $\xi \in (x_{i-1}, x_{i+1})$*

$$871 \quad (B.83) \quad |y_{j-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_j^\theta - x_i|^{1-\alpha}$$

$$872 \quad (B.84) \quad |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'| \leq C|y_j^\theta - x_i|^{-\alpha}(|2T - x_i - y_j^\theta| + h_N)$$

$$\begin{aligned} 873 \quad (B.85) \quad &|(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})''| \leq C(r-1)|y_j^\theta - x_i|^{-\alpha} + C|y_j^\theta - x_i|^{-1-\alpha}(|2T - x_i - y_j^\theta| + h_N)^2 \end{aligned}$$

Proof. Since $y_{j-i-1}(\xi) > x_{j-2} \geq x_N > \xi$

$$(B.86) \quad y_{j-i}^\theta(\xi) - \xi = (1 - \theta)(y_{j-i-1}(\xi) - \xi) + \theta(y_{j-i}(\xi) - \xi) > 0$$

$$(B.87) \quad \begin{aligned} (y_{j-i}(\xi) - \xi)'' &= y_{j-i}''(\xi) \\ &= \frac{1-r}{r} \xi^{1/r-2} (2T - y_{j-i}(\xi))^{1-2/r} Z_{2N-(j-i)} \leq 0 \end{aligned}$$

It's concave, so

$$(B.88) \quad y_{j-i}(\xi) - \xi \geq \min_{\xi \in \{x_{i-1}, x_{i+1}\}} y_{j-i}(\xi) - \xi = \min\{x_{j+1} - x_{i+1}, x_{j-1} - x_{i-1}\} \geq C(x_j - x_i)$$

With (B.86), we have

$$(B.89) \quad |y_{j-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_j^\theta - x_i|^{1-\alpha}$$

By Lemma A.4

$$(B.90) \quad \begin{aligned} |y_{j-i}'(\xi) - 1| &= \xi^{1/r-1} |(2T - y_{j-i}(\xi))^{1-1/r} - \xi^{1-1/r}| \\ &\leq \xi^{-1} |2T - y_{j-i}(\xi) - \xi| \end{aligned}$$

$$(B.91) \quad \begin{aligned} |2T - \xi - y_{j-i}(\xi)| &\leq |2T - x_i - x_j| + |x_i - \xi| + |x_j - y_{j-i}(\xi)| \\ &\leq |2T - x_i - x_j| + h_{i+1} + h_j \\ &\leq C(|2T - x_i - x_j| + h_N) \end{aligned}$$

With $\xi \simeq x_i \simeq 1$,

$$(B.92) \quad |y_{j-i}'(\xi) - 1| \leq C(|2T - x_i - x_j| + h_N)$$

Thus,

$$(B.93) \quad \begin{aligned} |(y_{j-i}^\theta(\xi))' - 1| &\leq (1 - \theta)|y_{j-i-1}'(\xi) - 1| + \theta|y_{j-i}'(\xi) - 1| \\ &\leq C((1 - \theta)|2T - x_i - x_{j-1}| + \theta|2T - x_i - x_j| + h_N) \\ &= C(|2T - x_i - y_j^\theta| + h_N) \end{aligned}$$

So

$$(B.94) \quad \begin{aligned} |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'| &= |1 - \alpha||y_{j-i}^\theta(\xi) - \xi|^{-\alpha} |(y_{j-i}^\theta(\xi))' - 1| \\ &\leq C|y_j^\theta - x_i|^{-\alpha} (|2T - x_i - y_j^\theta| + h_N) \end{aligned}$$

(B.95)

$$\begin{aligned} |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})''| &\leq |1 - \alpha||y_{j-i}^\theta(\xi) - \xi|^{-\alpha} |(y_{j-i}^\theta(\xi) - \xi)''| + \alpha(\alpha - 1)|y_{j-i}^\theta(\xi) - \xi|^{-1-\alpha} (y_{j-i}^\theta(\xi))' - 1|^2 \\ &\leq C(r - 1)|y_j^\theta - x_i|^{-\alpha} + C|y_j^\theta - x_i|^{-1-\alpha} (|2T - x_i - y_j^\theta| + h_N)^2 \end{aligned}$$

Acknowledgments. We would like to acknowledge the assistance of volunteers in putting together this example manuscript and supplement.

REFERENCES

- [1] D. GILBARG AND N. S. TRUDINGER, *Poisson's Equation and the Newtonian Potential*, Springer Berlin Heidelberg, Berlin, Heidelberg, 1977, pp. 50–67, https://doi.org/10.1007/978-3-642-96379-7_4, https://doi.org/10.1007/978-3-642-96379-7_4.
- [2] X. ROS-OTON AND J. SERRA, *The dirichlet problem for the fractional laplacian: Regularity up to the boundary*, Journal de Mathématiques Pures et Appliquées, 101 (2014), pp. 275–302, <https://doi.org/https://doi.org/10.1016/j.matpur.2013.06.003>, <https://www.sciencedirect.com/science/article/pii/S0021782413000895>.