

AN EXAMPLE ARTICLE*

DIANNE DOE[†], PAUL T. FRANK[‡], AND JANE E. SMITH[‡]

Abstract. This is an example SIAM L^AT_EX article. This can be used as a template for new articles. Abstracts must be able to stand alone and so cannot contain citations to the paper's references, equations, etc. An abstract must consist of a single paragraph and be concise. Because of online formatting, abstracts must appear as plain as possible. Any equations should be inline.

Key words. example, L^AT_EX

MSC codes. 68Q25, 68R10, 68U05

1. Introduction. The introduction introduces the context and summarizes the manuscript. It is importantly to clearly state the contributions of this piece of work.

For $\Omega = (0, 2T)$, $1 < \alpha < 2$, suppose $f \in C^\beta(\Omega)$, $\beta > 4 - \alpha$, $\|f\|_\beta^{(\alpha/2)} < \infty$

$$(1.1) \quad \begin{cases} (-\Delta)^{\frac{\alpha}{2}} u(x) = f(x), & x \in \Omega \\ u(x) = 0, & x \in \mathbb{R} \setminus \Omega \end{cases}$$

where

$$(1.2) \quad (-\Delta)^{\frac{\alpha}{2}} u(x) = -\frac{\partial^\alpha u}{\partial |x|^\alpha} = -\kappa_\alpha \frac{d^2}{dx^2} \int_\Omega \frac{|x-y|^{1-\alpha}}{\Gamma(2-\alpha)} u(y) dy$$

$$(1.3) \quad \kappa_\alpha = -\frac{1}{2 \cos(\alpha\pi/2)} > 0$$

and the solution $u \in C^{\alpha/2}(\Omega)$.

2. Regularity.

Remark 2.1. 1. $C^k(U)$ is the set of all k -times continuously differentiable functions on open set U .

2. $C^\beta(U)$ is the collection of function f which for any $V \subset\subset U$ $f|_V \in C^\beta(\bar{V})$.

THEOREM 2.2. If $f \in C^\beta(\Omega)$, $\beta > 2$ and $\|f\|_\beta^{(\alpha/2)} < \infty$, then for $l = 0, 1, 2$

$$(2.1) \quad |f^{(l)}(x)| \leq \|f\|_\beta^{(\alpha/2)} \begin{cases} x^{-l-\alpha/2}, & \text{if } 0 < x \leq T \\ (2T-x)^{-l-\alpha/2}, & \text{if } T \leq x < 2T \end{cases}$$

THEOREM 2.3 (Regularity up to the boundary [1]).

$$(2.2) \quad \|u\|_{\beta+\alpha}^{(-\alpha/2)} \leq C \left(\|u\|_{C^{\alpha/2}(\mathbb{R})} + \|f\|_\beta^{(\alpha/2)} \right)$$

*Submitted to the editors DATE.

Funding: This work was funded by the Fog Research Institute under contract no. FRI-454.

[†]Imagination Corp., Chicago, IL (ddoe@imag.com, <http://www.imag.com/~ddoe/>).

[‡]Department of Applied Mathematics, Fictional University, Boise, ID (ptfrank@fictional.edu, jesmith@fictional.edu).

29 **COROLLARY 2.4.** *Let u be a solution of (1.1) on Ω . Then, for any $x \in \Omega$ and*
 30 *$l = 0, 1, 2, 3, 4$*

$$31 \quad (2.3) \quad |u^{(l)}(x)| \leq \|u\|_{\beta+\alpha}^{(-\alpha/2)} \begin{cases} x^{\alpha/2-l}, & \text{if } 0 < x \leq T \\ (2T-x)^{\alpha/2-l}, & \text{if } T \leq x < 2T \end{cases}$$

32 The paper is organized as follows. Our main results are in section 4, experimental
 33 results are in section 7, and the conclusions follow in section 8.

3. Numeric Format.

$$34 \quad (3.1) \quad x_i = \begin{cases} T \left(\frac{i}{N} \right)^r, & 0 \leq i \leq N \\ 2T - T \left(\frac{2N-i}{N} \right)^r, & N \leq i \leq 2N \end{cases}$$

35 where $r \geq 1$. And let

$$36 \quad (3.2) \quad h_j = x_j - x_{j-1}, \quad 1 \leq j \leq 2N$$

37 Let $\{\phi_j(x)\}_{j=1}^{2N-1}$ be standard hat functions, which are basis of the piecewise linear
 38 function space.

$$39 \quad (3.3) \quad \phi_j(x) = \begin{cases} \frac{1}{h_j}(x - x_{j-1}), & x_{j-1} \leq x \leq x_j \\ \frac{1}{h_{j+1}}(x_{j+1} - x), & x_j \leq x \leq x_{j+1} \\ 0, & \text{otherwise} \end{cases}$$

40 And then, we can approximate $u(x)$ with

$$41 \quad (3.4) \quad u_h(x) := \sum_{j=1}^{2N-1} u(x_j) \phi_j(x)$$

42 For convience, we denote

$$43 \quad (3.5) \quad I_h^{2-\alpha}(x_i) := \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u_h(y) dy$$

44 And now, we can approximate the operator (1.2) at x_i with

$$45 \quad (3.6) \quad \begin{aligned} D_h^\alpha u_h(x_i) &:= D_h^2 I_h^{2-\alpha}(x_i) \\ &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} I_h^{2-\alpha}(x_{i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) I_h^{2-\alpha}(x_i) + \frac{1}{h_{i+1}} I_h^{2-\alpha}(x_{i+1}) \right) \end{aligned}$$

46 Finally, we approximate the equation (1.1) with

$$47 \quad (3.7) \quad -\kappa_\alpha D_h^\alpha u_h(x_i) = f(x_i), \quad 1 \leq i \leq 2N-1$$

48 The discrete equation (3.7) can be written in matrix form

$$49 \quad (3.8) \quad AU = F$$

where U is unknown, $F = (f(x_1), \dots, f(x_{2N-1}))$. The matrix A is constructed as follows: Since

(3.9)

$$\begin{aligned}
 I_h^{2-\alpha}(x_i) &= \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u_h(y) dy \\
 &= \sum_{j=1}^{2N-1} \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u(x_j) \phi_j(y) dy \\
 &= \sum_{j=1}^{2N-1} u(x_j) \frac{1}{\Gamma(2-\alpha)} \int_{x_{j-1}}^{x_{j+1}} |x_i - y|^{1-\alpha} \phi_j(y) dy \\
 &= \sum_{j=1}^{2N-1} \frac{u(x_j)}{\Gamma(4-\alpha)} \left(\frac{|x_i - x_{j-1}|^{3-\alpha}}{h_j} - \frac{h_j + h_{j+1}}{h_j h_{j+1}} |x_i - x_j|^{3-\alpha} + \frac{|x_i - x_{j+1}|^{3-\alpha}}{h_{j+1}} \right) \\
 &=: \sum_{j=1}^{2N-1} \tilde{a}_{ij} u(x_j), \quad 0 \leq i \leq 2N
 \end{aligned}$$

Then, substitute in (3.6), we have

$$(3.10) \quad -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) = \sum_{j=1}^{2N-1} a_{ij} u(x_j)$$

where

$$(3.11) \quad a_{ij} = -\kappa_{\alpha} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \tilde{a}_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{a}_{i,j} + \frac{1}{h_{i+1}} \tilde{a}_{i+1,j} \right)$$

4. Main results. Here we state our main results; the proof is deferred to section 5 and section 6.

Let's denote $h = \frac{1}{N}$, we have

THEOREM 4.1 (Truncation Error). *If $f \in C^2(\Omega)$ and $\alpha \in (1, 2)$, and $u(x)$ is a solution of the equation (1.1), then there exists a constant $C_1, C_2 = C_1(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)}, \|f\|_{C^2(\Omega)}), C_2(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$, such that the truncation error of the discrete format satisfies*

$$\begin{aligned}
 |-\kappa_{\alpha} D_h^{\alpha} u_h(x_i) - f(x_i)| &\leq C_1 h^{\min\{\frac{r\alpha}{2}, 2\}} (x_i^{-\alpha} + (2T - x_i)^{-\alpha}) \\
 &\quad + C_2 h^2 \begin{cases} |T - x_{i-1}|^{1-\alpha}, & 1 \leq i \leq N \\ |T - x_{i+1}|^{1-\alpha}, & N < i \leq 2N-1 \end{cases}
 \end{aligned}
 \tag{4.1}$$

where $C_2 = 0$ if $r = 1$.

THEOREM 4.2 (Convergence). *The discrete equation (3.7) has solution U , and there exists a positive constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)}, \|f\|_{\beta}^{(\alpha/2)})$ such that the error between the numerical solution U with the exact solution $u(x_i)$ satisfies*

$$(4.2) \quad \max_{1 \leq i \leq 2N-1} |U_i - u(x_i)| \leq C h^{\min\{\frac{r\alpha}{2}, 2\}}$$

That means the numerical method has convergence order $\min\{\frac{r\alpha}{2}, 2\}$.

5. Proof of Theorem 4.1. For convience, let's denote

$$(5.1) \quad I^{2-\alpha}(x) = \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x-y|^{1-\alpha} u(y) dy$$

Then, the truncation error of the discrete format can be written as

$$(5.2) \quad \begin{aligned} -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) - f(x_i) &= -\kappa_{\alpha} (D_h^2 I_h^{2-\alpha}(x_i) - \frac{d^2}{dx^2} I^{2-\alpha}(x_i)) \\ &= -\kappa_{\alpha} D_h^2 (I_h^{2-\alpha} - I^{2-\alpha})(x_i) - \kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) \end{aligned}$$

5.1. Estimate of $-\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i)$.

THEOREM 5.1. *There exists a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$ such that*

$$(5.3) \quad \left| -\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) \right| \leq Ch^2 (x_i^{-\alpha/2-2/r} + (2T-x_i)^{-\alpha/2-2/r})$$

Proof. Since $f \in C^2(\Omega)$ and

$$(5.4) \quad \frac{d^2}{dx^2} (-\kappa_{\alpha} I^{2-\alpha}(x)) = f(x), \quad x \in \Omega,$$

we have $I^{2-\alpha} \in C^4(\Omega)$. Therefore, using equation (A.3) of Lemma A.1, for $1 \leq i \leq 2N-1$, we have

$$(5.5) \quad -\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) = \frac{h_{i+1} - h_i}{3} f'(x_i) + \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2))$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$. By Lemma B.2 and Theorem 2.2 we have 1.

$$(5.6) \quad \left| \frac{h_{i+1} - h_i}{3} f'(x_i) \right| \leq \frac{\|f\|_{\beta}^{(\alpha/2)}}{3} Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T-x_i)^{-\alpha/2-2/r}, & N < i \leq 2N-1 \end{cases}$$

2. See Proof 17, there is a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$ such that

$$(5.7) \quad \begin{aligned} &\left| \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \right| \\ &\leq Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T-x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N-1 \end{cases} \end{aligned}$$

Summarizes, we get the result. \square

5.2. Estimate of R_i . Now, we study the first part of (5.2)

$$(5.8) \quad D_h^2 (I^{2-\alpha} - I_h^{2-\alpha})(x_i) = D_h^2 \left(\int_0^{2T} (u(y) - u_h(y)) \frac{|y-x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \right)$$

For convience, let's denote

$$(5.9) \quad T_{ij} = \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) \frac{|y-x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy$$

92 And define

$$\begin{aligned}
 R_i &:= D_h^2(I^{2-\alpha} - I_h^{2-\alpha})(x_i) \\
 (5.10) \quad &= \frac{2}{h_i + h_{i+1}} \sum_{j=1}^{2N} \left(\frac{1}{h_i} T_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_{i+1}} T_{i+1,j} \right)
 \end{aligned}$$

94 We have some results about the estimate of R_i

95 **THEOREM 5.2.** *For $1 \leq i < N/2$, there exists $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that*

$$(5.11) \quad R_i \leq \begin{cases} Ch^2 x_i^{-\alpha/2-2/r}, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 (x_i^{-1-\alpha} \ln(i) + \ln(N)), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2} x_i^{-1-\alpha}, & \alpha/2 - 2/r < 0 \end{cases}$$

97

98 **THEOREM 5.3.** *For $N/2 \leq i \leq N$, there exists constant C, C_2 such that*

$$(5.12) \quad R_i \leq Ch^2 x_i^{-\alpha/2-2/r} + C_2 h^2 |T - x_{i-1}|^{1-\alpha}$$

100 where $C_2 = 0$ if $r = 1$.

101 And for $N < i \leq 2N - 1$, it is symmetric to the previous case.

102 To prove these results, we need some utils. Also for simplicity, we denote

DEFINITION 5.4.

$$(5.13) \quad S_{ij} = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} T_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_{i+1}} T_{i+1,j} \right)$$

104 then

$$(5.14) \quad R_i = \sum_{j=1}^{2N} S_{ij}$$

106 5.3. Proof of Theorem 5.2.

107 **LEMMA 5.5.** *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $1 \leq$
 108 $i < N/2$,*

$$(5.15) \quad \sum_{j=\max\{2i+1, i+3\}}^N S_{ij} \leq Ch^2 x_i^{-\alpha/2-2/r}$$

110 *Proof.* For $\max\{2i + 1, i + 3\} \leq j \leq N$, by Lemma C.1 and Lemma C.2

$$\begin{aligned}
 S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) D_h^2 \left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)} \right) (x_i) dy \\
 (5.16) \quad &\leq Ch^2 \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} dy \\
 &= Ch^2 \int_{x_{j-1}}^{x_j} y^{-\alpha/2-2/r-1} dy
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 \sum_{j=\max\{2i+1, i+3\}}^N S_{ij} &\leq Ch^2 \int_{x_{2i}}^{x_N} y^{-\alpha/2-2/r-1} dy \\
 &= \frac{C}{\alpha/2+2/r} h^2 (x_{2i}^{-\alpha/2-2/r} - T^{-\alpha/2-2/r}) \\
 &\leq \frac{C}{\alpha/2+2/r} 2^{r(-\alpha/2-2/r)} h^2 x_i^{-\alpha/2-2/r}
 \end{aligned}
 \tag{5.17}$$

LEMMA 5.6. *Thert exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $1 \leq i < N/2$,*

$$\sum_{j=N+1}^{2N} S_{ij} \leq \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}
 \tag{5.18}$$

Proof. For $1 \leq i < N/2, N+1 \leq j \leq 2N-1$, by equation (C.2) and Lemma C.2

$$\begin{aligned}
 S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) D_h^2 \left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)} \right) (x_i) dy \\
 &\leq \int_{x_{j-1}}^{x_j} Ch^2 (2T - y)^{\alpha/2-2/r} y^{-1-\alpha} dy \\
 &\leq Ch^2 T^{-1-\alpha} \int_{x_{j-1}}^{x_j} (2T - y)^{\alpha/2-2/r} dy
 \end{aligned}
 \tag{5.19}$$

$$\begin{aligned}
 \sum_{j=N+1}^{2N-1} S_{ij} &\leq CT^{-1-\alpha} h^2 \int_{x_N}^{x_{2N-1}} (2T - y)^{\alpha/2-2/r} dy \\
 &\leq CT^{-1-\alpha} h^2 \begin{cases} \frac{1}{\alpha/2-2/r+1} T^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 > 0 \\ \ln(T) - \ln(h_{2N}), & \alpha/2 - 2/r + 1 = 0 \\ \frac{1}{|\alpha/2-2/r+1|} h_{2N}^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \\
 &= \begin{cases} \frac{C}{\alpha/2-2/r+1} T^{-\alpha/2-2/r} h^2, & \alpha/2 - 2/r + 1 > 0 \\ CrT^{-1-\alpha} h^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ \frac{C}{|\alpha/2-2/r+1|} T^{-\alpha/2-2/r} h^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}
 \end{aligned}
 \tag{5.19}$$

And by Lemma A.3

$$S_{i,2N} \leq CT^{-1-\alpha} h_{2N}^{\alpha/2+1} = CT^{-\alpha/2} h^{r\alpha/2+r}$$

And when $\alpha/2 - 2/r + 1 \geq 0$,

$$h^{r\alpha/2+r} \leq h^2$$

Summarizes, we get the result.

For $i = 1, 2$.

LEMMA 5.7. *By Lemma C.5 , Lemma 5.5 and Lemma 5.6 we get*

$$R_1 = \sum_{j=1}^3 S_{1j} + \sum_{j=4}^{2N} S_{1j}$$

$$\leq Ch^2 x_1^{-\alpha/2-2/r} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

$$R_2 = \sum_{j=1}^4 S_{2j} + \sum_{j=5}^{2N} S_{2j}$$

$$\leq Ch^2 x_2^{-\alpha/2-2/r} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

For $3 \leq i < N/2$, we have a new separation of R_i , Let's denote $k = \lceil \frac{i}{2} \rceil$.

$$R_i = \sum_{j=1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right)$$

$$= \sum_{j=1}^{k-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right)$$

$$+ \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right)$$

$$+ \sum_{j=k+1}^{2i-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right)$$

$$+ \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1,2i} + T_{i-1,2i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2i} \right)$$

$$+ \sum_{j=2i+1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right)$$

$$= I_1 + I_2 + I_3 + I_4 + I_5$$

LEMMA 5.8. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $3 \leq i \leq N, k = \lceil \frac{i}{2} \rceil$*

$$|I_1| = \left| \sum_{j=1}^{k-1} S_{ij} \right| \leq \begin{cases} Ch^2 x_i^{-\alpha/2-2/r}, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 x_i^{-1-\alpha} \ln(i), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r} x_i^{-1-\alpha}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

Proof. For $2 \leq j \leq k-1$, by Lemma C.1 and Lemma C.3

$$\begin{aligned} S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) D_h^2 \left(\frac{|\cdot - y|^{1-\alpha}}{\Gamma(2-\alpha)} \right) (x_i) dy \\ &\leq Ch^2 \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)} dy \\ &= Ch^2 x_i^{-1-\alpha} \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} dy \end{aligned}$$

And by Lemma A.3 , Lemma C.3

$$S_{i1} \leq C x_1^{\alpha/2} x_1 x_i^{-1-\alpha} = C x_1^{\alpha/2+1} x_i^{-1-\alpha} = C T^{\alpha/2+1} h^{r\alpha/2+r} x_i^{-1-\alpha}$$

Therefore,

$$\begin{aligned} I_1 &= \sum_{j=1}^{k-1} S_{ij} = S_{i1} + \sum_{j=2}^{k-1} S_{ij} \\ &\leq Ch^{r\alpha/2+r} x_i^{-1-\alpha} + Ch^2 x_i^{-1-\alpha} \int_{x_1}^{x_{\lceil \frac{k}{2} \rceil - 1}} y^{\alpha/2-2/r} dy \\ &\leq Ch^{r\alpha/2+r} x_i^{-1-\alpha} + Ch^2 x_i^{-1-\alpha} \int_{x_1}^{2^{-r} x_i} y^{\alpha/2-2/r} dy \end{aligned}$$

But

$$\int_{x_1}^{2^{-r} x_i} y^{\alpha/2-2/r} dy \leq \begin{cases} \frac{1}{\alpha/2-2/r+1} (2^{-r} x_i)^{\alpha/2-2/r+1}, & \alpha/2-2/r+1 > 0 \\ \ln(2^{-r} x_i) - \ln(x_1), & \alpha/2-2/r+1 = 0 \\ \frac{1}{|\alpha/2-2/r+1|} x_1^{\alpha/2-2/r+1}, & \alpha/2-2/r+1 < 0 \end{cases}$$

So we have

$$I_1 \leq \begin{cases} \frac{C}{\alpha/2-2/r+1} h^2 x_i^{-\alpha/2-2/r}, & \alpha/2-2/r+1 > 0 \\ Ch^2 x_i^{-1-\alpha} \ln(i), & \alpha/2-2/r+1 = 0 \\ \frac{C}{|\alpha/2-2/r+1|} h^{r\alpha/2+r} x_i^{-1-\alpha}, & \alpha/2-2/r+1 < 0 \end{cases} \quad \square$$

DEFINITION 5.9. For convience, let's denote

$$V_{ij} = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right)$$

THEOREM 5.10. There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for

$$3 \leq i < N/2, k = \lceil \frac{i}{2} \rceil,$$

$$I_3 = \sum_{j=k+1}^{2i-1} V_{ij} \leq Ch^2 x_i^{-\alpha/2-2/r}$$

To estimate V_{ij} , we need some preparations.

155 LEMMA 5.11. Denote $y_j^\theta = \theta x_{j-1} + (1 - \theta)x_j, \theta \in [0, 1]$, by Lemma A.2

$$\begin{aligned}
 T_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\
 &= \int_{x_{j-1}}^{x_j} -\frac{\theta(1-\theta)}{2} h_j^2 u''(y_j^\theta) \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} \\
 &\quad + \frac{\theta(1-\theta)}{3!} h_j^3 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} (\theta^2 u'''(\eta_{j1}^\theta) - (1-\theta)^2 u'''(\eta_{j2}^\theta)) dy_j^\theta \\
 &= \int_0^1 -\frac{\theta(1-\theta)}{2} h_j^3 u''(y_j^\theta) \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} \\
 &\quad + \frac{\theta(1-\theta)}{3!} h_j^4 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} (\theta^2 u'''(\eta_{j1}^\theta) - (1-\theta)^2 u'''(\eta_{j2}^\theta)) d\theta
 \end{aligned}
 \tag{5.31}$$

157 where $\eta_{j1}^\theta \in [x_{j-1}, y_j^\theta], \eta_{j2}^\theta \in [y_j^\theta, x_j]$.

158 Now Let's construct a series of functions to represent T_{ij} .

DEFINITION 5.12.

$$y_{j-i}(x) = (x^{1/r} + Z_{j-i})^r, \quad Z_{j-i} = T^{1/r} \frac{j-i}{N}
 \tag{5.32}$$

160

$$y_{j-i}^\theta(x) = \theta y_{j-1-i}(x) + (1-\theta) y_{j-i}(x)
 \tag{5.33}$$

162

$$h_{j-i}(x) = y_{j-i}(x) - y_{j-i-1}(x)
 \tag{5.34}$$

164 Now, we define

$$P_{j-i}^\theta(x) = (h_{j-i}(x))^3 u''(y_{j-i}^\theta(x)) \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}
 \tag{5.35}$$

166

$$Q_{j-i}^\theta(x) = (h_{j-i}(x))^4 \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}
 \tag{5.36}$$

168 And now we can rewrite T_{ij}

169 LEMMA 5.13. For $2 \leq i \leq N, 2 \leq j \leq N$,

$$\begin{aligned}
 T_{ij} &= \int_0^1 -\frac{\theta(1-\theta)}{2} P_{j-i}^\theta(x_i) d\theta \\
 &\quad + \int_0^1 \frac{\theta(1-\theta)}{3!} (\theta^2 Q_{j-i}^\theta(x_i) u'''(\eta_{j1}^\theta) - (1-\theta)^2 Q_{j-i}^\theta(x_i) u'''(\eta_{j2}^\theta)) d\theta
 \end{aligned}
 \tag{5.37}$$

171 Immediately, we can see from (5.29) that

LEMMA 5.14. For $3 \leq i \leq N-1$, $3 \leq j \leq N-1$,

$$\begin{aligned}
 (5.38) \quad V_{ij} &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\
 &= \int_0^1 -\frac{\theta(1-\theta)}{2} D_h^2 P_{j-i}^\theta(x_i) d\theta \\
 &\quad + \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1}) u'''(\eta_{j+1,1}^\theta) - Q_{j-i}^\theta(x_i) u'''(\eta_{j,1}^\theta)}{h_{i+1}} \right) d\theta \\
 &\quad - \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i) u'''(\eta_{j,1}^\theta) - Q_{j-i}^\theta(x_{i-1}) u'''(\eta_{j-1,1}^\theta)}{h_i} \right) d\theta \\
 &\quad - \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1}) u'''(\eta_{j+1,2}^\theta) - Q_{j-i}^\theta(x_i) u'''(\eta_{j,2}^\theta)}{h_{i+1}} \right) d\theta \\
 &\quad + \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i) u'''(\eta_{j,2}^\theta) - Q_{j-i}^\theta(x_{i-1}) u'''(\eta_{j-1,2}^\theta)}{h_i} \right) d\theta
 \end{aligned}$$

To estimate V_{ij} , we first estimate $D_h^2 P_{j-i}^\theta(x_i)$, but By Lemma A.1,

$$(5.39) \quad D_h^2 P_{j-i}^\theta(x_i) = P_{j-i}^{\theta''}(\xi), \quad \xi \in [x_{i-1}, x_{i+1}]$$

By Leibniz formula, we calculate and estimate the derivations of h_{j-i}^3 , $u''(y_{j-i}^\theta(x))$

and $\frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$ separately.

Firstly, we have

LEMMA 5.15. There exists a constant $C = C(T, r)$ such that For $3 \leq i \leq N-1$, $\lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i-1, N-1\}$, $\xi \in [x_{i-1}, x_{i+1}]$,

$$(5.40) \quad h_{j-i}^3(\xi) \leq C h^2 x_i^{2-2/r} h_j$$

$$(5.41) \quad (h_{j-i}^3(\xi))' \leq C(r-1) h^2 x_i^{1-2/r} h_j$$

$$(5.42) \quad (h_{j-i}^3(\xi))'' \leq C(r-1) h^2 x_i^{-2/r} h_j$$

The proof of this theorem see Lemma C.6 and Lemma C.7

Second,

LEMMA 5.16. There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $3 \leq i \leq N-1$, $\lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i-1, N-1\}$, $\xi \in [x_{i-1}, x_{i+1}]$,

$$(5.43) \quad u''(y_{j-i}^\theta(\xi)) \leq C x_i^{\alpha/2-2}$$

$$(5.44) \quad (u''(y_{j-i}^\theta(\xi)))' \leq C x_i^{\alpha/2-3}$$

$$(5.45) \quad (u''(y_{j-i}^\theta(\xi)))'' \leq C x_i^{\alpha/2-4}$$

The proof of this theorem see Proof 24

And Finally, we have

LEMMA 5.17. There exists a constant $C = C(T, \alpha, r)$ such that For $3 \leq i \leq N-1$, $1 \leq j \leq \min\{2i-1, N-1\}$, $\xi \in [x_{i-1}, x_{i+1}]$,

$$(5.46) \quad |y_{j-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C |y_j^\theta - x_i|^{1-\alpha}$$

$$(5.47) \quad (|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})' \leq C|y_j^\theta - x_i|^{1-\alpha}x_i^{-1}$$

$$(5.48) \quad (|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'' \leq C|y_j^\theta - x_i|^{1-\alpha}x_i^{-2}$$

where $y_j^\theta = \theta x_{j-1} + (1-\theta)x_j$

The proof of this theorem see Proof 25

LEMMA 5.18. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For*
 $3 \leq i \leq N-1, \lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i-1, N-1\},$

$$(5.49) \quad D_h^2 P_{j-i}^\theta(x_i) \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j$$

where $y_j^\theta = \theta x_{j-1} + (1-\theta)x_j$

Proof. Since

$$(5.50) \quad D_h^2 P_{j-i}^\theta(x_i) = P_{j-i}^{\theta''}(\xi), \quad \xi \in [x_{i-1}, x_{i+1}]$$

From (5.35), using Leibniz formula and Lemma 5.15, Lemma 5.16 and Lemma 5.17□

LEMMA 5.19. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for*
 $3 \leq i < N, k = \lceil \frac{i}{2} \rceil.$
For $k \leq j \leq \min\{2i-1, N-1\},$

$$(5.51) \quad \begin{aligned} & \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \right) \\ & \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j \end{aligned}$$

And for $k+1 \leq j \leq \min\{2i, N\},$

$$(5.52) \quad \begin{aligned} & \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta) - Q_{j-i}^\theta(x_{i-1})u'''(\eta_{j-1}^\theta)}{h_i} \right) \\ & \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j \end{aligned}$$

where $\eta_j^\theta \in [x_{j-1}, x_j].$

proof see Proof 26

LEMMA 5.20. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for*
 $3 \leq i < N, k = \lceil \frac{i}{2} \rceil, k+1 \leq j \leq \min\{2i-1, N-1\},$

$$(5.53) \quad \begin{aligned} V_{ij} & \leq Ch^2 \int_0^1 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j d\theta \\ & = Ch^2 \int_{x_{j-1}}^{x_j} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} dy \end{aligned}$$

Proof. Since Lemma 5.14, by Lemma 5.18 and Lemma 5.19, we get the result immediately. \square

Now we can prove Theorem 5.10 using Lemma 5.20, $k = \lceil \frac{i}{2} \rceil$

$$\begin{aligned}
 I_3 &= \sum_{k+1}^{2i-1} V_{ij} \leq Ch^2 \int_{x_k}^{x_{2i-1}} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} dy \\
 &= Ch^2 \left(\frac{|x_k - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} + \frac{|x_{2i-1} - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} \right) x_i^{\alpha/2-2-2/r} \\
 &\leq Ch^2 x_i^{2-\alpha} x_i^{\alpha/2-2-2/r} = Ch^2 x_i^{-\alpha/2-2/r}
 \end{aligned}$$

LEMMA 5.21.

$$D_h P_{j-i}^\theta(x_i) := \frac{P_{k-i}^\theta(x_{i+1}) - P_{k-i}^\theta(x_i)}{h_{i+1}} = P_{j-i}^{\theta'}(\xi), \quad \xi \in [x_i, x_{i+1}]$$

Then, for $3 \leq i \leq N-1$, $k = \lceil \frac{i}{2} \rceil$,

$$D_h P_{k-i}^\theta(x_i) \leq Ch^2 x_i^{-\alpha/2-2/r} h_j$$

Proof. Using Leibniz formula, by Lemma 5.15, Lemma 5.16 and Lemma 5.17, we take $j = k+1$, $i = i+1$, we get

$$\begin{aligned}
 D_h P_{k-i}^\theta(x_i) &\leq Ch^2 x_{i+1}^{\alpha/2-2/r-1} |y_{k+1}^\theta - x_{i+1}|^{1-\alpha} h_{j+1} \\
 &\leq Ch^2 x_i^{\alpha/2-2/r-1} |y_k^\theta - x_i|^{1-\alpha} h_j \\
 &\leq Ch^2 x_i^{-\alpha/2-2/r} h_j
 \end{aligned}$$

LEMMA 5.22. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $3 \leq i < N$, $k = \lceil \frac{i}{2} \rceil$,*

$$I_2 = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right) \leq Ch^2 x_i^{-\alpha/2-2/r}$$

And for $3 \leq i < N/2$,

$$I_4 = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1,2i} + T_{i-1,2i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2i} \right) \leq Ch^2 x_i^{-\alpha/2-2/r}$$

Proof. In fact,

$$\begin{aligned}
 &\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \\
 &= \frac{1}{h_{i+1}} (T_{i+1,k} - T_{i,k}) + \frac{1}{h_{i+1}} (T_{i+1,k+1} - T_{i,k}) + \left(\frac{1}{h_{i+1}} - \frac{1}{h_i} \right) T_{i,k}
 \end{aligned}$$

241 While, by Lemma A.2

$$\begin{aligned}
 \frac{1}{h_{i+1}}(T_{i+1,k} - T_{i,k}) &= \int_{x_{k-1}}^{x_k} (u(y) - u_h(y)) \frac{|x_{i+1} - y|^{1-\alpha} - |x_i - y|^{1-\alpha}}{h_{i+1}\Gamma(2-\alpha)} dy \\
 &\leq \int_{x_{k-1}}^{x_k} h_j^2 u''(\eta) \frac{|\xi - y|^{-\alpha}}{\Gamma(1-\alpha)} dy \\
 &\leq Ch_j h^2 x_j^{2-2/r} x_{k-1}^{\alpha/2-2} |x_i - x_k|^{-\alpha} \\
 &\leq Ch_j h^2 x_i^{-\alpha/2-2/r}
 \end{aligned}
 \tag{5.61}$$

243 Thus,

$$\frac{2}{h_i + h_{i+1}} \frac{1}{h_{i+1}} (T_{i+1,k} - T_{i,k}) \leq Ch^2 x_i^{-\alpha/2-2/r}
 \tag{5.62}$$

245 For

$$\begin{aligned}
 \frac{1}{h_{i+1}}(T_{i+1,k+1} - T_{i,k}) &= \int_0^1 -\frac{\theta(1-\theta)}{2} \frac{P_{k-i}^\theta(x_{i+1}) - P_{k-i}^\theta(x_i)}{h_{i+1}} d\theta \\
 &+ \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{Q_{k-i}^\theta(x_{i+1})u'''(\eta_{j+1,1}^\theta) - Q_{k-i}^\theta(x_i)u'''(\eta_{j,1}^\theta)}{h_{i+1}} d\theta \\
 &- \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{Q_{k-i}^\theta(x_{i+1})u'''(\eta_{j+1,2}^\theta) - Q_{k-i}^\theta(x_i)u'''(\eta_{j,2}^\theta)}{h_{i+1}} d\theta
 \end{aligned}
 \tag{5.63}$$

247 And by Lemma 5.21

$$\frac{P_{k-i}^\theta(x_{i+1}) - P_{k-i}^\theta(x_i)}{h_{i+1}} \leq Ch^2 x_i^{-\alpha/2-2/r} h_j
 \tag{5.64}$$

249 And with Lemma 5.19, we can get

$$\frac{2}{h_i + h_{i+1}} \frac{1}{h_{i+1}} (T_{i+1,k+1} - T_{i,k}) \leq Ch^2 x_i^{-\alpha/2-2/r}
 \tag{5.65}$$

251 For the third term, by Lemma B.1, Lemma B.2 and Lemma A.2

$$\begin{aligned}
 \frac{2}{h_i + h_{i+1}} \frac{h_{i+1} - h_i}{h_i h_{i+1}} T_{i,k} &\leq h_i^{-3} h^2 x_i^{1-2/r} h_k Ch_k^2 x_{k-1}^{\alpha/2-2} |x_k - x_i|^{1-\alpha} \\
 &\leq Ch^2 x_i^{-\alpha/2-2/r}
 \end{aligned}
 \tag{5.66}$$

253 Summarizes, we have

$$I_2 \leq Ch^2 x_i^{-\alpha/2-2/r}
 \tag{5.67}$$

255 The case for I_4 is similar. □

256 Now combine Lemma 5.8, Lemma 5.22, Theorem 5.10, Lemma 5.5 and Lemma 5.6
 257 to get the final result.

258 For $3 \leq i < N/2$

$$\begin{aligned}
 R_i &= I_1 + I_2 + I_3 + I_4 + I_5 \\
 &\leq Ch^2 x_i^{-\alpha/2-2/r} + \begin{cases} Ch^2 x_i^{-\alpha/2-2/r}, & r\alpha/2 + r - 2 > 0 \\ Ch^2 (x_i^{1-\alpha} \ln(i) + \ln(N)), & r\alpha/2 + r - 2 = 0 \\ Ch^{r\alpha/2+r} x_i^{1-\alpha}, & r\alpha/2 + r - 2 < 0 \end{cases}
 \end{aligned}
 \tag{5.68}$$

Combine with $i = 1, 2$, we get for $1 \leq i \leq N/2$

$$(5.69) \quad R_i \leq \begin{cases} Ch^2 x_i^{-\alpha/2-2/r}, & r\alpha/2 + r - 2 > 0 \\ Ch^2 (x_i^{-1-\alpha} \ln(i) + \ln(N)), & r\alpha/2 + r - 2 = 0 \\ Ch^{r\alpha/2+r} x_i^{-1-\alpha}, & r\alpha/2 + r - 2 < 0 \end{cases}$$

5.4. Proof of Theorem 5.3. For $N/2 \leq i < N, k = \lceil \frac{i}{2} \rceil$, we have

$$(5.70) \quad \begin{aligned} R_i &= \sum_{j=1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\ &= \sum_{j=1}^{k-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\ &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right) \\ &\quad + \sum_{j=k+1}^{N-1} + \sum_{j=N}^{N+1} + \sum_{j=N+2}^{2N-\lceil \frac{N}{2} \rceil} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\ &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1,2i} + T_{i-1,2i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2i} \right) \\ &\quad + \sum_{j=2N-\lceil \frac{N}{2} \rceil+2}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\ &= I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7 \end{aligned}$$

We have estimate I_1 in Lemma 5.8 and I_2 in Lemma 5.22. We can control I_3 in similar with Theorem 5.10 by Lemma 5.20 where $2i - 1 \geq N - 1$

$$(5.71) \quad \begin{aligned} I_3 &= \sum_{j=k+1}^{N-1} V_{ij} \leq Ch^2 \int_{x_k}^{x_{N-1}} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} dy \\ &= Ch^2 \left(\frac{|x_k - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} + \frac{|x_{N-1} - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} \right) x_i^{\alpha/2-2-2/r} \\ &\leq Ch^2 x_i^{2-\alpha} x_i^{\alpha/2-2-2/r} = Ch^2 x_i^{-\alpha/2-2/r} \end{aligned}$$

For I_4 , we have

LEMMA 5.23. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that, for $N/2 \leq i < N - 1$*

$$(5.72) \quad \begin{aligned} V_{iN} &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,N+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,N} + \frac{1}{h_i} T_{i-1,N-1} \right) \\ &\leq Ch^2 |T - x_{i-1}|^{1-\alpha} \end{aligned}$$

Proof. We use the similar skill in the last section, but more complicated. for $j = N$, Let

$$(5.73) \quad y_{i \rightarrow N}(x) = \frac{x^{1/r} - Z_i}{Z_1} h_N + T, \quad Z_i = T^{1/r} \frac{i}{N}, x_N = T$$

274 and

$$275 \quad (5.74) \quad y_{i \rightarrow N-1}(x) = (x^{1/r} + Z_{N-1-i})^r, \quad Z_{N-1-i} = T^{1/r} \frac{N-1-i}{N}$$

276 Thus,

$$\begin{aligned} 277 \quad y_{i \rightarrow N}(x_{i-1}) &= x_{N-1}, & y_{i \rightarrow N}(x_i) &= x_N, & y_{i \rightarrow N}(x_{i+1}) &= x_{N+1} \\ 278 \quad y_{i \rightarrow N-1}(x_{i-1}) &= x_{N-2}, & y_{i \rightarrow N}(x_i) &= x_{N-1}, & y_{i \rightarrow N}(x_{i+1}) &= x_N \end{aligned}$$

279 Then, define

$$280 \quad (5.75) \quad y_{i \rightarrow N}^\theta(x) = \theta y_{i \rightarrow N-1}(x) + (1-\theta) y_{i \rightarrow N}(x)$$

281

$$282 \quad (5.76) \quad h_{i \rightarrow N}(x) = y_{i \rightarrow N}(x) - y_{i \rightarrow N-1}(x)$$

283 We have

$$284 \quad (5.77) \quad y_{i \rightarrow N-1}'(x) = y_{i \rightarrow N-1}^{1-1/r}(x) x^{1/r-1}$$

$$285 \quad (5.78) \quad y_{i \rightarrow N-1}''(x) = \frac{1-r}{r} y_{i \rightarrow N-1}^{1-2/r}(x) x^{1/r-2} Z_{N-1-i}$$

$$286 \quad (5.79) \quad y_{i \rightarrow N}'(x) = \frac{1}{r} \frac{h_N}{Z_1} x^{1/r-1}$$

$$287 \quad (5.80) \quad y_{i \rightarrow N}''(x) = \frac{1-r}{r^2} \frac{h_N}{Z_1} x^{1/r-2}$$

288

$$289 \quad (5.81) \quad P_{i \rightarrow N}^\theta(x) = (h_{i \rightarrow N}(x))^3 \frac{|y_{i \rightarrow N}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)} u''(y_{i \rightarrow N}^\theta(x))$$

290

$$291 \quad (5.82) \quad Q_{i \rightarrow N}^\theta(x) = (h_{i \rightarrow N}(x))^4 \frac{|y_{i \rightarrow N}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

292 Similar with Lemma 5.13, we can get for $l = -1, 0, 1$,

$$\begin{aligned} 293 \quad (5.83) \quad T_{i+l, N+l} &= \int_0^1 -\frac{\theta(1-\theta)}{2} P_{i \rightarrow N}^\theta(x_{i+l}) d\theta \\ &+ \int_0^1 \frac{\theta(1-\theta)}{3!} Q_{i \rightarrow N}^\theta(x_{i+l}) (\theta^2 u'''(\eta_{N+l,1}^\theta) - (1-\theta)^2 u'''(\eta_{N+l,2}^\theta)) d\theta \end{aligned}$$

294 So we have

(5.84)

$$\begin{aligned}
 V_{iN} &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1, N+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i, N} + \frac{1}{h_i} T_{i-1, N-1} \right) \\
 &= \int_0^1 -\frac{\theta(1-\theta)}{2} D_h^2 P_{i \rightarrow N}^\theta(x_i) d\theta \\
 &\quad + \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{i \rightarrow N}^\theta(x_{i+1}) u'''(\eta_{N+1,1}^\theta) - Q_{i \rightarrow N}^\theta(x_i) u'''(\eta_{N,1}^\theta)}{h_{i+1}} \right) d\theta \\
 &\quad - \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{i \rightarrow N}^\theta(x_i) u'''(\eta_{N,1}^\theta) - Q_{i \rightarrow N}^\theta(x_{i-1}) u'''(\eta_{N-1,1}^\theta)}{h_i} \right) d\theta \\
 &\quad - \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{i \rightarrow N}^\theta(x_{i+1}) u'''(\eta_{N+1,2}^\theta) - Q_{i \rightarrow N}^\theta(x_i) u'''(\eta_{N,2}^\theta)}{h_{i+1}} \right) d\theta \\
 &\quad + \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{i \rightarrow N}^\theta(x_i) u'''(\eta_{N,2}^\theta) - Q_{i \rightarrow N}^\theta(x_{i-1}) u'''(\eta_{N-1,2}^\theta)}{h_i} \right) d\theta
 \end{aligned}$$

296 To estimate $D_h^2 P_{i \rightarrow N}^\theta(x_i) = P_{i \rightarrow N}^{\theta''}(\xi), \xi \in [x_{i-1}, x_{i+1}]$,

$$297 \quad h_{i \rightarrow N}(\xi) \leq 2h_N$$

298

LEMMA 5.24.

$$299 \quad (5.85) \quad h_{i \rightarrow N}^3(\xi) \leq Ch_N^3 \leq Ch^3$$

$$300 \quad (5.86) \quad (h_{i \rightarrow N}^3(\xi))' \leq C(r-1)h_N^2 h \leq C(r-1)h^3$$

$$301 \quad (5.87) \quad (h_{i \rightarrow N}^3(\xi))'' \leq C(r-1)h_N^2 \leq C(r-1)h^2$$

Proof.

$$\begin{aligned}
 (h_{i \rightarrow N}^l(\xi))' &= l h_{i \rightarrow N}^{l-1}(\xi) (y_{i \rightarrow N}'(\xi) - y_{i \rightarrow N-1}'(\xi)) \\
 &= l h_{i \rightarrow N}^{l-1}(\xi) x_i^{1/r-1} \left(\frac{1}{r} \frac{h_N}{Z_1} - y_{i \rightarrow N-1}^{1-1/r}(\xi) \right)
 \end{aligned}$$

303 while

(5.89)

$$\begin{aligned}
 \left| \frac{1}{r} \frac{h_N}{Z_1} - y_{i \rightarrow N-1}^{1-1/r}(\xi) \right| &= \left| \frac{1}{r} \frac{x_N - (x_N^{1/r} - Z_1)^r}{Z_1} - \eta^{1-1/r} \right| \quad \eta \in [x_{N-2}, x_N] \\
 &= T^{1-1/r} \left| \left(\frac{N-t}{N} \right)^{r-1} - \left(\frac{N-s}{N} \right)^{r-1} \right| \quad t \in [0, 1], s \in [0, 2] \\
 &\leq T^{1-1/r} \left| 1 - \left(\frac{N-2}{N} \right)^{r-1} \right| \leq CT^{1-1/r} (r-1) \frac{2}{N}
 \end{aligned}$$

305 Thus,

$$306 \quad (5.90) \quad (h_{i \rightarrow N}^l(\xi))' \leq C(r-1)h_N^{l-1} x_i^{1/r-1} h$$

307 And

(5.91)

$$\begin{aligned} (h_{i \rightarrow N}^3(\xi))'' &= 6h_{i \rightarrow N}(\xi)(y_{i \rightarrow N}'(\xi) - y_{i \rightarrow N-1}'(\xi))^2 \\ &\quad + 3h_{i \rightarrow N}^2(\xi)(y_{i \rightarrow N}''(\xi) - y_{i \rightarrow N-1}''(\xi)) \\ &\leq C(r-1)^2 h_N x_i^{2/r-2} h^2 + C h_N^2 \frac{r-1}{r} x_i^{1/r-2} \left(\frac{h_N}{r Z_1} - y_{i \rightarrow N-1}^{1-2/r}(\xi) Z_{N-1-i} \right) \end{aligned}$$

308

309

310

$$\left| \frac{h_N}{r Z_1} - y_{i \rightarrow N-1}^{1-2/r}(\xi) Z_{N-1-i} \right| \leq T^{1-1/r} + C x_N^{1-2/r} x_N^{1/r} = C T^{1-1/r}$$

311 So

312 (5.92)

$$\begin{aligned} (h_{i \rightarrow N}^3(\xi))'' &\leq C(r-1)^2 h_N x_i^{2/r-2} h^2 + C h_N^2 \frac{1-r}{r} x_i^{1/r-2} \\ &\leq C(r-1) h_N^2 x_i^{1/r-1} \end{aligned}$$

□

LEMMA 5.25.

313 (5.93)

$$u''(y_{i \rightarrow N}^\theta(\xi)) \leq C x_{N-2}^{-\alpha/2-2} \leq C$$

314 (5.94)

$$(u''(y_{i \rightarrow N}^\theta(\xi)))' \leq C$$

315 (5.95)

$$(u''(y_{i \rightarrow N}^\theta(\xi)))'' \leq C$$

Proof.

316 (5.96)

$$\begin{aligned} (u''(y_{i \rightarrow N}^\theta(\xi)))' &= u'''(y_{i \rightarrow N}^\theta(\xi)) y_{i \rightarrow N}'^\theta(\xi) \\ &\leq C(\theta y_{i \rightarrow N-1}'(\xi) + (1-\theta) y_{i \rightarrow N}'(\xi)) \\ &\leq C x_i^{1/r-1} (\theta y_{i \rightarrow N-1}^{1-1/r}(\xi) + (1-\theta) \frac{h_N}{r Z_1}) \\ &\leq C x_i^{1/r-1} x_N^{1-1/r} \end{aligned}$$

317 And

(5.97)

$$\begin{aligned} (u''(y_{i \rightarrow N}^\theta(\xi)))'' &= u''''(y_{i \rightarrow N}^\theta(\xi)) (y_{i \rightarrow N}'^\theta(\xi))^2 + u'''(y_{i \rightarrow N}^\theta(\xi)) y_{i \rightarrow N}''^\theta(\xi) \\ &\leq C x_i^{2/r-2} x_N^{2-2/r} + C \frac{r-1}{r} x_i^{1/r-2} (\theta x_N^{1-2/r} Z_{N-1-i} + (1-\theta) \frac{h_N}{r Z_1}) \\ &\leq C x_i^{2/r-2} + C(r-1) x_i^{1/r-2} T^{1-1/r} \end{aligned}$$

318

□

LEMMA 5.26.

319 (5.98)

$$|y_{i \rightarrow N}^\theta(\xi) - \xi|^{1-\alpha} \leq C |y_N^\theta - x_i|^{1-\alpha}$$

320 (5.99)

$$(|y_{i \rightarrow N}^\theta(\xi) - \xi|^{1-\alpha})' \leq C |y_N^\theta - x_i|^{1-\alpha}$$

321 (5.100)

$$(|y_{i \rightarrow N}^\theta(\xi) - \xi|^{1-\alpha})'' \leq C(r-1) |y_N^\theta - x_i|^{-\alpha} + |y_N^\theta - x_i|^{1-\alpha}$$

Proof.

(5.101)

$$\begin{aligned} (y_{i \rightarrow N}^\theta(\xi) - \xi)' &= (\theta(y_{i \rightarrow N-1}(\xi) - \xi) + (1-\theta)(y_{i \rightarrow N}(\xi) - \xi))' \\ &= \theta(y_{i \rightarrow N-1}'(\xi) - 1) + (1-\theta)(y_{i \rightarrow N}'(\xi) - 1) \\ &= \theta \xi^{1/r-1} (y_{i \rightarrow N-1}^{1-1/r}(\xi) - \xi^{1-1/r}) + (1-\theta) \xi^{1/r-1} \left(\frac{h_N}{r Z_1} - \xi^{1-1/r} \right) \end{aligned}$$

322

$$(y_{i \rightarrow N}^\theta(\xi) - \xi)'' = \theta(y_{i \rightarrow N-1}''(\xi)) + (1 - \theta)(y_{i \rightarrow N}''(\xi))$$

$$= \frac{1-r}{r} \xi^{1/r-2} (\theta y_{i \rightarrow N-1}^{1-2/r}(\xi) Z_{N-1-i} + (1 - \theta) \frac{h_N}{r Z_1}) \leq 0$$

And

$$|(y_{i \rightarrow N}^\theta(\xi) - \xi)'| \leq C(r-1) \xi^{1/r-2} T^{1-1/r}$$

We have known

$$C|x_{N-1} - x_i| \leq |y_{i \rightarrow N-1}(\xi) - \xi| \leq C|x_{N-1} - x_i|$$

If $\xi \leq x_{N-1}$, then $(y_{i \rightarrow N}(\xi) - \xi)' \geq 0$, so

$$C|x_N - x_i| \leq |x_{N-1} - x_{i-1}| \leq |y_{i \rightarrow N}^\theta(\xi) - \xi| \leq |x_{N+1} - x_{i+1}| \leq C|x_N - x_i|$$

If $i = N-1$ and $\xi \in [x_{N-1}, x_N]$, then $y_{i \rightarrow N}(\xi) - \xi$ is concave, bigger than its two neighboring points, which are equal to h_N , so

$$h_N = |x_N - x_{N-1}| \leq |y_{i \rightarrow N}(\xi) - \xi| \leq |x_{N+1} - x_{N-1}| = 2h_N$$

So we have

$$|y_{i \rightarrow N}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_N^\theta - x_i|^{1-\alpha}$$

While

$$y_{i \rightarrow N-1}^{1-1/r}(\xi) - \xi^{1-1/r} \leq (y_{i \rightarrow N-1}(\xi) - \xi) \xi^{-1/r}$$

and

$$|\frac{h_N}{r Z_1} - \xi^{1-1/r}| \leq \max\{|\frac{h_N}{r Z_1} - x_{i-1}^{1-1/r}|, |\frac{h_N}{r Z_1} - x_{i+1}^{1-1/r}|\}$$

$$\leq \max \begin{cases} T^{1-1/r} - x_{i-1}^{1-1/r} \leq |x_N - x_{i-1}| T^{-1/r} \leq C|x_N - x_i| \\ |x_{i+1}^{1-1/r} - x_{N-1}^{1-1/r}| \leq |x_{i+1} - x_{N-1}| x_{N-1}^{-1/r} \leq C|x_N - x_i| \end{cases}$$

So we have

$$(y_{i \rightarrow N}^\theta(\xi) - \xi)' \leq C|y_N^\theta - x_i|$$

$$(|y_{i \rightarrow N}^\theta(\xi) - \xi|^{1-\alpha})' = |y_{i \rightarrow N}^\theta(\xi) - \xi|^{-\alpha} (y_{i \rightarrow N}^\theta(\xi) - \xi)'$$

$$\leq |y_N^\theta - x_i|^{1-\alpha}$$

Finally,

$$(|y_{i \rightarrow N}^\theta(\xi) - \xi|^{1-\alpha})'' = (1-\alpha)|y_{i \rightarrow N}^\theta(\xi) - \xi|^{-\alpha} (y_{i \rightarrow N}^\theta(\xi) - \xi)''$$

$$+ \alpha(\alpha-1)|y_{i \rightarrow N}^\theta(\xi) - \xi|^{-1-\alpha} ((y_{i \rightarrow N}^\theta(\xi) - \xi)')^2 \quad \square$$

$$\leq C(r-1)|y_N^\theta - x_i|^{-\alpha} + C|y_N^\theta - x_i|^{1-\alpha}$$

By the three lemmas above, for $N/2 \leq i \leq N-1$, we have

$$(5.113) \quad D_h^2 P_{i \rightarrow N}^\theta(x_i) = P_{i \rightarrow N}^\theta(\xi) \quad \xi \in [x_{i-1}, x_{i+1}] \quad \square$$

$$\leq C h^3 |y_N^\theta - x_i|^{1-\alpha} + C(r-1)(h^3 |y_N^\theta - x_i|^{-\alpha} + h^2 |y_N^\theta - x_i|^{1-\alpha})$$

6. Proof of Theorem 4.2.

7. Experimental results.

8. Conclusions. Some conclusions here.

Appendix A. Approximate of difference quotients.

LEMMA A.1. *If $g(x)$ is twice differentiable continuous function on open set Ω , there exists $\xi \in [x_{i-1}, x_{i+1}]$ such that*

$$(A.1) \quad D_h^2 g(x_i) := \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = g''(\xi), \quad \xi \in [x_{i-1}, x_{i+1}]$$

$$(A.2) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \int_{x_{i-1}}^{x_i} g''(y)(y - x_{i-1}) dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} g''(y)(x_{i+1} - y) dy \right)$$

And if $g(x) \in C^4(\Omega)$, then

$$(A.3) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = g''(x_i) + \frac{h_{i+1} - h_i}{3} g'''(x_i) + \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 g''''(\eta_1) + h_{i+1}^3 g''''(\eta_2))$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$.

Proof.

$$g(x_{i-1}) = g(x_i) - (x_i - x_{i-1})g'(x_i) + \frac{(x_i - x_{i-1})^2}{2} g''(\xi_1), \quad \xi_1 \in [x_{i-1}, x_i]$$

$$g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i)g'(x_i) + \frac{(x_{i+1} - x_i)^2}{2} g''(\xi_2), \quad \xi_2 \in [x_i, x_{i+1}]$$

Substitute them in the left side of (A.1), we have

$$\frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = \frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2)$$

Now, using intermediate value theorem, there exists $\xi \in [\xi_1, \xi_2]$ such that

$$\frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2) = g''(\xi)$$

For the second equation, similarly

$$g(x_{i-1}) = g(x_i) - (x_i - x_{i-1})g'(x_i) + \int_{x_{i-1}}^{x_i} g''(y)(y - x_{i-1}) dy$$

$$g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i)g'(x_i) + \int_{x_i}^{x_{i+1}} g''(y)(x_{i+1} - y)dy$$

And the last equation can be obtained by

$$g(x_{i-1}) = g(x_i) - h_i g'(x_i) + \frac{h_i^2}{2} g''(x_i) - \frac{h_i^3}{3!} g'''(x_i) + \frac{h_i^4}{4!} g''''(\eta_1)$$

$$g(x_{i+1}) = g(x_i) + h_{i+1} g'(x_i) + \frac{h_{i+1}^2}{2} g''(x_i) + \frac{h_{i+1}^3}{3!} g'''(x_i) + \frac{h_{i+1}^4}{4!} g''''(\eta_2)$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$. Expecially,

$$\begin{aligned} \frac{h_i^4}{4!} g''''(\eta_1) &= \int_{x_{i-1}}^{x_i} g''''(y) \frac{(y - x_{i-1})^3}{3!} dy \\ \frac{h_{i+1}^4}{4!} g''''(\eta_2) &= \int_{x_i}^{x_{i+1}} g''''(y) \frac{(x_{i+1} - y)^3}{3!} dy \end{aligned}$$

Substitute them to the left side of (A.3), we can get the result. \square

LEMMA A.2. If $y \in [x_{j-1}, x_j]$, denote $y = \theta x_{j-1} + (1 - \theta)x_j$, $\theta \in [0, 1]$,

$$u(y_j^\theta) - u_h(y_j^\theta) = -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [x_{j-1}, x_j]$$

(A.6)

$$u(y_j^\theta) - u_h(y_j^\theta) = -\frac{\theta(1-\theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1-\theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1-\theta)^2 u'''(\eta_2))$$

where $\eta_1 \in [x_{j-1}, y_j^\theta]$, $\eta_2 \in [y_j^\theta, x_j]$.

Proof. By Taylor expansion, we have

$$u(x_{j-1}) = u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(\xi_1), \quad \xi_1 \in [x_{j-1}, y_j^\theta]$$

$$u(x_j) = u(y_j^\theta) + (1-\theta) h_j u'(y_j^\theta) + \frac{(1-\theta)^2 h_j^2}{2!} u''(\xi_2), \quad \xi_2 \in [y_j^\theta, x_j]$$

Thus

$$\begin{aligned} u(y_j^\theta) - u_h(y_j^\theta) &= u(y_j^\theta) - (1-\theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1-\theta)}{2} h_j^2 (\theta u''(\xi_1) + (1-\theta)u''(\xi_2)) \\ &= -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [\xi_1, \xi_2] \end{aligned}$$

The second equation is similar,

$$u(x_{j-1}) = u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(y_j^\theta) - \frac{\theta^3 h_j^3}{3!} u'''(\eta_1)$$

$$u(x_j) = u(y_j^\theta) + (1-\theta) h_j u'(y_j^\theta) + \frac{(1-\theta)^2 h_j^2}{2!} u''(y_j^\theta) + \frac{(1-\theta)^3 h_j^3}{3!} u'''(\eta_2)$$

where $\eta_1 \in [x_{j-1}, y_j^\theta]$, $\eta_2 \in [y_j^\theta, x_j]$. Thus \square

$$u(y_j^\theta) - u_h(y_j^\theta) = u(y_j^\theta) - (1-\theta)u(x_{j-1}) - \theta u(x_j)$$

$$= -\frac{\theta(1-\theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1-\theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1-\theta)^2 u'''(\eta_2))$$

LEMMA A.3. For $x \in [x_{j-1}, x_j]$

$$(A.7) \quad |u(x) - u_h(x)| = \left| \frac{x_j - x}{h_j} \int_{x_{j-1}}^x u'(y) dy - \frac{x - x_{j-1}}{h_j} \int_x^{x_j} u'(y) dy \right| \\ \leq \int_{x_{j-1}}^{x_j} |u'(y)| dy$$

If $x \in [0, x_1]$, with Corollary 2.4, we have

$$(A.8) \quad |u(x) - u_h(x)| \leq \int_0^{x_1} |u'(y)| dy \leq \int_0^{x_1} C y^{\alpha/2-1} dy \leq C \frac{2}{\alpha} x_1^{\alpha/2}$$

Similarly, if $x \in [x_{2N-1}, 1]$, we have

$$(A.9) \quad |u(x) - u_h(x)| \leq C \frac{2}{\alpha} (2T - x_{2N-1})^{\alpha/2} = C \frac{2}{\alpha} x_1^{\alpha/2}$$

Appendix B. Inequality.

LEMMA B.1.

$$(B.1) \quad h_i \leq rT^{1/r} h \begin{cases} x_i^{1-1/r}, & 1 \leq i \leq N \\ (2T - x_{i-1})^{1-1/r}, & N < i \leq 2N-1 \end{cases}$$

$$(B.2) \quad h_i \geq rT^{1/r} h \begin{cases} x_{i-1}^{1-1/r}, & 1 \leq i \leq N \\ (2T - x_i)^{1-1/r}, & N < i \leq 2N-1 \end{cases}$$

Proof. For $1 \leq i \leq N$,

$$h_i = T \left(\left(\frac{i}{N} \right)^r - \left(\frac{i-1}{N} \right)^r \right) \\ \leq rT \frac{1}{N} \left(\frac{i}{N} \right)^{r-1} = rT^{1/r} h x_i^{1-1/r}$$

$$h_i \geq rT \frac{1}{N} \left(\frac{i-1}{N} \right)^{r-1} = rT^{1/r} h x_{i-1}^{1-1/r}$$

For $N < i \leq 2N$,

$$h_i = T \left(\left(\frac{2N-i+1}{N} \right)^r - \left(\frac{2N-i}{N} \right)^r \right) \\ \leq rT \frac{1}{N} \left(\frac{2N-i+1}{N} \right)^{r-1} = rT^{1/r} h (2T - x_{i-1})^{1-1/r}$$

$$h_i \geq rT \frac{1}{N} \left(\frac{2N-i}{N} \right)^{r-1} = rT^{1/r} h (2T - x_i)^{1-1/r}$$

□

409 LEMMA B.2. *There is a constant $C = 2^{|r-2|}r(r-1)T^{2/r}$ such that for all $i \in$*
 410 $\{1, 2, \dots, 2N-1\}$

$$411 \quad (B.3) \quad |h_{i+1} - h_i| \leq Ch^2 \begin{cases} x_i^{1-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{1-2/r}, & N < i \leq 2N-1 \end{cases}$$

Proof.

$$412 \quad h_{i+1} - h_i = \begin{cases} T \left(\left(\frac{i+1}{N} \right)^r - 2 \left(\frac{i}{N} \right)^r + \left(\frac{i-1}{N} \right)^r \right), & 1 \leq i \leq N-1 \\ 0, & i = N \\ -T \left(\left(\frac{2N-i-1}{N} \right)^r - 2 \left(\frac{2N-i}{N} \right)^r + \left(\frac{2N-i+1}{N} \right)^r \right), & N+1 \leq i \leq 2N-1 \end{cases}$$

413 For $i = 1$,

$$414 \quad h_2 - h_1 = T(2^r - 2) \left(\frac{1}{N} \right)^r = (2^r - 2)T^{2/r}h^2x_1^{1-2/r}$$

415 For $2 \leq i \leq N-1$,

$$416 \quad h_{i+1} - h_i = r(r-1)T N^{-2}\eta^{r-2}, \quad \eta \in \left[\frac{i-1}{N}, \frac{i+1}{N} \right]$$

417 If $r \in [1, 2]$,

$$\begin{aligned} 418 \quad h_{i+1} - h_i &= r(r-1)T N^{-2}\eta^{r-2} \leq r(r-1)T h^2 \left(\frac{i-1}{N} \right)^{r-2} \\ &\leq r(r-1)T h^2 2^{2-r} \left(\frac{i}{N} \right)^{r-2} \\ &= 2^{2-r}r(r-1)T^{2/r}h^2x_i^{1-2/r} \end{aligned}$$

419 else if $r > 2$,

$$\begin{aligned} 420 \quad h_{i+1} - h_i &= r(r-1)T N^{-2}\eta^{r-2} \leq r(r-1)T h^2 \left(\frac{i+1}{N} \right)^{r-2} \\ &\leq r(r-1)T h^2 2^{r-2} \left(\frac{i}{N} \right)^{r-2} \\ &= 2^{r-2}r(r-1)T^{2/r}h^2x_i^{1-2/r} \end{aligned}$$

421 Since

$$422 \quad 2^r - 2 \leq 2^{|r-2|}r(r-1), \quad r \geq 1$$

423 we have

$$424 \quad h_{i+1} - h_i \leq 2^{|r-2|}r(r-1)T^{2/r}h^2x_i^{1-2/r}, \quad 1 \leq i \leq N-1$$

425 For $i = N$, $h_{N+1} - h_N = 0$. For $N < i \leq 2N-1$, it's central symmetric to the first
 426 half of the proof, which is

$$427 \quad h_i - h_{i+1} \leq 2^{|r-2|}r(r-1)T^{2/r}h^2(2T - x_i)^{1-2/r}$$

428 Summarizes the inequalities, we can get

$$429 \quad (B.4) \quad |h_{i+1} - h_i| \leq 2^{|r-2|} r(r-1) T^{2/r} h^2 \begin{cases} x_i^{1-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{1-2/r}, & N < i \leq 2N-1 \end{cases} \quad \square$$

430 **Appendix C. Proofs of some technical details.**

431 *Additional proof of Theorem 5.1.* For $2 \leq i \leq N-1$,

$$\begin{aligned} & \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \\ 432 \quad & \leq C \frac{2}{h_i + h_{i+1}} (h_i^3 x_{i-1}^{-2-\alpha/2} + h_{i+1}^3 x_i^{-2-\alpha/2}) \\ & \leq 2C (h_i^2 x_{i-1}^{-2-\alpha/2} + h_{i+1}^2 x_i^{-2-\alpha/2}) \end{aligned}$$

433 Since Lemma B.1, we have

$$\begin{aligned} 434 \quad & h_i \leq r T^{1/r} h x_i^{1-1/r}, \quad 1 \leq i \leq N \\ 435 \quad & h_{i+1} \leq r T^{1/r} h x_{i+1}^{1-1/r}, \quad 1 \leq i \leq N-1 \end{aligned}$$

436 and

$$\begin{aligned} 437 \quad & x_{i-1}^{-2-\alpha/2} \leq 2^{-r(-2-\alpha/2)} x_i^{-2-\alpha/2} \quad 2 \leq i \leq N-1 \\ 438 \quad & x_{i+1}^{1-1/r} \leq 2^{r-1} x_i^{1-1/r} \quad 1 \leq i \leq N-1 \end{aligned}$$

439 So there is a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{\alpha/2})$ such that

$$440 \quad \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \leq C h^2 x_i^{-\alpha/2-2/r}, \quad 2 \leq i \leq N-1$$

441 For $i = 1$, by (A.4)

$$\begin{aligned} & \frac{1}{4!} \frac{2}{h_1 + h_2} (h_1^3 f''(\eta_1) + h_2^3 f''(\eta_2)) \\ 442 \quad & = \frac{2}{h_1 + h_2} \left(\frac{1}{h_1} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy + \frac{1}{4!} h_2^3 f''(\eta_2) \right) \end{aligned}$$

443 We have proved above that

$$444 \quad \frac{2}{h_1 + h_2} h_2^3 f''(\eta_2) \leq C h^2 x_1^{-\alpha/2-2/r}$$

445 and we can get

$$\begin{aligned} & \int_0^{x_1} f''(y) \frac{y^3}{3!} dy \leq C \frac{1}{3!} \int_0^{x_1} y^{1-\alpha/2} dy \\ 446 \quad & = C \frac{1}{3!(2-\alpha/2)} x_1^{2-\alpha/2} \end{aligned}$$

447 so

$$448 \quad \frac{2}{h_1 + h_2} \frac{1}{h_1} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy = \frac{C 2^{1-r}}{3!(2-\alpha/2)} x_1^{-\alpha/2} = \frac{C 2^{1-r}}{3!(2-\alpha/2)} T^{2/r} h^2 x_1^{-\alpha/2-2/r}$$

And for $i = N$, we have

$$\begin{aligned} & \frac{2}{h_N + h_{N+1}} (h_N^3 f''(\eta_1) + h_{N+1}^3 f''(\eta_2)) \\ &= h_N^2 (f''(\eta_1) + f''(\eta_2)) \\ &\leq r^2 T^{2/r} h^2 x_N^{2-2/r} 2C x_{N-1}^{-2-\alpha/2} \\ &\leq 2r^2 T^{2/r} C 2^{-r(-2-\alpha/2)} h^2 x_N^{-\alpha/2-2/r} \end{aligned}$$

Finally, $N + 1 \leq i \leq 2N - 1$ is symmetric to the first half of the proof, so we can conclude that \square

$$\frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \leq C h^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T - x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N - 1 \end{cases}$$

LEMMA C.1. *There is a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ for $2 \leq j \leq N$, if $y \in [x_{j-1}, x_j]$,*

$$(C.1) \quad |u(y) - u_h(y)| \leq C h^2 y^{\alpha/2-2/r}$$

Proof. For $2 \leq j \leq N$, we have

$$x_j \leq 2^r y, \quad x_{j-1} \geq 2^{-r} y$$

And by Lemma A.2, Lemma B.1 and Corollary 2.4, we have

$$\begin{aligned} u(y) - u_h(y) &= -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [x_{j-1}, x_j] \\ &\leq \frac{\|u\|_{\beta+\alpha}^{(-\alpha/2)}}{4} r^2 T^{2/r} h^2 x_j^{2-2/r} x_{j-1}^{\alpha/2-2} \\ &\leq C h^2 2^{2r-2} y^{2-2/r} 2^{-r(\alpha/2-2)} y^{\alpha/2-2} \\ &= C 2^{-r\alpha/2+4r-2} h^2 y^{\alpha/2-2/r} \end{aligned}$$

symmetricly, for $N < j \leq 2N - 1$, we have

$$(C.2) \quad |u(y) - u_h(y)| \leq C h^2 (2T - y)^{\alpha/2-2/r} \quad \square$$

LEMMA C.2. *There is a constant $C = C(\alpha, r)$ such that for all $1 \leq i < N/2$, $\max\{2i + 1, i + 3\} \leq j \leq 2N$ and $y \in [x_{j-1}, x_j]$, we have*

$$(C.3) \quad D_h^2 \left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)} \right) (x_i) \leq C \frac{y^{-1-\alpha}}{\Gamma(-\alpha)}$$

Proof. Since $y \geq x_{j-1} > x_{i+1}$, by Lemma A.1, if $j - 1 > i + 1$ \square

$$\begin{aligned} D_h^2 \left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)} \right) (x_i) &= \frac{|y - \xi|^{-1-\alpha}}{\Gamma(-\alpha)}, \quad \xi \in [x_{i-1}, x_{i+1}] \\ &\leq \frac{(y - x_{i+1})^{-1-\alpha}}{\Gamma(-\alpha)} \\ &\leq \left(1 - \left(\frac{2}{3}\right)^r\right)^{-1-\alpha} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} \end{aligned}$$

LEMMA C.3. *There is a constant $C = C(\alpha, r)$ such that for all $3 \leq i < N/2, k = \lceil \frac{i}{2} \rceil, 1 \leq j \leq k-1$ and $y \in [x_{j-1}, x_j]$, we have*

$$(C.4) \quad D_h^2\left(\frac{|\cdot - y|^{1-\alpha}}{\Gamma(2-\alpha)}\right)(x_i) \leq C \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)}$$

Proof. Since $y \leq x_j < x_{i-1}$, by Lemma A.1,

$$\begin{aligned} D_h^2\left(\frac{|\cdot - y|^{1-\alpha}}{\Gamma(2-\alpha)}\right)(x_i) &= \frac{|\xi - y|^{-1-\alpha}}{\Gamma(-\alpha)}, \quad \xi \in [x_{i-1}, x_{i+1}] \\ &\leq \frac{(x_{i-1} - x_j)^{-1-\alpha}}{\Gamma(-\alpha)} \leq \frac{(x_{i-1} - x_{k-1})^{-1-\alpha}}{\Gamma(-\alpha)} \\ &\leq \left(\left(\frac{2}{3}\right)^r - \left(\frac{1}{2}\right)^r\right)^{-1-\alpha} \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)} \end{aligned}$$

473

LEMMA C.4. *While $0 \leq i < N/2$, By Lemma A.3*

$$\begin{aligned} |T_{i1}| &\leq C \int_0^{x_1} x_1^{\alpha/2} \frac{|x_i - y|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\ (C.5) \quad &= C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2} |x_i^{2-\alpha} - |x_i - x_1|^{2-\alpha}| \\ &\leq C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2+2-\alpha} = C \frac{1}{\Gamma(3-\alpha)} x_1^{2-\alpha/2} \quad 0 < 2-\alpha < 1 \end{aligned}$$

For $2 \leq j \leq N$, by Lemma A.2 and Corollary 2.4

$$\begin{aligned} |T_{ij}| &\leq \frac{C}{4} \int_{x_{j-1}}^{x_j} h_j^2 x_{j-1}^{\alpha/2-2} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\ (C.6) \quad &\leq \frac{C}{4\Gamma(3-\alpha)} h_j^2 x_{j-1}^{\alpha/2-2} ||x_j - x_i|^{2-\alpha} - |x_{j-1} - x_i|^{2-\alpha}| \end{aligned}$$

LEMMA C.5. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that*

$$(C.7) \quad \sum_{j=1}^3 S_{1j} \leq Ch^2 x_1^{-\alpha/2-2/r}$$

480

$$(C.8) \quad \sum_{j=1}^4 S_{2j} \leq Ch^2 x_2^{-\alpha/2-2/r}$$

482

Proof.

$$S_{1j} = \frac{2}{x_2} \left(\frac{1}{x_1} T_{0j} - \left(\frac{1}{x_1} + \frac{1}{h_2} \right) T_{1j} + \frac{1}{h_2} T_{2j} \right)$$

So, by Lemma C.4

$$S_{11} \leq \frac{2}{x_2 x_1} 4 \frac{C}{\Gamma(3-\alpha)} x_1^{2-\alpha/2} \leq C x_1^{-\alpha/2}$$

485

$$S_{12} \leq \frac{2}{x_2 x_1} \frac{C}{4\Gamma(3-\alpha)} h_2^2 x_1^{\alpha/2-2} (x_2^{2-\alpha} + 2h_2^{2-\alpha} + h_2^{2-\alpha}) \leq C x_1^{-\alpha/2}$$

$$S_{13} \leq \frac{2}{x_2 x_1} \frac{C}{4\Gamma(3-\alpha)} h_3^2 x_2^{\alpha/2-2} (x_3^{2-\alpha} + 2h_3^{2-\alpha} + h_3^{2-\alpha}) \leq C x_1^{-\alpha/2}$$

But

$$x_1^{-\alpha/2} = T^{2/r} h^2 x_1^{-\alpha/2-2/r}$$

For $i = 2$, Sorry □

LEMMA C.6. *There exists a constant $C = C(T, r, l)$ such that For $3 \leq i \leq N - 1$, $k + 1 = \lceil \frac{i}{2} \rceil$, $k \leq j \leq \min\{2i - 1, N - 1\}$, $l = 3, 4$, when $\xi \in [x_{i-1}, x_{i+1}]$,*

$$(C.9) \quad (h_{j-i}^3(\xi))' \leq (r-1) C h^2 x_i^{1-2/r} h_j$$

$$(C.10) \quad (h_{j-i}^4(\xi))' \leq (r-1) C h^2 x_i^{1-2/r} h_j^2$$

Proof. From (5.32)

$$(C.11) \quad y'_{j-i}(x) = y_{j-i}^{1-1/r}(x) x^{1/r-1}$$

$$(C.12) \quad y''_{j-i}(x) = \frac{1-r}{r} y_{j-i}^{1-2/r}(x) x^{1/r-2} Z_{j-i}$$

for $l = 3, 4$, by (5.34)

$$(C.13) \quad \begin{aligned} (h_{j-i}^l(\xi))' &= l h_{j-i}^{l-1}(\xi) (y'_{j-i}(\xi) - y'_{j-i-1}(\xi)) \\ &= l h_{j-i}^{l-1}(\xi) \xi^{1/r-1} (y_{j-i}^{1-1/r}(\xi) - y_{j-i-1}^{1-1/r}(\xi)) \geq 0 \end{aligned}$$

For $\xi \in [x_{i-1}, x_{i+1}]$ and $2 \leq k \leq j \leq \min\{2i - 1, N - 1\}$, using Lemma B.1

$$\begin{aligned} h_{j-i}(\xi) &\leq h_{j-i}(x_{i+1}) = h_{j+1} \\ &\leq r T^{1/r} h x_{j+1}^{1-1/r} \leq r T^{1/r} 2^{r-1} h x_i^{1-1/r} \end{aligned}$$

And

$$(C.14) \quad 2^{-r} x_i \leq x_{i-1} \leq \xi \leq x_{i+1} \leq 2^r x_i$$

We have

$$(C.15) \quad \xi^{1/r-m} \leq 2^{|mr-1|} x_i^{1/r-m}, \quad m = 1, 2$$

but

$$(C.16) \quad \begin{aligned} y_{j-i}^{1-1/r}(\xi) - y_{j-i-1}^{1-1/r}(\xi) &= (\xi^{1/r} + Z_{j-i})^{r-1} - (\xi^{1/r} + Z_{j-i-1})^{r-1} \\ &= (r-1) Z_1 (\xi^{1/r} + Z_{j-i-\gamma})^{r-2}, \quad \gamma \in [0, 1] \\ &= (r-1) T^{1/r} h y_{j-i-\gamma}^{1-2/r}(\xi) \end{aligned}$$

And
(C.17)

$$4^{-r}x_i \leq x_{\lceil \frac{i}{2} \rceil - 1} \leq x_{j-2} = y_{j-i-1}(x_{i-1}) \leq y_{j-i-\gamma}(\xi) \leq y_{j-i}(x_{i+1}) = x_{j+1} \leq x_{2i} \leq 2^r x_i$$

Therefore,

$$(C.18) \quad y_{j-i-\gamma}^{1-2/r}(\xi) \leq 2^{2|r-2|} x_i^{1-2/r}$$

So we can get

$$(C.19) \quad y'_{j-i}(\xi) - y'_{j-i-1}(\xi) \leq (r-1)C(T, r) h x_i^{-1/r}$$

We get

$$(C.20) \quad (h_{j-i}^l(\xi))' \leq l(r-1)C h_{j+1}^{l-1} h x_i^{-1/r}$$

And by Lemma B.1,

$$(C.21) \quad h_{j+1} \leq rTh \left(\frac{j+1}{N} \right)^{r-1} \leq rTh 2^{r-1} \left(\frac{j-1}{N} \right) = 2^{r-1} h_j$$

$$(C.22) \quad h_{j+1} \leq rT^{1/r} h x_{j+1}^{1-1/r} \leq rT^{1/r} h x_{2i}^{1-1/r} \leq rT^{1/r} 2^{r-1} h x_i^{1-1/r}$$

We can get

$$(C.23) \quad \begin{aligned} (h_{j-i}^l(\xi))' &\leq l(r-1)C h_j^{l-2} h_{j+1} h x_i^{-1/r} \\ &\leq l(r-1)C h h_j^{l-2} (h x_i^{1-1/r}) x_i^{-1/r} \\ &= (r-1)C h^2 x_i^{1-2/r} h_j^{l-2} \end{aligned}$$

Meanwhile, we can get

$$(C.24) \quad h_{j-i}^3(\xi) \leq h_{j+1}^3 \leq C h^2 x_i^{2-2/r} h_j$$

$$(C.25) \quad h_{j-i}^4(\xi) \leq h_{j+1}^4 \leq C h^2 x_i^{2-2/r} h_j^2$$

□

LEMMA C.7. *There exists a constant $C = C(T, r, l)$ such that For $3 \leq i \leq N - 1$, $\lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i - 1, N - 1\}$, when $\xi \in [x_{i-1}, x_{i+1}]$,*

$$(C.26) \quad (h_{j-i}^3(\xi))'' \leq C(r-1) h^2 x_i^{-2/r} h_j$$

Proof. From (C.11)

$$(C.27) \quad \begin{aligned} (h_{j-i}^3(\xi))'' &= 6h_{j-i}(\xi)(y'_{j-i}(\xi) - y'_{j-i-1}(\xi))^2 + 3h_{j-i}^2(\xi)(y''_{j-i}(\xi) - y''_{j-i-1}(\xi)) \\ &= 6h_{j-i}(\xi)\xi^{1/r-1}(y_{j-i}^{1-1/r}(\xi) - y_{j-i-1}^{1-1/r}(\xi)) \\ &\quad + 3\frac{1-r}{r}h_{j-i}^2(\xi)\xi^{1/r-2}(y_{j-i}^{1-2/r}(\xi)Z_{j-i} - y_{j-i-1}^{1-2/r}(\xi)Z_{j-i-1}) \end{aligned}$$

537 Using the inequalities of the proof of Lemma C.6

$$\begin{aligned}
 & 6h_{j-i}(\xi)(y'_{j-i}(\xi) - y'_{j-i-1}(\xi))^2 \\
 538 \quad (C.28) \quad & \leq 6h_{j+1}((r-1)Chx_i^{-1/r})^2 \\
 & \leq C(r-1)^2 h^2 x_i^{-2/r} h_j
 \end{aligned}$$

539 For the second partial

$$\begin{aligned}
 & h_{j-i}^2(\xi)\xi^{1/r-2}(y_{j-i}^{1-2/r}(\xi)Z_{j-i} - y_{j-i-1}^{1-2/r}(\xi)Z_{j-i-1}) \\
 540 \quad (C.29) \quad & \leq Ch_{j+1}^2 x_i^{1/r-2}((y_{j-i}^{1-2/r}(\xi) - y_{j-i-1}^{1-2/r}(\xi))Z_{j-i} + y_{j-i-1}^{1-2/r}(\xi)Z_1)
 \end{aligned}$$

541 but

$$\begin{aligned}
 & y_{j-i}^{1-2/r}(\xi) - y_{j-i-1}^{1-2/r}(\xi) = (\xi^{1/r} + Z_{j-i})^{r-2} - (\xi^{1/r} + Z_{j-i-1})^{r-2} \\
 & = (r-2)Z_1(\xi^{1/r} + Z_{j-i-\gamma})^{r-3} \\
 542 \quad (C.30) \quad & = (r-2)T^{-r}hy_{j-i-\gamma}^{1-3/r}(\xi) \\
 & \leq C(r-2)hx_i^{1-3/r}
 \end{aligned}$$

543 So we can get

$$\begin{aligned}
 & h_{j-i}^2(\xi)\xi^{1/r-2}(y_{j-i}^{1-2/r}(\xi)Z_{j-i} - y_{j-i-1}^{1-2/r}(\xi)Z_{j-i-1}) \\
 & \leq Ch_jhx_i^{1-1/r}x_i^{1/r-2}(C(r-2)hx_i^{1-3/r}Z_{j-i} + Cx_i^{1-2/r}T^{1/r}h) \\
 544 \quad (C.31) \quad & \leq Ch^2((r-2)x_i^{-3/r}x_{|j-i|}^{1/r} + x_i^{-2/r})h_j \\
 & \leq Ch^2x_i^{-2/r}h_j
 \end{aligned}$$

545 Summarizes, we have

$$546 \quad (C.32) \quad (h_{j-i}^3(\xi))'' \leq C(r-1)h^2x_i^{-2/r}h_j \quad \square$$

547 *proof of Lemma 5.16.* From (5.32)

$$548 \quad (C.33) \quad y'_{j-i}(x) = y_{j-i}^{1-1/r}(x)x^{1/r-1}$$

$$549 \quad (C.34) \quad y''_{j-i}(x) = \frac{1-r}{r}y_{j-i}^{1-2/r}(x)x^{1/r-2}Z_{j-i}$$

550 Since

$$551 \quad x_{j-2} \leq y_{j-i-1}(x_{i-1}) \leq y_{j-i}^\theta(\xi) \leq y_{j-i-1}^\theta(x_{i+1}) \leq x_{j+1}$$

552 We have known (C.17)

$$553 \quad (C.35) \quad u''(y_{j-i}^\theta(\xi)) \leq C(y_{j-i}^\theta(\xi))^{\alpha/2-2} \leq Cx_{j-2}^{\alpha/2-2} \leq Cx_{\lfloor \frac{i}{2} \rfloor -1}^{\alpha/2-2} \leq C4^{r(2-\alpha/2)}x_i^{\alpha/2-2}$$

554

$$\begin{aligned}
 & (u''(y_{j-i}^\theta(\xi)))' = u'''(y_{j-i}^\theta(\xi))y_{j-i}^{\theta'}(\xi) \\
 555 \quad (C.36) \quad & \leq Cx_i^{\alpha/2-3}\xi^{1/r-1}y_{j-i}^{1-1/r}(\xi) \\
 & \leq Cx_i^{\alpha/2-3}x_i^{1/r-1}x_i^{1-1/r} = Cx_i^{\alpha/2-3}
 \end{aligned}$$

556

$$\begin{aligned}
& (u''(y_{j-i}^\theta(\xi)))'' = u''''(y_{j-i}^\theta(\xi))(y_{j-i}^{\theta''}(\xi))^2 + u'''(y_{j-i}^\theta(\xi))y_{j-i}^{\theta'''}(\xi) \\
& \leq Cx_i^{\alpha/2-4} + Cx_i^{\alpha/2-3}\frac{r-1}{r}x_i^{1-2/r}x_i^{1/r-2}Z_{|j-i|+1} \\
& \leq Cx_i^{\alpha/2-4} + C\frac{r-1}{r}x_i^{\alpha/2-3}x_i^{-1/r}x_i^{1/r} \\
& = Cx_i^{\alpha/2-4}
\end{aligned}
\tag{C.37}$$

□

Proof of Lemma 5.17.

$$\begin{aligned}
& |y_{j-i}^\theta(\xi) - \xi| = |\theta(y_{j-i-1}(\xi) - \xi) + (1-\theta)(y_{j-i}(\xi) - \xi)| \\
& = \theta|y_{j-i-1}(\xi) - \xi| + (1-\theta)|y_{j-i}(\xi) - \xi|
\end{aligned}
\tag{C.38}$$

Since $|y_{j-i}(\xi) - \xi|$ is increasing about ξ , we have

$$\left(\frac{i-1}{i}\right)^r |x_j - x_i| \leq |x_{j-1} - x_{i-1}| \leq |y_{j-i}(\xi) - \xi| \leq |x_{j+1} - x_{i+1}| \leq \left(\frac{i+1}{i}\right)^r |x_j - x_i|
\tag{C.39}$$

Thus,

$$\left(\frac{2}{3}\right)^r |y_j^\theta - x_i| \leq |y_{j-i}^\theta(\xi) - \xi| \leq \left(\frac{3}{4}\right)^r (\theta|x_j - x_i| + (1-\theta)|x_{j-1} - x_i|) = \left(\frac{3}{4}\right)^r |y_j^\theta - x_i|
\tag{C.40}$$

563

$$|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_j^\theta - x_i|^{1-\alpha}
\tag{C.41}$$

Next,

$$\begin{aligned}
& (|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})' = (1-\alpha)|y_{j-i}^\theta(\xi) - \xi|^{-\alpha}\xi^{1/r-1}(\theta y_{j-i-1}^{1-1/r}(\xi) + (1-\theta)y_{j-i}^{1-1/r}(\xi)) - 1| \\
& \leq C|y_j^\theta - x_i|^{-\alpha}\xi^{1/r-1}|\theta y_{j-i-1}^{1-1/r}(\xi) + (1-\theta)y_{j-i}^{1-1/r}(\xi) - \xi^{1-1/r}|
\end{aligned}
\tag{C.42}$$

Similar with (C.40), we have

$$\begin{aligned}
& |y_{j-i}^{1-1/r}(\xi) - \xi^{1-1/r}| \leq C|x_j^{1-1/r} - x_i^{1-1/r}| \\
& \leq C|x_j - x_i|x_i^{-1/r}
\end{aligned}
\tag{C.43}$$

So we can get

$$\begin{aligned}
& |\theta y_{j-i-1}^{1-1/r}(\xi) + (1-\theta)y_{j-i}^{1-1/r}(\xi) - \xi^{1-1/r}| \\
& \leq Cx_i^{-1/r}(\theta|x_{j-1} - x_i| + (1-\theta)|x_j - x_i|) \\
& = Cx_i^{-1/r}|y_j^\theta - x_i|
\end{aligned}
\tag{C.44}$$

Combine them, we get

$$\begin{aligned}
& (|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})' \leq C|y_j^\theta - x_i|^{-\alpha}x_i^{1/r-1}x_i^{-1/r}|y_j^\theta - x_i| \\
& = C|y_j^\theta - x_i|^{1-\alpha}x_i^{-1}
\end{aligned}
\tag{C.45}$$

Finally, we have

$$\begin{aligned}
& (|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'' = \alpha(\alpha-1)|y_{j-i}^\theta(\xi) - \xi|^{-\alpha-1}(\xi^{1/r-1}(\theta y_{j-i-1}^{1-1/r}(\xi) + (1-\theta)y_{j-i}^{1-1/r}(\xi)) - 1)^2 \\
& + (1-\alpha)|y_{j-i}^\theta(\xi) - \xi|^{-\alpha}\frac{1-r}{r}\xi^{1/r-2}|\theta y_{j-i-1}^{1-2/r}(\xi)Z_{j-i-1} + (1-\theta)y_{j-i}^{1-2/r}(\xi)Z_{j-i}|
\end{aligned}
\tag{C.46}$$

575 Using the inequalities above ,we have

$$\begin{aligned}
 & |y_{j-i}^\theta(\xi) - \xi|^{-\alpha-1}(\xi^{1/r-1}(\theta y_{j-i-1}^{1-1/r}(\xi) + (1-\theta)y_{j-i}^{1-1/r}(\xi)) - 1)^2 \\
 576 \quad (C.47) \quad & \leq C|y_j^\theta - x_i|^{-\alpha-1}(x_i^{-1}|y_j^\theta - x_i|)^2 \\
 & = C|y_j^\theta - x_i|^{1-\alpha}x_i^{-2}
 \end{aligned}$$

577 And by

$$578 \quad (C.48) \quad |Z_{j-i}| = |x_j^{1/r} - x_i^{1/r}| \leq |x_j - x_i|x_i^{1/r-1}$$

579 we have

$$\begin{aligned}
 & |y_{j-i}^\theta(\xi) - \xi|^{-\alpha}\xi^{1/r-2}|\theta y_{j-i-1}^{1-2/r}(\xi)Z_{j-i-1} + (1-\theta)y_{j-i}^{1-2/r}(\xi)Z_{j-i}| \\
 580 \quad (C.49) \quad & \leq C|y_j^\theta - x_i|^{-\alpha}x_i^{1/r-2}x_i^{1-2/r}|\theta Z_{j-i-1} + (1-\theta)Z_{j-i}| \\
 & \leq C|y_j^\theta - x_i|^{-\alpha}x_i^{-2}|y_j^\theta - x_i| \\
 & = C|y_j^\theta - x_i|^{1-\alpha}x_i^{-2}
 \end{aligned}$$

□

581 *proof of Lemma 5.19.* For $k \leq j < \min\{2i-1, N-1\}$

$$\begin{aligned}
 & \frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \\
 582 \quad (C.50) \quad & \frac{Q_{j-i}^\theta(x_{i+1}) - Q_{j-i}^\theta(x_i)}{h_{i+1}}u'''(\eta_{j+1}^\theta) + Q_{j-i}^\theta(x_i)\frac{u'''(\eta_{j+1}^\theta) - u'''(\eta_j^\theta)}{h_{i+1}} \\
 & \leq Q_{j-i}^{\theta'}(\xi)Cx_j^{\alpha/2-3} + Q_{j-i}^\theta(x_i)Cu''''(\eta)\frac{h_i + h_{i+1}}{h_{i+1}}
 \end{aligned}$$

583 where $\xi \in [x_i, x_{i+1}]$, $\eta \in [x_{j-1}, x_{j+1}]$.

584 From (5.36), by Lemma C.6 and Lemma 5.17, we have

$$\begin{aligned}
 & Q_{j-i}^{\theta'}(\xi) \leq Ch^2 \frac{|y_{j+1}^\theta - x_{i+1}|^{1-\alpha}}{\Gamma(2-\alpha)} x_{i+1}^{1-2/r} h_{j+1}^2 \\
 585 \quad (C.51) \quad & \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{1-2/r} h_j^2
 \end{aligned}$$

586 And by defination

$$587 \quad (C.52) \quad Q_{j-i}^\theta(x_i) = h_j^4 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} \leq Ch^2 x_i^{2-2/r} \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} h_j^2$$

588 With , we have

$$589 \quad (C.53) \quad 4^{-r}x_i \leq x_{k-1} \leq x_{j-1} < x_j \leq x_{2i-1} \leq 2^r x_i$$

590 So we have

$$\begin{aligned}
 & \frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \\
 591 \quad (C.54) \quad & \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{1-2/r} h_j^2 x_i^{\alpha/2-3} + Ch^2 x_i^{2-2/r} \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} h_j^2 x_{j-1}^{\alpha/2-4} \\
 & = Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j^2
 \end{aligned}$$

592 while

$$593 \quad h_j \leq h_{2i-1} \leq 2^r h_i$$

594 Subsitute into the inequality above, we get the goal

$$\begin{aligned} & \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \right) \\ 595 \quad (C.55) \quad & \leq \frac{1}{h_i} Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j 2^r h_i \\ & = Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j \end{aligned}$$

596 While, the later is similar. □

597 **Acknowledgments.** We would like to acknowledge the assistance of volunteers
598 in putting together this example manuscript and supplement.

599 REFERENCES

- 600 [1] X. ROS-OTON AND J. SERRA, *The dirichlet problem for the fractional laplacian: Regular-*
601 *ity up to the boundary*, Journal de Mathématiques Pures et Appliquées, 101 (2014),
602 pp. 275–302, <https://doi.org/https://doi.org/10.1016/j.matpur.2013.06.003>, <https://www.sciencedirect.com/science/article/pii/S0021782413000895>.
603