

A SECOND ORDER NUMERICAL METHODS FOR REISZ-FRACTIONAL ELLIPTIC EQUATION ON GRADED MESH*

JIANXING HAN[†] AND MINGHUA CHEN[‡]

Abstract. This is an example SIAM L^AT_EX article. This can be used as a template for new articles. Abstracts must be able to stand alone and so cannot contain citations to the paper's references, equations, etc. An abstract must consist of a single paragraph and be concise. Because of online formatting, abstracts must appear as plain as possible. Any equations should be inline.

Key words. example, L^AT_EX

MSC codes. ?????????????????

1. Introduction. For $\Omega = (0, 2T)$, $1 < \alpha < 2$,

$$(1.1) \quad \begin{cases} (-\Delta)^{\frac{\alpha}{2}} u(x) = f(x), & x \in \Omega \\ u(x) = 0, & x \in \mathbb{R} \setminus \Omega \end{cases}$$

where

$$(1.2) \quad (-\Delta)^{\frac{\alpha}{2}} u(x) = -\frac{\partial^\alpha u}{\partial |x|^\alpha} = -\kappa_\alpha \frac{d^2}{dx^2} \int_\Omega \frac{|x-y|^{1-\alpha}}{\Gamma(2-\alpha)} u(y) dy$$

$$(1.3) \quad \kappa_\alpha = -\frac{1}{2 \cos(\alpha\pi/2)} > 0$$

2. Preliminaries: Numeric scheme and main results.

2.1. Numeric Format.

$$(2.1) \quad x_i = \begin{cases} T \left(\frac{i}{N} \right)^r, & 0 \leq i \leq N \\ 2T - T \left(\frac{2N-i}{N} \right)^r, & N \leq i \leq 2N \end{cases}$$

where $r \geq 1$. And let

$$(2.2) \quad h_j = x_j - x_{j-1}, \quad 1 \leq j \leq 2N$$

Let $\{\phi_j(x)\}_{j=1}^{2N-1}$ be standard hat functions, which are basis of the piecewise linear function space.

$$(2.3) \quad \phi_j(x) = \begin{cases} \frac{1}{h_j}(x - x_{j-1}), & x_{j-1} \leq x \leq x_j \\ \frac{1}{h_{j+1}}(x_{j+1} - x), & x_j \leq x \leq x_{j+1} \\ 0, & \text{otherwise} \end{cases}$$

And then, define the piecewise linear interpolant of the true solution u to be

$$(2.4) \quad \Pi_h u(x) := \sum_{j=1}^{2N-1} u(x_j) \phi_j(x)$$

*Submitted to the editors DATE.

[†]School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, PR China (hanjx2023@mail.lzu.edu.cn).

[‡]School of Mathematics and Statistics, Lanzhou University, Lanzhou 730000, PR China (chen@mail.lzu.edu.cn).

For convenience, we denote

$$(2.5) \quad I^{2-\alpha}u(x) := \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x-y|^{1-\alpha} u(y) dy$$

and

$$(2.6) \quad D_h^2 u(x_i) := \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} u(x_{i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) u(x_i) + \frac{1}{h_{i+1}} u(x_{i+1}) \right)$$

Now, we discretise (1.1) by replacing $u(x)$ by a continuous piecewise linear function

$$(2.7) \quad u_h(x) := \sum_{j=1}^{2N-1} u_j \phi_j(x)$$

whose nodal values u_j are to be determined by collocation at each mesh point x_i for $i = 1, 2, \dots, 2N-1$:

$$(2.8) \quad -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) := -\kappa_{\alpha} D_h^2 I^{2-\alpha} u_h(x_i) = f(x_i) =: f_i$$

Here,

$$(2.9) \quad -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) = \sum_{j=1}^{2N-1} -\kappa_{\alpha} D_h^2 I^{2-\alpha} \phi_j(x_i) u_j = \sum_{j=1}^{2N-1} a_{ij} u_j$$

where

$$(2.10) \quad a_{ij} = -\kappa_{\alpha} D_h^2 I^{2-\alpha} \phi_j(x_i) \quad \text{for } i, j = 1, 2, \dots, 2N-1$$

We have replaced $(-\Delta)^{\alpha/2} u(x_i) = f(x_i)$ in (1.1) by $-\kappa_{\alpha} D_h^{\alpha} u_h(x_i) = f(x_i)$ in (2.8), with truncation error

$$(2.11) \quad \tau_i := -\kappa_{\alpha} \left(D_h^{\alpha} \Pi_h u(x_i) - \frac{d^2}{dx^2} I^{2-\alpha} u(x_i) \right) \quad \text{for } i = 1, 2, \dots, 2N-1$$

where $-\kappa_{\alpha} D_h^{\alpha} \Pi_h u(x_i) = \sum_{j=1}^{2N-1} -\kappa_{\alpha} D_h^{\alpha} \phi_j(x_i) u(x_j) = \sum_{j=1}^{2N-1} a_{ij} u(x_j)$.

The discrete equation (2.8) can be written in matrix form

$$(2.12) \quad AU = F$$

where $A = (a_{ij}) \in \mathbb{R}^{(2N-1) \times (2N-1)}$, $U = (u_1, \dots, u_{2N-1})^T$ is unknown and $F = (f_1, \dots, f_{2N-1})^T$.

We can deduce a_{ij} ,

$$(2.13) \quad \begin{aligned} a_{ij} &= -\kappa_{\alpha} D_h^2 I^{2-\alpha} \phi_j(x_i) \\ &= -\kappa_{\alpha} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \tilde{a}_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{a}_{i,j} + \frac{1}{h_{i+1}} \tilde{a}_{i+1,j} \right) \end{aligned}$$

where

$$(2.14) \quad \begin{aligned} \tilde{a}_{ij} &= I^{2-\alpha} \phi_j(x_i) \\ &= \frac{1}{\Gamma(4-\alpha)} \left(\frac{|x_i - x_{j-1}|^{3-\alpha}}{h_j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) |x_i - x_j|^{3-\alpha} + \frac{|x_i - x_{j+1}|^{3-\alpha}}{h_{j+1}} \right) \end{aligned}$$

2.2. Regularity of the true solution. For any $\beta > 0$, we use the standard notation $C^\beta(\Omega)$, $C^\beta(\mathbb{R})$, etc., for Hölder spaces and their norms and seminorms. When no confusion is possible, we use the notation $C^\beta(\Omega)$ to refer to $C^{k,\beta'}(\Omega)$, where k is the greatest integer such that $k < \beta$ and where $\beta' = \beta - k$. The Hölder spaces $C^{k,\beta'}(\Omega)$ are defined as the subspaces of $C^k(\Omega)$ consisting of functions whose k -th order partial derivatives are locally Hölder continuous[1] with exponent β' in Ω , where $C^k(\Omega)$ is the set of all k -times continuously differentiable functions on open set Ω .

DEFINITION 2.1 (delta dependent norm [2]). ...

THEOREM 2.2. Let $f \in C^\beta(\Omega)$, $\beta > 2$ be such that $\|f\|_\beta^{(\alpha/2)} < \infty$, then for $l = 0, 1, 2$

$$(2.15) \quad |f^{(l)}(x)| \leq \|f\|_\beta^{(\alpha/2)} \begin{cases} x^{-l-\alpha/2}, & \text{if } 0 < x \leq T \\ (2T-x)^{-l-\alpha/2}, & \text{if } T \leq x < 2T \end{cases}$$

THEOREM 2.3 (Regularity up to the boundary [2]). Let Ω be a bounded domain, and $\beta > 0$ be such that neither β nor $\beta + \alpha$ is an integer. Let $f \in C^\beta(\Omega)$ be such that $\|f\|_\beta^{(\alpha/2)} < \infty$, and $u \in C^{\alpha/2}(\mathbb{R}^n)$ be a solution of (1.1). Then, $u \in C^{\beta+\alpha}(\Omega)$ and

$$(2.16) \quad \|u\|_{\beta+\alpha}^{(-\alpha/2)} \leq C \left(\|u\|_{C^{\alpha/2}(\mathbb{R})} + \|f\|_\beta^{(\alpha/2)} \right)$$

COROLLARY 2.4. Let u be a solution of (1.1) where $f \in L^\infty(\Omega)$ and $\|f\|_\beta^{(\alpha/2)} < \infty$. Then, for any $x \in \Omega$ and $l = 0, 1, 2, 3, 4$

$$(2.17) \quad |u^{(l)}(x)| \leq \|u\|_{\beta+\alpha}^{(-\alpha/2)} \begin{cases} x^{\alpha/2-l}, & \text{if } 0 < x \leq T \\ (2T-x)^{\alpha/2-l}, & \text{if } T \leq x < 2T \end{cases}$$

And in this paper bellow, without special instructions, we allways assume that

$$(2.18) \quad f \in L^\infty(\Omega) \cap C^\beta(\Omega) \quad \text{and} \quad \|f\|_\beta^{(\alpha/2)} < \infty, \quad \text{with } \alpha + \beta > 4$$

2.3. Main results. Here we state our main results; the proof is deferred to section 3 and section 4.

Let's denote $h = \frac{1}{N}$, we have

THEOREM 2.5 (Local Truncation Error). If $u(x)$ is a solution of the equation (1.1) where f satisfy the regular condition (2.18), then there exists $C_1(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)}, \|f\|_\beta^{(\alpha/2)})$ and $C_2(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$, such that the truncation error (2.11) satisfies

$$(2.19) \quad \begin{aligned} |\tau_i| &:= |-\kappa_\alpha D_h^\alpha \Pi_h u(x_i) - f(x_i)| \\ &\leq C_1 h^{\min\{\frac{r\alpha}{2}, 2\}} \begin{cases} x_i^{-\alpha}, & 1 \leq i \leq N \\ (2T-x_i)^{-\alpha}, & N < i \leq 2N-1 \end{cases} \\ &\quad + C_2(r-1)h^2 \begin{cases} |T-x_{i-1}|^{1-\alpha}, & 1 \leq i \leq N \\ |T-x_{i+1}|^{1-\alpha}, & N < i \leq 2N-1 \end{cases} \end{aligned}$$

THEOREM 2.6 (Global Error). *The discrete equation (2.8) has solution and there exists a positive constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)}, \|f\|_{\beta}^{(\alpha/2)})$ such that the error between the numerical solution U with the exact solution $u(x_i)$ satisfies*

$$(2.20) \quad \max_{1 \leq i \leq 2N-1} |u_i - u(x_i)| \leq Ch^{\min\{\frac{r\alpha}{2}, 2\}}$$

That means the numerical method has convergence order $\min\{\frac{r\alpha}{2}, 2\}$.

3. Local Truncation Error.

3.1. Proof of Theorem 2.5. The truncation error of the discrete format can be written as

$$(3.1) \quad \begin{aligned} -\kappa_{\alpha} D_h^{\alpha} \Pi_h u(x_i) - f(x_i) &= -\kappa_{\alpha} (D_h^2 I^{2-\alpha} \Pi_h u(x_i) - \frac{d^2}{dx^2} I^{2-\alpha} u(x_i)) \\ &= -\kappa_{\alpha} D_h^2 I^{2-\alpha} (\Pi_h u - u)(x_i) - \kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha} u(x_i) \end{aligned}$$

THEOREM 3.1. *There exists a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$ such that*

$$(3.2) \quad \left| -\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha} u(x_i) \right| \leq Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T - x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N-1 \end{cases}$$

Proof. Since $f \in C^2(\Omega)$ and

$$(3.3) \quad \frac{d^2}{dx^2} (-\kappa_{\alpha} I^{2-\alpha} u(x)) = f(x), \quad x \in \Omega,$$

we have $I^{2-\alpha} u \in C^4(\Omega)$. Therefore, using equation (A.3) of Lemma A.1, for $1 \leq i \leq 2N-1$, we have

$$(3.4) \quad \begin{aligned} -\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha} u(x_i) &= \frac{h_{i+1} - h_i}{3} f'(x_i) \\ &+ \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \int_{x_{i-1}}^{x_i} f''(y) \frac{(y - x_{i-1})^3}{3!} dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} f''(y) \frac{(y - x_{i+1})^3}{3!} dy \right) \end{aligned}$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$. By Lemma B.2 and Theorem 2.2 we have 1.

$$(3.5) \quad \left| \frac{h_{i+1} - h_i}{3} f'(x_i) \right| \leq \frac{C(r-1)\|f\|_{\beta}^{(\alpha/2)}}{3} h^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{-\alpha/2-2/r}, & N < i \leq 2N-1 \end{cases}$$

2. See Proof 24, there is a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$ such that

$$(3.6) \quad \begin{aligned} &\frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \int_{x_{i-1}}^{x_i} f''(y) \frac{(y - x_{i-1})^3}{3!} dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} f''(y) \frac{(y - x_{i+1})^3}{3!} dy \right) \\ &\leq Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T - x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N-1 \end{cases} \end{aligned}$$

Summarizes, we get the result. \square

And define

$$(3.7) \quad R_i := D_h^2 I^{2-\alpha}(u - \Pi_h u)(x_i)$$

We have some results about the estimate of R_i

THEOREM 3.2. *For $1 \leq i < N/2$, there exists $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that*

$$(3.8) \quad R_i \leq \begin{cases} Ch^2 x_i^{-\alpha/2-2/r}, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2(x_i^{-1-\alpha} \ln(i) + \ln(N)), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r} x_i^{-1-\alpha}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

THEOREM 3.3. *For $N/2 \leq i \leq N$, there exists constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that*

$$(3.9) \quad R_i \leq C(r-1)h^2|T - x_{i-1}|^{1-\alpha} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

And for $N < i \leq 2N - 1$, it is symmetric to the previous case.

Combine Theorem 3.1, Theorem 3.2 and Theorem 3.3, the proof of Theorem 2.5 completed.

We prove Theorem 3.2 and Theorem 3.3 in next subsections below.

3.2. Mesh Transport Functions.

DEFINITION 3.4 (Mesh Transport Functions).

$$(3.10) \quad y_{i,j}(x) = \begin{cases} (x^{1/r} + Z_{j-i})^r & i < N, j < N \\ \frac{x^{1/r} - Z_i}{Z_1} h_N + x_N & i < N, j = N \\ 2T - (Z_{2N-(j-i)} - x^{1/r})^r & i < N, j > N \\ \left(\frac{Z_1}{h_N} (x - x_N) + Z_j \right)^r & i = N, j < N \\ x, & i = N, j = N \end{cases}$$

where

$$(3.11) \quad Z_j := T^{1/r} \frac{j}{N}$$

We give some properties of mesh transport functions.

LEMMA 3.5. y

3.3. Proof of Theorem 3.2.

$$(3.12) \quad D_h^2 I^{2-\alpha}(u - \Pi_h u)(x_i) = D_h^2 \left(\int_0^{2T} (u(y) - \Pi_h u(y)) \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \right)$$

For convience, let's denote

$$(3.13) \quad T_{ij} = \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy, \quad i = 0, \dots, 2N, j = 1, \dots, 2N$$

Also for simplicity, we denote

DEFINITION 3.6.

$$(3.14) \quad S_{ij} = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} T_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_{i+1}} T_{i+1,j} \right)$$

then

$$(3.15) \quad R_i = \sum_{j=1}^{2N} S_{ij}$$

131

132 LEMMA 3.7. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $1 \leq$*
 133 *$i < N/2$,*

$$(3.16) \quad \sum_{j=\max\{2i+1, i+3\}}^N S_{ij} \leq Ch^2 x_i^{-\alpha/2-2/r}$$

135 *Proof.* Let

$$136 \quad K_y(x) = \frac{|y-x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

137 For $\max\{2i+1, i+3\} \leq j \leq N$, by Lemma C.1 and Lemma C.2

$$(3.17) \quad \begin{aligned} S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) D_h^2 K_y(x_i) dy \\ &\leq Ch^2 \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} dy \\ &= Ch^2 \int_{x_{j-1}}^{x_j} y^{-\alpha/2-2/r-1} dy \end{aligned}$$

139 Therefore,

$$(3.18) \quad \begin{aligned} \sum_{j=\max\{2i+1, i+3\}}^N S_{ij} &\leq Ch^2 \int_{x_{2i}}^{x_N} y^{-\alpha/2-2/r-1} dy \\ &= \frac{C}{\alpha/2+2/r} h^2 (x_{2i}^{-\alpha/2-2/r} - T^{-\alpha/2-2/r}) \\ &\leq \frac{C}{\alpha/2+2/r} 2^{r(-\alpha/2-2/r)} h^2 x_i^{-\alpha/2-2/r} \end{aligned} \quad \square$$

141

142 LEMMA 3.8. *Thert exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $1 \leq$*
 143 *$i < N/2$,*

$$(3.19) \quad \sum_{j=N+1}^{2N} S_{ij} \leq \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

Proof. For $1 \leq i < N/2, N+1 \leq j \leq 2N-1$, by equation (C.2) and Lemma C.2

$$\begin{aligned} S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) D_h^2 K_y(x_i) dy \\ &\leq \int_{x_{j-1}}^{x_j} Ch^2 (2T - y)^{\alpha/2-2/r} y^{-1-\alpha} dy \\ &\leq Ch^2 T^{-1-\alpha} \int_{x_{j-1}}^{x_j} (2T - y)^{\alpha/2-2/r} dy \end{aligned}$$

$$\begin{aligned} \sum_{j=N+1}^{2N-1} S_{ij} &\leq CT^{-1-\alpha} h^2 \int_{x_N}^{x_{2N-1}} (2T - y)^{\alpha/2-2/r} dy \\ &\leq CT^{-1-\alpha} h^2 \begin{cases} \frac{1}{\alpha/2-2/r+1} T^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 > 0 \\ \ln(T) - \ln(h_{2N}), & \alpha/2 - 2/r + 1 = 0 \\ \frac{1}{|\alpha/2-2/r+1|} h_{2N}^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \\ &= \begin{cases} \frac{C}{\alpha/2-2/r+1} T^{-\alpha/2-2/r} h^2, & \alpha/2 - 2/r + 1 > 0 \\ CrT^{-1-\alpha} h^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ \frac{C}{|\alpha/2-2/r+1|} T^{-\alpha/2-2/r} h^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \end{aligned} \quad (3.20)$$

And by Lemma A.3

$$S_{i,2N} \leq CT^{-1-\alpha} h_{2N}^{\alpha/2+1} = CT^{-\alpha/2} h^{r\alpha/2+r}$$

And when $\alpha/2 - 2/r + 1 \geq 0$,

$$h^{r\alpha/2+r} \leq h^2$$

Summarizes, we get the result. □


For $i = 1, 2$.

LEMMA 3.9. *By Lemma C.5, Lemma 3.7 and Lemma 3.8 we get*

$$\begin{aligned} R_1 &= \sum_{j=1}^3 S_{1j} + \sum_{j=4}^{2N} S_{1j} \\ &\leq Ch^2 x_1^{-\alpha/2-2/r} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \end{aligned} \quad (3.21)$$

$$\begin{aligned} R_2 &= \sum_{j=1}^4 S_{2j} + \sum_{j=5}^{2N} S_{2j} \\ &\leq Ch^2 x_2^{-\alpha/2-2/r} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \end{aligned} \quad (3.22)$$

For $3 \leq i < N/2$, we have a new separation of R_i , Let's denote $k = \lceil \frac{i}{2} \rceil$.



160

$$\begin{aligned}
 R_i &= \sum_{j=1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &= \sum_{j=1}^{k-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right) \\
 &\quad + \sum_{j=k+1}^{2i-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\
 &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1,2i} + T_{i-1,2i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2i} \right) \\
 &\quad + \sum_{j=2i+1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &= I_1 + I_2 + I_3 + I_4 + I_5
 \end{aligned}
 \tag{3.23}$$

162

163 **LEMMA 3.10.** *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for*
 164 $3 \leq i \leq N, k = \lceil \frac{i}{2} \rceil$

$$|I_1| = \left| \sum_{j=1}^{k-1} S_{ij} \right| \leq \begin{cases} Ch^2 x_i^{-\alpha/2-2/r}, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 x_i^{-1-\alpha} \ln(i), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r} x_i^{-1-\alpha}, & \alpha/2 - 2/r + 1 < 0 \end{cases}
 \tag{3.24}$$

166 *Proof.* by Lemma A.3 , Lemma C.3

$$S_{i1} \leq C x_1^{\alpha/2} x_1 x_i^{-1-\alpha} = C x_1^{\alpha/2+1} x_i^{-1-\alpha} = C T^{\alpha/2+1} h^{r\alpha/2+r} x_i^{-1-\alpha}
 \tag{3.25}$$

167

For $2 \leq j \leq k-1$, by Lemma C.1 and Lemma C.3

$$\begin{aligned}
 S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) D_h^2 K_y(x_i) dy \\
 &\leq Ch^2 \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)} dy \\
 &= Ch^2 x_i^{-1-\alpha} \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} dy
 \end{aligned}
 \tag{3.26}$$

Therefore,

$$\begin{aligned}
 I_1 &= \sum_{j=1}^{k-1} S_{ij} = S_{i1} + \sum_{j=2}^{k-1} S_{ij} \\
 &\leq Ch^{r\alpha/2+r} x_i^{-1-\alpha} + Ch^2 x_i^{-1-\alpha} \int_{x_1}^{x_{\lceil \frac{j}{2} \rceil-1}} y^{\alpha/2-2/r} dy \\
 &\leq Ch^{r\alpha/2+r} x_i^{-1-\alpha} + Ch^2 x_i^{-1-\alpha} \int_{x_1}^{2^{-r} x_i} y^{\alpha/2-2/r} dy
 \end{aligned}
 \tag{3.27}$$

But

$$\int_{x_1}^{2^{-r} x_i} y^{\alpha/2-2/r} dy \leq \begin{cases} \frac{1}{\alpha/2-2/r+1} (2^{-r} x_i)^{\alpha/2-2/r+1}, & \alpha/2-2/r+1 > 0 \\ \ln(2^{-r} x_i) - \ln(x_1), & \alpha/2-2/r+1 = 0 \\ \frac{1}{|\alpha/2-2/r+1|} x_1^{\alpha/2-2/r+1}, & \alpha/2-2/r+1 < 0 \end{cases}
 \tag{3.28}$$

So we have

$$I_1 \leq \begin{cases} \frac{C}{\alpha/2-2/r+1} h^2 x_i^{-\alpha/2-2/r}, & \alpha/2-2/r+1 > 0 \\ Ch^2 x_i^{-1-\alpha} \ln(i), & \alpha/2-2/r+1 = 0 \\ \frac{C}{|\alpha/2-2/r+1|} h^{r\alpha/2+r} x_i^{-1-\alpha}, & \alpha/2-2/r+1 < 0 \end{cases} \quad \square
 \tag{3.29}$$

DEFINITION 3.11. For convience, let's denote

$$V_{ij} = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right)
 \tag{3.30}$$

THEOREM 3.12. There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $3 \leq i < N/2, k = \lceil \frac{i}{2} \rceil$,

$$I_3 = \sum_{j=k+1}^{2i-1} V_{ij} \leq Ch^2 x_i^{-\alpha/2-2/r}
 \tag{3.31}$$

To estimate V_{ij} , we need some preparations.

LEMMA 3.13. For $y \in (x_{j-1}, x_j)$, we can rewrite

$$y = x_{j-1} + \theta h_j = (1-\theta)x_{j-1} + \theta x_j =: y_j^\theta, \quad \theta \in (0, 1)
 \tag{3.32}$$

185 by Lemma A.2,

$$\begin{aligned}
 T_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\
 &= \int_0^1 (u(y_j^\theta) - \Pi_h u(y_j^\theta)) \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} h_j d\theta \\
 &= \int_0^1 -\frac{\theta(1-\theta)}{2} h_j^3 u''(y_j^\theta) \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} \\
 &\quad + \frac{\theta(1-\theta)}{3!} h_j^4 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} (\theta^2 u'''(\eta_{j,1}^\theta) - (1-\theta)^2 u'''(\eta_{j,2}^\theta)) d\theta
 \end{aligned}
 \tag{3.33}$$

187 where $\eta_{j,1}^\theta \in (x_{j-1}, y_j^\theta)$, $\eta_{j,2}^\theta \in (y_j^\theta, x_j)$.

188 Now Let's construct a series of functions to represent T_{ij} .

DEFINITION 3.14.

$$189 \quad (3.34) \quad y_{j-i}(x) = (x^{1/r} + Z_{j-i})^r, \quad Z_{j-i} = T^{1/r} \frac{j-i}{N}$$

190 Particularly, for $i, j \leq N-1$,

$$191 \quad y_{j-i}(x_{i-1}) = x_{j-1}, \quad y_{j-i}(x_i) = x_j, \quad y_{j-i}(x_{i+1}) = x_{j+1}$$

$$193 \quad (3.35) \quad y_{j-i}'(x) = y_{j-i}(x)^{1-1/r} x^{1/r-1}$$

$$194 \quad (3.36) \quad y_{j-i}''(x) = \frac{1-r}{r} y_{j-i}(x)^{1-2/r} x^{1/r-2} Z_{j-i}$$

$$195 \quad (3.37)$$

$$197 \quad (3.38) \quad y_{j-i}^\theta(x) = (1-\theta)y_{j-1-i}(x) + \theta y_{j-i}(x)$$

198

$$199 \quad (3.39) \quad h_{j-i}(x) = y_{j-i}(x) - y_{j-i-1}(x)$$

200 Now, we define

$$201 \quad (3.40) \quad P_{j-i}^\theta(x) = (h_{j-i}(x))^3 u''(y_{j-i}^\theta(x)) \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

202

$$203 \quad (3.41) \quad Q_{j-i}^\theta(x) = (h_{j-i}(x))^4 \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

204 And now we can rewrite T_{ij}

205 LEMMA 3.15. For $2 \leq i \leq N$, $2 \leq j \leq N$,

$$\begin{aligned}
 T_{ij} &= \int_0^1 -\frac{\theta(1-\theta)}{2} P_{j-i}^\theta(x_i) d\theta \\
 &\quad + \int_0^1 \frac{\theta(1-\theta)}{3!} Q_{j-i}^\theta(x_i) [\theta^2 u'''(\eta_{j,1}^\theta) - (1-\theta)^2 u'''(\eta_{j,2}^\theta)] d\theta
 \end{aligned}
 \tag{3.42}$$

Immediately, we can see from (3.30) that

LEMMA 3.16. For $3 \leq i, j \leq N-1$,

$$\begin{aligned}
 (3.43) \quad V_{ij} &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\
 &= \int_0^1 -\frac{\theta(1-\theta)}{2} D_h^2 P_{j-i}^\theta(x_i) d\theta \\
 &\quad + \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1}) u'''(\eta_{j+1,1}^\theta) - Q_{j-i}^\theta(x_i) u'''(\eta_{j,1}^\theta)}{h_{i+1}} \right) d\theta \\
 &\quad - \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i) u'''(\eta_{j,1}^\theta) - Q_{j-i}^\theta(x_{i-1}) u'''(\eta_{j-1,1}^\theta)}{h_i} \right) d\theta \\
 &\quad - \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1}) u'''(\eta_{j+1,2}^\theta) - Q_{j-i}^\theta(x_i) u'''(\eta_{j,2}^\theta)}{h_{i+1}} \right) d\theta \\
 &\quad + \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i) u'''(\eta_{j,2}^\theta) - Q_{j-i}^\theta(x_{i-1}) u'''(\eta_{j-1,2}^\theta)}{h_i} \right) d\theta
 \end{aligned}$$

To estimate V_{ij} , we first estimate $D_h^2 P_{j-i}^\theta(x_i)$, but By Lemma A.1,

$$(3.44) \quad D_h^2 P_{j-i}^\theta(x_i) = P_{j-i}^{\theta''}(\xi), \quad \xi \in (x_{i-1}, x_{i+1})$$

By Leibniz formula, we calculate and estimate the derivations of $h_{j-i}^3(x)$, $u''(y_{j-i}^\theta(x))$ and $\frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$ separately.

Firstly, we have

LEMMA 3.17. There exists a constant $C = C(T, r)$ such that For $3 \leq i \leq N-1$, $\lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\}$, $\xi \in (x_{i-1}, x_{i+1})$,

$$(3.45) \quad h_{j-i}^3(\xi) \leq C h^2 x_i^{2-2/r} h_j$$

$$(3.46) \quad (h_{j-i}^3(\xi))' \leq C(r-1) h^2 x_i^{1-2/r} h_j$$

$$(3.47) \quad (h_{j-i}^3(\xi))'' \leq C(r-1) h^2 x_i^{-2/r} h_j$$

The proof of this theorem see Lemma C.6 and Lemma C.7

Second,

LEMMA 3.18. There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $3 \leq i \leq N-1$, $\lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\}$, $\xi \in (x_{i-1}, x_{i+1})$,

$$(3.48) \quad u''(y_{j-i}^\theta(\xi)) \leq C x_i^{\alpha/2-2}$$

$$(3.49) \quad (u''(y_{j-i}^\theta(\xi)))' \leq C x_i^{\alpha/2-3}$$

$$(3.50) \quad (u''(y_{j-i}^\theta(\xi)))'' \leq C x_i^{\alpha/2-4}$$

The proof of this theorem see Proof 30

And Finally, we have

LEMMA 3.19. *There exists a constant $C = C(T, \alpha, r)$ such that For $3 \leq i \leq N - 1, \lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\}, \xi \in (x_{i-1}, x_{i+1}),$*

$$(3.51) \quad |y_{j-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_j^\theta - x_i|^{1-\alpha}$$

$$(3.52) \quad |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'| \leq C|y_j^\theta - x_i|^{1-\alpha}x_i^{-1}$$

$$(3.53) \quad |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})''| \leq C|y_j^\theta - x_i|^{1-\alpha}x_i^{-2}$$

where $y_j^\theta = \theta x_{j-1} + (1 - \theta)x_j$

The proof of this theorem see Proof 31

LEMMA 3.20. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $3 \leq i \leq N - 1, \lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i - 1, N - 1\},$*

$$(3.54) \quad D_h^2 P_{j-i}^\theta(x_i) \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j$$

where $y_j^\theta = \theta x_{j-1} + (1 - \theta)x_j$

Proof. Since Lemma A.1

$$(3.55) \quad D_h^2 P_{j-i}^\theta(x_i) = P_{j-i}^{\theta''}(\xi), \quad \xi \in (x_{i-1}, x_{i+1})$$

From (3.40), using Leibniz formula and Lemma 3.17, Lemma 3.18 and Lemma 3.19□

LEMMA 3.21. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $3 \leq i \leq N - 1.$
For $\lceil \frac{i}{2} \rceil \leq j \leq \min\{2i - 1, N - 1\},$*

$$(3.56) \quad \begin{aligned} & \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \right) \\ & \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j \end{aligned}$$

And for $\lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i, N\},$

$$(3.57) \quad \begin{aligned} & \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta) - Q_{j-i}^\theta(x_{i-1})u'''(\eta_{j-1}^\theta)}{h_i} \right) \\ & \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j \end{aligned}$$

where $\eta_j^\theta \in (x_{j-1}, x_j).$

proof see Proof 32

LEMMA 3.22. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $3 \leq i \leq N - 1, \lceil \frac{i}{2} \rceil + 1 \leq j \leq \min\{2i - 1, N - 1\},$*

$$(3.58) \quad \begin{aligned} V_{ij} & \leq Ch^2 \int_0^1 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j d\theta \\ & = Ch^2 \int_{x_{j-1}}^{x_j} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} dy \end{aligned}$$

257 *Proof.* Since Lemma 3.16, by Lemma 3.20 and Lemma 3.21, we get the result
 258 immediately. \square

259 Now we can prove Theorem 3.12 using Lemma 3.22, $k = \lceil \frac{i}{2} \rceil$

$$\begin{aligned}
 I_3 &= \sum_{k+1}^{2i-1} V_{ij} \leq Ch^2 \int_{x_k}^{x_{2i-1}} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} dy \\
 (3.59) \quad &= Ch^2 \left(\frac{|x_k - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} + \frac{|x_{2i-1} - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} \right) x_i^{\alpha/2-2-2/r} \\
 &\leq Ch^2 x_i^{2-\alpha} x_i^{\alpha/2-2-2/r} = Ch^2 x_i^{-\alpha/2-2/r}
 \end{aligned}$$

261 Now we study I_2, I_4 .

262 LEMMA 3.23. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for*
 263 $3 \leq i \leq N-1, k = \lceil \frac{i}{2} \rceil$,
 (3.60)

$$I_2 = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right) \leq Ch^2 x_i^{-\alpha/2-2/r}$$

265 And for $3 \leq i < N/2$,
 (3.61)

$$I_4 = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1,2i} + T_{i-1,2i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2i} \right) \leq Ch^2 x_i^{-\alpha/2-2/r}$$

267 *Proof.* In fact,

$$\begin{aligned}
 (3.62) \quad &\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \\
 &= \frac{1}{h_{i+1}} (T_{i+1,k} - T_{i,k}) + \frac{1}{h_{i+1}} (T_{i+1,k+1} - T_{i,k}) + \left(\frac{1}{h_{i+1}} - \frac{1}{h_i} \right) T_{i,k}
 \end{aligned}$$

269 While, by Lemma A.2 and Lemma B.1
 (3.63)

$$\begin{aligned}
 \frac{1}{h_{i+1}} (T_{i+1,k} - T_{i,k}) &= \int_{x_{k-1}}^{x_k} (u(y) - \Pi_h u(y)) \frac{|x_{i+1} - y|^{1-\alpha} - |x_i - y|^{1-\alpha}}{h_{i+1} \Gamma(2-\alpha)} dy \\
 (3.63) \quad &\leq h_k^2 \max_{\eta \in (x_{k-1}, x_k)} |u''(\eta)| \int_{x_{k-1}}^{x_k} \frac{|\xi - y|^{-\alpha}}{\Gamma(1-\alpha)} dy, \quad \xi \in (x_i, x_{i+1}) \\
 &\leq Ch^2 x_k^{2-2/r} x_{k-1}^{\alpha/2-2} h_k |x_i - x_k|^{-\alpha} \\
 &\leq Ch^2 x_i^{-\alpha/2-2/r} h_k
 \end{aligned}$$

271 Thus,

$$(3.64) \quad \frac{2}{h_i + h_{i+1}} \frac{1}{h_{i+1}} |T_{i+1,k} - T_{i,k}| \leq Ch^2 x_i^{-\alpha/2-2/r}$$

From Lemma 3.15
(3.65)

$$\begin{aligned} \frac{1}{h_{i+1}}(T_{i+1,k+1} - T_{i,k}) &= \int_0^1 -\frac{\theta(1-\theta)}{2} \frac{P_{k-i}^\theta(x_{i+1}) - P_{k-i}^\theta(x_i)}{h_{i+1}} d\theta \\ &+ \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{Q_{k-i}^\theta(x_{i+1})u'''(\eta_{k+1,1}^\theta) - Q_{k-i}^\theta(x_i)u'''(\eta_{k,1}^\theta)}{h_{i+1}} d\theta \\ &- \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{Q_{k-i}^\theta(x_{i+1})u'''(\eta_{k+1,2}^\theta) - Q_{k-i}^\theta(x_i)u'''(\eta_{k,2}^\theta)}{h_{i+1}} d\theta \end{aligned}$$

and

$$(3.66) \quad D_h P_{k-i}^\theta(x_i) := \frac{P_{k-i}^\theta(x_{i+1}) - P_{k-i}^\theta(x_i)}{h_{i+1}} = P_{k-i}^{\theta'}(\xi), \quad \xi \in (x_i, x_{i+1})$$

Similar with Lemma 3.20, from Lemma 3.15, using Leibniz formula, by Lemma C.6,
Lemma 3.18 and Lemma 3.19 we get

$$(3.67) \quad |D_h P_{k-i}^\theta(x_i)| \leq Ch^2 x_i^{-\alpha/2-2/r} h_k$$

And with Lemma 3.21, we can get

$$(3.68) \quad \frac{2}{h_i + h_{i+1}} \frac{1}{h_{i+1}} |T_{i+1,k+1} - T_{i,k}| \leq Ch^2 x_i^{-\alpha/2-2/r}$$

For the third term, by Lemma B.1, Lemma B.2 and Lemma A.2

$$\begin{aligned} (3.69) \quad \frac{2}{h_i + h_{i+1}} \frac{h_{i+1} - h_i}{h_i h_{i+1}} T_{i,k} &\leq h_i^{-3} h^2 x_i^{1-2/r} h_k Ch_k^2 x_{k-1}^{\alpha/2-2} |x_k - x_i|^{1-\alpha} \\ &\leq Ch^2 x_i^{-\alpha/2-2/r} \end{aligned}$$

Summarizes, we have

$$(3.70) \quad I_2 \leq Ch^2 x_i^{-\alpha/2-2/r}$$

The case for I_4 is similar. \square

Now combine Lemma 3.9, Lemma 3.10, Lemma 3.23, Theorem 3.12, Lemma 3.7
and Lemma 3.8, we get Theorem 3.2.

3.4. Proof of Theorem 3.3. For $N/2 \leq i < N, k = \lceil \frac{i}{2} \rceil$, we have

$$\begin{aligned}
 (3.71) \quad R_i &= \sum_{j=1}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &= \sum_{j=1}^{k-1} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} (T_{i+1,k} + T_{i+1,k+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,k} \right) \\
 &\quad + \sum_{j=k+1}^{N-1} + \sum_{j=N}^{N+1} + \sum_{j=N+2}^{2N-\lceil \frac{N}{2} \rceil} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j-1} \right) \\
 &\quad + \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1,2N-\lceil \frac{N}{2} \rceil+1} + T_{i-1,2N-\lceil \frac{N}{2} \rceil}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,2N-\lceil \frac{N}{2} \rceil+1} \right) \\
 &\quad + \sum_{j=2N-\lceil \frac{N}{2} \rceil+2}^{2N} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_i} T_{i-1,j} \right) \\
 &= I_1 + I_2 + I_3^1 + I_3^2 + I_3^3 + I_4 + I_5
 \end{aligned}$$

We have estimate I_1 in Lemma 3.10 and I_2 in Lemma 3.23. We can control I_3^1 similar with Theorem 3.12 by Lemma 3.22 where $2i - 1 \geq N - 1$

LEMMA 3.24. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $N/2 \leq i < N, k = \lceil \frac{i}{2} \rceil$,*

$$\begin{aligned}
 (3.72) \quad I_3^1 &= \sum_{j=k+1}^{N-1} V_{ij} \leq Ch^2 \int_{x_k}^{x_{N-1}} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} dy \\
 &= Ch^2 \left(\frac{|x_k - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} + \frac{|x_{N-1} - x_i|^{2-\alpha}}{\Gamma(3-\alpha)} \right) x_i^{\alpha/2-2-2/r} \\
 &\leq Ch^2 x_i^{2-\alpha} x_i^{\alpha/2-2-2/r} = Ch^2 x_i^{-\alpha/2-2/r}
 \end{aligned}$$

Let's study I_3^3 before I_3^2 .

$$(3.73) \quad I_3^3 = \sum_{j=N+2}^{2N-\lceil \frac{N}{2} \rceil} V_{ij}$$

Similarly, Let's define a new series of functions

DEFINITION 3.25. *For $i \leq N - 1, j \geq N + 1$, with no confusion, we also denote in this section*

$$(3.74) \quad y_{j-i}(x) = 2T - (Z_{2N-j+i} - x^{1/r})^r, \quad Z_{2N-j+i} = T^{1/r} \frac{2N-j+i}{N}$$

Particularly

$$y_{j-i}(x_{i-1}) = x_{j-1}, \quad y_{j-i}(x_i) = x_j, \quad y_{j-i}(x_{i+1}) = x_{j+1}$$

304 $y \rightarrow z?$

305 (3.75) $y_{j-i}'(x) = (2T - y_{j-i}(x))^{1-1/r} x^{1/r-1}$

306 (3.76) $y_{j-i}''(x) = \frac{1-r}{r} (2T - y_{j-i}(x))^{1-2/r} x^{1/r-2} Z_{2N-j+i}$

307 (3.77)

308

309 (3.78) $y_{j-i}^\theta(x) = (1 - \theta)y_{j-i-1}(x) + \theta y_{j-i}(x)$

310

311 (3.79) $h_{j-i}(x) = y_{j-i}(x) - y_{j-i-1}(x)$

312

313 (3.80) $P_{j-i}^\theta(x) = (h_{j-i}(x))^3 u''(y_{j-i}^\theta(x)) \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$

314

315 (3.81) $Q_{j-i}^\theta(x) = (h_{j-i}(x))^4 \frac{|y_{j-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$

316 Now we have the same formula Lemma 3.16 for $i \leq N-1, j \geq N+2$,

317 Similarly, we first estimate

318 (3.82) $D_h^2 P_{j-i}^\theta(\xi) = P_{j-i}^{\theta}{}''(\xi), \quad \xi \in (x_{i-1}, x_{i+1})$

319 Combine Definition 3.25, Lemma C.8, Lemma C.9 and Lemma C.10, using Leibniz
320 formula, we have

321 LEMMA 3.26. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For*
322 *$N/2 \leq i \leq N-1, N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1$, we have*

323 (3.83)
$$\begin{aligned} |D_h^2 P_{j-i}^\theta(\xi)| &\leq Ch_j h^2 \left(|y_j^\theta - x_i|^{1-\alpha} \right. \\ &\quad + |y_j^\theta - x_i|^{-\alpha} (|2T - x_i - y_j^\theta| + h_N) \\ &\quad + |y_j^\theta - x_i|^{-1-\alpha} (|2T - x_i - y_j^\theta| + h_N)^2 \\ &\quad \left. + (r-1) |y_j^\theta - x_i|^{-\alpha} \right) \end{aligned}$$

324 And

325 LEMMA 3.27. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For*
326 *$N/2 \leq i \leq N-1, N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil, \xi \in (x_{i-1}, x_{i+1})$, we have*

327 (3.84)
$$\begin{aligned} \frac{2}{h_i + h_{i+1}} &\left| \frac{Q_{j-i}^\theta(x_{i+1}) u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i) u'''(\eta_j^\theta)}{h_{i+1}} \right| \\ &\leq Ch^2 h_j (|y_j^\theta - x_i|^{1-\alpha} + |y_j^\theta - x_i|^{-\alpha} (|2T - x_i - y_j^\theta| + h_N)) \end{aligned}$$

328 and

329 (3.85)
$$\begin{aligned} \frac{2}{h_i + h_{i+1}} &\left(\frac{Q_{j-i}^\theta(x_i) u'''(\eta_j^\theta) - Q_{j-i}^\theta(x_{i-1}) u'''(\eta_{j-1}^\theta)}{h_{i+1}} \right) \\ &\leq Ch^2 h_j (|y_j^\theta - x_i|^{1-\alpha} + |y_j^\theta - x_i|^{-\alpha} (|2T - x_i - y_j^\theta| + h_N)) \end{aligned}$$

Proof. From Definition 3.25, by Lemma C.8 and Lemma C.10, for $\xi \in (x_i, x_{i+1})$, by Leibniz formula, we have

$$(3.86) \quad |Q_{j-i}^\theta(\xi)| \leq Ch^2 h_j^2 ((r-1)|y_j^\theta - x_i|^{1-\alpha} + |y_j^\theta - x_i|^{-\alpha}(|2T - x_i - y_j^\theta| + h_N))$$

$$(3.87) \quad |Q_{j-i}^\theta(\xi)| \leq Ch^2 h_j^2 |y_j^\theta - x_i|^{1-\alpha}$$

So use the skill in Proof 32 with Lemma C.9

$$(3.88) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \right) \leq Ch^2 h_j (|y_j^\theta - x_i|^{1-\alpha} + |y_j^\theta - x_i|^{-\alpha}(|2T - x_i - y_j^\theta| + h_N)) \quad \square$$

Combine Lemma 3.26, Lemma 3.27 and formula Lemma 3.16 for $i \leq N-1, j \geq N+2$, we have

LEMMA 3.28. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq i \leq N-1, N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1$*

$$(3.89) \quad V_{ij} \leq Ch^2 \int_{x_{j-1}}^{x_j} \left(|y - x_i|^{1-\alpha} + |y - x_i|^{-\alpha}(|2T - x_i - y| + h_N) + |y - x_i|^{-1-\alpha}(|2T - x_i - y| + h_N)^2 + (r-1)|y - x_i|^{-\alpha} \right) dy$$

We can estimate I_3^3 Now.

LEMMA 3.29. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq i \leq N-1$, we have*

$$(3.90) \quad I_3^3 = \sum_{j=N+2}^{2N - \lceil \frac{N}{2} \rceil} V_{ij} \leq Ch^2 + C(r-1)h^2 |T - x_{i-1}|^{1-\alpha}$$

Proof.

$$(3.91) \quad I_3^3 = \sum_{j=N+2}^{2N - \lceil \frac{N}{2} \rceil} V_{ij} \leq Ch^2 \int_{x_{N+1}}^{x_{2N - \lceil \frac{N}{2} \rceil}} \left(|y - x_i|^{1-\alpha} + |y - x_i|^{-\alpha}(|2T - x_i - y| + h_N) + |y - x_i|^{-1-\alpha}(|2T - x_i - y| + h_N)^2 + (r-1)|y - x_i|^{-\alpha} \right) dy$$

Since

$$(3.92) \quad |2T - x_i - y| + h_N \leq y - x_i$$

$$(3.93) \quad \begin{aligned} I_3^3 &\leq Ch^2 \int_{x_{N+1}}^{x_{2N - \lceil \frac{N}{2} \rceil}} |y - x_i|^{1-\alpha} + (r-1)|y - x_i|^{-\alpha} \\ &\leq Ch^2 (T^{2-\alpha} + (r-1)|x_{N+1} - x_i|^{1-\alpha}) \\ &\leq Ch^2 + C(r-1)h^2 |T - x_{i-1}|^{1-\alpha} \end{aligned} \quad \square$$

For I_3^2 , we have

THEOREM 3.30. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that, for $N/2 \leq i \leq N-1$*

$$(3.94) \quad \begin{aligned} V_{iN} &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} T_{i+1, N+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i, N} + \frac{1}{h_i} T_{i-1, N-1} \right) \\ &\leq Ch^2 + C(r-1)h^2 |T - x_{i-1}|^{1-\alpha} \end{aligned}$$

Proof. We use the similar skill in the last section, but more complicated. for $j = N$, Let

$$(3.95) \quad {}_L y_{N-1-i}(x) = (x^{1/r} + Z_{N-1-i})^r, \quad Z_{N-1-i} = T^{1/r} \frac{N-1-i}{N}$$

$$(3.96) \quad {}_0 y_{N-i}(x) = \frac{x^{1/r} - Z_i}{Z_1} h_N + T, \quad Z_i = T^{1/r} \frac{i}{N}, x_N = T$$

and

$$(3.97) \quad {}_R y_{N+1-i}(x) = 2T - (Z_{N-1+i} - x^{1/r})^r, \quad Z_{N-1+i} = T^{1/r} \frac{N-1+i}{N}$$

Thus,

$${}_L y_{N-1-i}(x_{i-1}) = x_{N-2}, \quad {}_L y_{N-1-i}(x_i) = x_{N-1}, \quad {}_L y_{N-1-i}(x_{i+1}) = x_N$$

$${}_0 y_{N-i}(x_{i-1}) = x_{N-1}, \quad {}_0 y_{N-i}(x_i) = x_N, \quad {}_0 y_{N-i}(x_{i+1}) = x_{N+1}$$

$${}_R y_{N+1-i}(x_{i-1}) = x_N, \quad {}_R y_{N+1-i}(x_i) = x_{N+1}, \quad {}_R y_{N+1-i}(x_{i+1}) = x_{N+2}$$

Then, define

$$(3.98) \quad {}_L y_{N-i}^\theta(x) = \theta {}_L y_{N-1-i}(x) + (1-\theta) {}_0 y_{N-i}(x)$$

$$(3.99) \quad {}_R y_{N+1-i}^\theta(x) = \theta {}_0 y_{N-i}(x) + (1-\theta) {}_R y_{N+1-i}(x)$$

$$(3.100) \quad {}_L h_{N-i}(x) = {}_0 y_{N-i}(x) - {}_L y_{N-1-i}(x)$$

$$(3.101) \quad {}_R h_{N+1-i}(x) = {}_R y_{N+1-i}(x) - {}_0 y_{N-i}(x)$$

We have

$$(3.102) \quad {}_L y_{N-1-i}'(x) = {}_L y_{N-1-i}^{1-1/r}(x) x^{1/r-1}$$

$$(3.103) \quad {}_L y_{N-1-i}''(x) = \frac{1-r}{r} {}_L y_{N-1-i}^{1-2/r}(x) x^{1/r-2} Z_{N-1-i}$$

$$(3.104) \quad {}_0 y_{N-i}'(x) = \frac{1}{r} \frac{h_N}{Z_1} x^{1/r-1}$$

$$(3.105) \quad {}_0 y_{N-i}''(x) = \frac{1-r}{r^2} \frac{h_N}{Z_1} x^{1/r-2}$$

$$(3.106) \quad {}_R y_{N+1-i}'(x) = (2T - {}_R y_{N+1-i}(x))^{1-1/r} x^{1/r-1}$$

$$(3.107) \quad {}_R y_{N+1-i}''(x) = \frac{1-r}{r} (2T - {}_R y_{N+1-i}(x))^{1-2/r} x^{1/r-2} Z_{N-1+i}$$

379

$$380 \quad (3.108) \quad {}_L P_{N-i}^\theta(x) = ({}_L h_{N-i}(x))^3 \frac{|{}_L y_{N-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)} u''({}_L y_{N-i}^\theta(x))$$

$$381 \quad (3.109) \quad {}_R P_{N+1-i}^\theta(x) = ({}_R h_{N+1-i}(x))^3 \frac{|{}_R y_{N+1-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)} u''({}_R y_{N+1-i}^\theta(x))$$

$$382 \quad (3.110) \quad {}_L Q_{N-i}^\theta(x) = ({}_L h_{N-i}(x))^4 \frac{|{}_L y_{N-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

$$383 \quad (3.111) \quad {}_R Q_{N+1-i}^\theta(x) = ({}_R h_{N+1-i}(x))^4 \frac{|{}_R y_{N+1-i}^\theta(x) - x|^{1-\alpha}}{\Gamma(2-\alpha)}$$

 384 Similar with Lemma 3.15, we can get for $l = -1, 0, 1$,

$$385 \quad (3.112) \quad \begin{aligned} T_{i+l, N+l} &= \int_0^1 -\frac{\theta(1-\theta)}{2} {}_L P_{N-i}^\theta(x_{i+l}) d\theta \\ &+ \int_0^1 \frac{\theta(1-\theta)}{3!} {}_L Q_{N-i}^\theta(x_{i+l}) (\theta^2 u'''(\eta_{N+l,1}^\theta) - (1-\theta)^2 u'''(\eta_{N+l,2}^\theta)) d\theta \end{aligned}$$

386

$$(3.113) \quad \begin{aligned} T_{i+l, N+1+l} &= \int_0^1 -\frac{\theta(1-\theta)}{2} {}_R P_{N+1-i}^\theta(x_{i+l}) d\theta \\ &+ \int_0^1 \frac{\theta(1-\theta)}{3!} {}_R Q_{N+1-i}^\theta(x_{i+l}) (\theta^2 u'''(\eta_{N+1+l,1}^\theta) - (1-\theta)^2 u'''(\eta_{N+1+l,2}^\theta)) d\theta \end{aligned}$$

387

388 So we have

$$(3.114) \quad \begin{aligned} V_{i,N} &= \int_0^1 -\frac{\theta(1-\theta)}{2} D_{hL}^2 {}_L P_{N-i}^\theta(x_i) d\theta \\ &+ \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_{i+1}) u'''(\eta_{N+1,1}^\theta) - {}_L Q_{N-i}^\theta(x_i) u'''(\eta_{N,1}^\theta)}{h_{i+1}} \right) d\theta \\ &- \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_i) u'''(\eta_{N,1}^\theta) - {}_L Q_{N-i}^\theta(x_{i-1}) u'''(\eta_{N-1,1}^\theta)}{h_i} \right) d\theta \\ &- \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_{i+1}) u'''(\eta_{N+1,2}^\theta) - {}_L Q_{N-i}^\theta(x_i) u'''(\eta_{N,2}^\theta)}{h_{i+1}} \right) d\theta \\ &+ \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_i) u'''(\eta_{N,2}^\theta) - {}_L Q_{N-i}^\theta(x_{i-1}) u'''(\eta_{N-1,2}^\theta)}{h_i} \right) d\theta \end{aligned}$$

389

 390 $N+1$ is similar.

 391 We estimate $D_{hL}^2 {}_L P_{N-i}^\theta(x_i) = {}_L P_{N-i}^{\theta''}(\xi), \xi \in (x_{i-1}, x_{i+1})$,

392

LEMMA 3.31.

$$393 \quad (3.115) \quad {}_L h_{N-i}^3(\xi) \leq Ch_N^3 \leq Ch^3$$

$$394 \quad (3.116) \quad {}_R h_{N+1-i}^3(\xi) \leq Ch_N^3 \leq Ch^3$$

$$(3.117) \quad ({}_L h_{N-i}^3(\xi))' \leq C(r-1)h_N^2 h \leq C(r-1)h^3$$

$$(3.118) \quad ({}_R h_{N+1-i}^3(\xi))' \leq C(r-1)h_N^2 h \leq C(r-1)h^3$$

$$(3.119) \quad ({}_L h_{N-i}^3(\xi))'' \leq C(r-1)h^2$$

$$(3.120) \quad ({}_R h_{N+1-i}^3(\xi))'' \leq C(r-1)h^2$$

Proof.

$$(3.121) \quad {}_L h_{N-i}(\xi) \leq 2(C?)h_N, \quad {}_R h_{N+1-i}(\xi) \leq 2h_N$$

400

$$(3.122) \quad \begin{aligned} ({}_L h_{N-i}^l(\xi))' &= {}_L h_{N-i}^{l-1}(\xi)({}_0 y_{N-i}'(\xi) - {}_L y_{N-1-i}'(\xi)) \\ &= {}_L h_{N-i}^{l-1}(\xi)\xi^{1/r-1}\left(\frac{1}{r}\frac{h_N}{Z_1} - {}_L y_{N-1-i}^{1-1/r}(\xi)\right) \end{aligned}$$

402 while

(3.123)

$$\begin{aligned} \left|\frac{1}{r}\frac{h_N}{Z_1} - {}_L y_{N-1-i}^{1-1/r}(\xi)\right| &= \left|\frac{1}{r}\frac{x_N - (x_N^{1/r} - Z_1)^r}{Z_1} - \eta^{1-1/r}\right| \quad \eta \in [x_{N-2}, x_N] \\ &= T^{1-1/r} \left| \left(\frac{N-t}{N}\right)^{r-1} - \left(\frac{N-s}{N}\right)^{r-1} \right| \quad t \in [0, 1], s \in [0, 2] \\ &\leq T^{1-1/r} \left| 1 - \left(\frac{N-2}{N}\right)^{r-1} \right| \leq CT^{1-1/r}(r-1)\frac{2}{N} \end{aligned}$$

404 Thus,

$$(3.124) \quad ({}_L h_{N-i}^l(\xi))' \leq C(r-1)h_N^{l-1}x_i^{1/r-1}h$$

406 And

(3.125)

$$\begin{aligned} ({}_L h_{N-i}^3(\xi))'' &= 3{}_L h_{N-i}^2(\xi){}_L h_{N-i}''(\xi) + 6{}_L h_{N-i}(\xi)({}_L h_{N-i}'(\xi))^2 \\ &\leq Ch_N^2 \frac{1-r}{r} x_i^{1/r-2} \left(\frac{1}{r}\frac{h_N}{Z_1} - {}_L y_{N-1-i}^{1-2/r}(\xi)Z_{N-1-i}\right) + Ch_N(r-1)^2 h^2 x_i^{2/r-2} \end{aligned}$$

$$(3.126) \quad \left|\frac{h_N}{rZ_1} - {}_L y_{N-1-i}^{1-2/r}(\xi)Z_{N-1-i}\right| \leq T^{1-1/r} + Cx_N^{1-2/r}x_N^{1/r} = CT^{1-1/r}$$

409 So

$$\begin{aligned} (3.126) \quad ({}_L h_{N-i}^3(\xi))'' &\leq Ch_N^2 \frac{1-r}{r} x_i^{1/r-2} + C(r-1)^2 h_N x_i^{2/r-2} h^2 \\ &\leq C(r-1)h_N^2 \end{aligned}$$

411 ${}_R h_{N+1-i}^3(\xi)$ is similar. □

LEMMA 3.32.

$$(3.127) \quad u''({}_L y_{N-i}^\theta(\xi)) \leq Cx_{N-2}^{-\alpha/2-2} \leq C$$

$$(3.128) \quad (u''({}_L y_{N-i}^\theta(\xi)))' \leq C$$

$$(3.129) \quad (u''({}_L y_{N-i}^\theta(\xi)))'' \leq C$$

Proof.

$$\begin{aligned}
 (u''(Ly_{N-i}^\theta(\xi)))' &= u'''(Ly_{N-i}^\theta(\xi))Ly_{N-i}^{\theta'}(\xi) \\
 &\leq C(\theta Ly_{N-1-i}'(\xi) + (1-\theta)_0 y_{N-i}'(\xi)) \\
 &\leq Cx_i^{1/r-1}(\theta Ly_{N-1-i}^{1-1/r}(\xi) + (1-\theta)\frac{h_N}{rZ_1}) \\
 &\leq Cx_i^{1/r-1}x_N^{1-1/r}
 \end{aligned}
 \tag{3.130}$$

And

$$\begin{aligned}
 (u''(Ly_{N-i}^\theta(\xi)))'' &= u''''(Ly_{N-i}^\theta(\xi))(Ly_{N-i}^{\theta'}(\xi))^2 + u'''(Ly_{N-i}^\theta(\xi))Ly_{N-i}^{\theta''}(\xi) \\
 &\leq Cx_i^{2/r-2}x_N^{2-2/r} + C\frac{r-1}{r}x_i^{1/r-2}(\theta x_N^{1-2/r}Z_{N-1-i} + (1-\theta)\frac{h_N}{rZ_1}) \\
 &\leq Cx_i^{2/r-2} + C(r-1)x_i^{1/r-2}T^{1-1/r}
 \end{aligned}
 \tag{3.131}$$

□

LEMMA 3.33.

$$|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_N^\theta - x_i|^{1-\alpha} \tag{3.132}$$

$$(|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha})' \leq C|y_N^\theta - x_i|^{1-\alpha} \tag{3.133}$$

$$(|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha})'' \leq C(r-1)|y_N^\theta - x_i|^{-\alpha} + |y_N^\theta - x_i|^{1-\alpha} \tag{3.134}$$

Proof.

$$\begin{aligned}
 (Ly_{N-i}^\theta(\xi) - \xi)' &= (\theta(Ly_{N-1-i}(\xi) - \xi) + (1-\theta)(_0y_{N-i}(\xi) - \xi))' \\
 &= \theta(Ly_{N-1-i}'(\xi) - 1) + (1-\theta)(_0y_{N-i}'(\xi) - 1) \\
 &= \theta\xi^{1/r-1}(Ly_{N-1-i}^{1-1/r}(\xi) - \xi^{1-1/r}) + (1-\theta)\xi^{1/r-1}(\frac{h_N}{rZ_1} - \xi^{1-1/r})
 \end{aligned}
 \tag{3.135}$$

$$\begin{aligned}
 (Ly_{N-i}^\theta(\xi) - \xi)'' &= \theta(Ly_{N-1-i}''(\xi)) + (1-\theta)(_0y_{N-i}''(\xi)) \\
 &= \frac{1-r}{r}\xi^{1/r-2}(\theta Ly_{N-1-i}^{1-2/r}(\xi)Z_{N-1-i} + (1-\theta)\frac{h_N}{rZ_1}) \leq 0
 \end{aligned}
 \tag{3.136}$$

And

$$|(Ly_{N-i}^\theta(\xi) - \xi)''| \leq C(r-1)\xi^{1/r-2}T^{1-1/r} \tag{3.137}$$

We have known

$$C|x_{N-1} - x_i| \leq |Ly_{N-1-i}(\xi) - \xi| \leq C|x_{N-1} - x_i| \tag{3.138}$$

If $\xi \leq x_{N-1}$, then $(_0y_{N-i}(\xi) - \xi)' \geq 0$, so

$$C|x_N - x_i| \leq |x_{N-1} - x_{i-1}| \leq |Ly_{N-i}^\theta(\xi) - \xi| \leq |x_{N+1} - x_{i+1}| \leq C|x_N - x_i| \tag{3.139}$$

If $i = N-1$ and $\xi \in [x_{N-1}, x_N]$, then $_0y_{N-i}(\xi) - \xi$ is concave, bigger than its two neighboring points, which are equal to h_N , so

$$h_N = |x_N - x_{N-1}| \leq |_0y_{N-i}(\xi) - \xi| \leq |x_{N+1} - x_{N-1}| = 2h_N \tag{3.140}$$

So we have

$$(3.141) \quad |Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_N^\theta - x_i|^{1-\alpha}$$

While

$$(3.142) \quad Ly_{N-1-i}^{1-1/r}(\xi) - \xi^{1-1/r} \leq (Ly_{N-1-i}(\xi) - \xi)\xi^{-1/r}$$

and

$$(3.143) \quad \left| \frac{h_N}{rZ_1} - \xi^{1-1/r} \right| \leq \max\left\{ \left| \frac{h_N}{rZ_1} - x_{i-1}^{1-1/r} \right|, \left| \frac{h_N}{rZ_1} - x_{i+1}^{1-1/r} \right| \right\}$$

$$\leq \max \begin{cases} T^{1-1/r} - x_{i-1}^{1-1/r} \leq |x_N - x_{i-1}|T^{-1/r} \leq C|x_N - x_i| \\ |x_{i+1}^{1-1/r} - x_{N-1}^{1-1/r}| \leq |x_{i+1} - x_{N-1}|x_{N-1}^{-1/r} \leq C|x_N - x_i| \end{cases}$$

So we have

$$(3.144) \quad (Ly_{N-i}^\theta(\xi) - \xi)' \leq C|y_N^\theta - x_i|$$

$$(3.145) \quad (|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha})' = |Ly_{N-i}^\theta(\xi) - \xi|^{-\alpha}(Ly_{N-i}^\theta(\xi) - \xi)' \leq |y_N^\theta - x_i|^{1-\alpha}$$

Finally,

$$(3.146) \quad \begin{aligned} (|Ly_{N-i}^\theta(\xi) - \xi|^{1-\alpha})'' &= (1-\alpha)|Ly_{N-i}^\theta(\xi) - \xi|^{-\alpha}(Ly_{N-i}^\theta(\xi) - \xi)'' \\ &\quad + \alpha(\alpha-1)|Ly_{N-i}^\theta(\xi) - \xi|^{-1-\alpha}((Ly_{N-i}^\theta(\xi) - \xi)')^2 \quad \square \\ &\leq C(r-1)|y_N^\theta - x_i|^{-\alpha} + C|y_N^\theta - x_i|^{1-\alpha} \end{aligned}$$

By the three lemmas above, for $N/2 \leq i \leq N-1$, we have

LEMMA 3.34.

$$(3.147) \quad \begin{aligned} D_{hL}^2 P_{N-i}^\theta(x_i) &= {}_L P_{N-i}^\theta''(\xi) \quad \xi \in (x_{i-1}, x_{i+1}) \\ &\leq Ch^3|y_N^\theta - x_i|^{1-\alpha} + C(r-1)(h^3|y_N^\theta - x_i|^{-\alpha} + h^2|y_N^\theta - x_i|^{1-\alpha}) \end{aligned}$$

while $\theta h_N = y_N^\theta - x_{N-1} \leq y_N^\theta - x_i$, we have

$$(3.148) \quad \theta D_{hL}^2 P_{N-i}^\theta(x_i) \leq Ch^3|y_N^\theta - x_i|^{1-\alpha} + C(r-1)(h^2|y_N^\theta - x_i|^{1-\alpha})$$

And

LEMMA 3.35.

$$(3.149) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{{}_L Q_{N-i}^\theta(x_{i+1})u'''(\eta_{N+1}^\theta) - {}_L Q_{N-i}^\theta(x_i)u'''(\eta_N^\theta)}{h_{i+1}} \right) \leq Ch^3|y_N^\theta - x_i|^{1-\alpha}$$

And immediately with Lemma 3.16, For $N/2 \leq i \leq N-1$

$$(3.150) \quad \begin{aligned} V_{iN} &\leq C \int_{x_{N-1}}^{x_N} h^2|y - x_i|^{1-\alpha} + C(r-1)h|y - x_i|^{1-\alpha} dy \\ &\leq Ch^2h_N|T - x_i|^{1-\alpha} + C(r-1)h^2|x_N - x_i|^{1-\alpha} \\ &\leq Ch^2 + C(r-1)h^2|T - x_{i-1}|^{1-\alpha} \end{aligned}$$

Similarly with $j = N+1$. □

I_4, I_5 is easy. Similar with Lemma 3.23 and Lemma 3.8, we have

THEOREM 3.36. *There is a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For*
 $N/2 \leq i \leq N,$
(3.151)

$$I_4 = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} (T_{i-1, 2N - \lceil \frac{N}{2} \rceil + 1} + T_{i-1, 2N - \lceil \frac{N}{2} \rceil}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i, 2N - \lceil \frac{N}{2} \rceil + 1} \right) \\ \leq Ch^2$$

Proof. Similar with Lemma 3.23. In fact, let $m = 2N - \lceil \frac{N}{2} \rceil + 1$

$$(3.152) \quad \begin{aligned} & \frac{1}{h_i} (T_{i-1, l} + T_{i-1, l-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i, l} \\ &= \frac{1}{h_i} (T_{i-1, l} - T_{i, l}) + \frac{1}{h_i} (T_{i-1, l-1} - T_{i, l}) + \left(\frac{1}{h_i} - \frac{1}{h_{i+1}} \right) T_{i, l} \end{aligned}$$

While, by Lemma A.2

$$(3.153) \quad \begin{aligned} \frac{1}{h_i} (T_{i-1, l} - T_{i, l}) &= \int_{x_{l-1}}^{x_l} (u(y) - \Pi_h u(y)) \frac{|x_{i-1} - y|^{1-\alpha} - |x_i - y|^{1-\alpha}}{h_i \Gamma(2-\alpha)} dy \\ &\leq C \int_{x_{l-1}}^{x_l} h_l^2 u''(\eta) \frac{|\xi - y|^{-\alpha}}{\Gamma(1-\alpha)} dy, \quad \xi \in (x_{i-1}, x_i) \\ &\leq Ch_l^3 (2T - x_{l-1})^{\alpha/2-2} T^{-\alpha} \\ &\leq Ch_l^3 \end{aligned}$$

Thus,

$$(3.154) \quad \frac{2}{h_i + h_{i+1}} \frac{1}{h_i} (T_{i-1, l} - T_{i, l}) \leq Ch_l^2$$

For

$$(3.155) \quad \frac{1}{h_i} (T_{i-1, l-1} - T_{i, l}) = \int_0^1 -\frac{\theta(1-\theta)}{2} \frac{h_{l-1}^3 |y_{l-1}^\theta - x_{i-1}|^{1-\alpha} u''(\eta_{l-1}^\theta) - h_l^3 |y_l^\theta - x_i|^{1-\alpha} u''(\eta_l^\theta)}{h_i} d\theta$$

And Similar with Lemma 3.21, we can get

$$(3.156) \quad \frac{h_{l-1}^3 |y_{l-1}^\theta - x_{i-1}|^{1-\alpha} u''(\eta_{l-1}^\theta) - h_l^3 |y_l^\theta - x_i|^{1-\alpha} u''(\eta_l^\theta)}{(h_i + h_{i+1}) h_i} \leq Ch_l^2 |y_l^\theta - x_i|^{1-\alpha}$$

So

$$(3.157) \quad \frac{2}{h_i + h_{i+1}} \frac{1}{h_i} (T_{i-1, l-1} - T_{i, l}) \leq Ch^2$$

For the third term, by Lemma B.1, Lemma B.2 and Lemma A.2

$$(3.158) \quad \begin{aligned} \frac{2}{h_i + h_{i+1}} \frac{h_{i+1} - h_i}{h_i h_{i+1}} T_{i, l} &\leq h_i^{-3} h^2 x_i^{1-2/r} h_l C h_l^2 x_{l-1}^{\alpha/2-2} |x_l - x_i|^{1-\alpha} \\ &\leq Ch^2 \end{aligned}$$

Summarizes, we have

$$(3.159) \quad I_4 \leq Ch^2$$

□

And

LEMMA 3.37. *There is a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq i \leq N$,*

$$I_5 = \sum_{j=2N-\lceil \frac{N}{2} \rceil+2}^{2N} S_{ij} \leq \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

Proof. For $i \leq N, j \geq 2N - \lceil \frac{N}{2} \rceil + 2$, we have

$$\begin{aligned} S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - \Pi_h u(y)) D_h^2 K_y(x_i) dy \\ &\leq \int_{x_{j-1}}^{x_j} Ch^2 (2T - y)^{\alpha/2-2/r} |y - x_{i+1}|^{-1-\alpha} dy \\ &\leq Ch^2 T^{-1-\alpha} \int_{x_{j-1}}^{x_j} (2T - y)^{\alpha/2-2/r} dy \end{aligned}$$

$$\begin{aligned} \sum_{j=2N-\lceil \frac{N}{2} \rceil+2}^{2N-1} S_{ij} &\leq CT^{-1-\alpha} h^2 \int_{(2-2^{-r})T}^{x_{2N-1}} (2T - y)^{\alpha/2-2/r} dy \\ &\leq CT^{-1-\alpha} h^2 \begin{cases} \frac{1}{\alpha/2-2/r+1} T^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 > 0 \\ \ln(2^{-r}T) - \ln(h_{2N}), & \alpha/2 - 2/r + 1 = 0 \\ \frac{1}{|\alpha/2-2/r+1|} h_{2N}^{\alpha/2-2/r+1}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \\ &= \begin{cases} \frac{C}{\alpha/2-2/r+1} T^{-\alpha/2-2/r} h^2, & \alpha/2 - 2/r + 1 > 0 \\ CrT^{-1-\alpha} h^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ \frac{C}{|\alpha/2-2/r+1|} T^{-\alpha/2-2/r} h^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \end{aligned}$$

Now we can conclude a part of the theorem Theorem 3.3 at the beginning of this section.

By Lemma 3.10 Lemma 3.23 Lemma 3.24 Theorem 3.30 Lemma 3.29 Theorem 3.36 Lemma 3.37, we have

THEOREM 3.38. *there exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that for $N/2 \leq i \leq N - 1$,*

$$\begin{aligned} R_i &= I_1 + I_2 + I_3^1 + I_3^2 + I_3^3 + I_4 + I_5 \\ &\leq C(r-1)h^2 |T - x_{i-1}|^{1-\alpha} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases} \end{aligned}$$

And what we left is the case $i = N$. Fortunately, we can use the same department of R_i above, and it is symmetric. Most of the item has been esitimated by Lemma 3.10 and Theorem 3.36, we just need to consider I_3, I_4 .

THEOREM 3.39. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that*

$$(3.163) \quad I_3 = \sum_{j=\lceil \frac{N}{2} \rceil + 1}^{N-1} V_{Nj} \leq Ch^2 + C(r-1)h^2|T - x_{N-1}|^{1-\alpha}$$

Proof. **DEFINITION 3.40.** *For $N/2 \leq j < N$, Let's define*

$$(3.164) \quad y_j(x) = \left(\frac{Z_1}{h_N}(x - x_N) + Z_j \right)^r, \quad Z_j = T^{1/r} \frac{j}{N}$$

We can see that is the inverse of the function ${}_0y_{N-i}(x)$ defined in Theorem 3.30.

$$(3.165) \quad y'_j(x) = y_j^{1-1/r}(x) \frac{rZ_1}{h_N}$$

$$(3.166) \quad y''_j(x) = y_j^{1-2/r}(x) \frac{r(r-1)Z_1}{h_N}$$

With the scheme we used several times, we can get

LEMMA 3.41. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq j < N$, $\xi \in [x_{N-1}, x_{N+1}]$,*

$$(3.167) \quad h_j(\xi)^3 \leq Ch^3$$

$$(3.168) \quad (h_j^3(\xi))' \leq C(r-1)h^3$$

$$(3.169) \quad (h_j^3(\xi))'' \leq C(r-1)h^3$$

$$(3.170) \quad u''(y_j^\theta(\xi)) \leq C$$

$$(3.171) \quad (u''(y_j^\theta(\xi)))' \leq C$$

$$(3.172) \quad (u''(y_j^\theta(\xi)))'' \leq C$$

$$(3.173) \quad |\xi - y_j^\theta(\xi)|^{1-\alpha} \leq C|x_N - y_j^\theta|^{1-\alpha}$$

$$(3.174) \quad (|\xi - y_j^\theta(\xi)|^{1-\alpha})' \leq C|x_N - y_j^\theta|^{1-\alpha}$$

$$(3.175) \quad (|\xi - y_j^\theta(\xi)|^{1-\alpha})'' \leq C|x_N - y_j^\theta|^{1-\alpha} + C(r-1)|x_N - y_j^\theta|^{-\alpha}$$

LEMMA 3.42. *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For $N/2 \leq j < N$,*

$$(3.176) \quad V_{Nj} \leq Ch^2 \int_{x_{j-1}}^{x_j} |x_N - y|^{1-\alpha} + (r-1)|x_N - y|^{-\alpha} dy$$

Therefore,

$$(3.177) \quad \begin{aligned} I_3 &\leq Ch^2 \int_{\lceil \frac{N}{2} \rceil}^{N-1} |x_N - y|^{1-\alpha} + (r-1)|x_N - y|^{-\alpha} dy \\ &\leq Ch^2(|T - x_{N-1}|^{2-\alpha} + (r-1)|T - x_{N-1}|^{1-\alpha}) \end{aligned}$$

□

For $j = N$,

LEMMA 3.43.

(3.178)

$$V_{N,N} = \frac{1}{h_N^2} (T_{N-1,N-1} - 2T_{N,N} + T_{N+1,N+1}) \leq Ch^2 + C(r-1)h^2|T - x_{N-1}|^{1-\alpha}$$

Proof.

(3.179)

□

$$\begin{aligned} V_{N,N} = & \int_0^1 -\frac{\theta(1-\theta)^{2-\alpha}}{2} \frac{1}{h_N^2} (h_{N-1}^{4-\alpha} u''(y_{N-1}^\theta) - 2h_N^{4-\alpha} u''(y_N^\theta) + h_{N+1}^{4-\alpha} u''(y_{N+1}^\theta)) d\theta \\ & + \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{1}{h_N} \left(\frac{Q_{N \rightarrow N}^\theta(x_{N+1}) u'''(\eta_{N+1,1}^\theta) - Q_{N \rightarrow N}^\theta(x_i) u'''(\eta_{N,1}^\theta)}{h_N} \right) d\theta \\ & - \int_0^1 \frac{\theta^3(1-\theta)}{3!} \frac{1}{h_N} \left(\frac{Q_{N \rightarrow N}^\theta(x_N) u'''(\eta_{N,1}^\theta) - Q_{N \rightarrow N}^\theta(x_{N-1}) u'''(\eta_{N-1,1}^\theta)}{h_N} \right) d\theta \\ & - \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{1}{h_N} \left(\frac{Q_{N \rightarrow N}^\theta(x_{N+1}) u'''(\eta_{N+1,2}^\theta) - Q_{N \rightarrow N}^\theta(x_N) u'''(\eta_{N,2}^\theta)}{h_N} \right) d\theta \\ & + \int_0^1 \frac{\theta(1-\theta)^3}{3!} \frac{1}{h_N} \left(\frac{Q_{N \rightarrow N}^\theta(x_N) u'''(\eta_{N,2}^\theta) - Q_{N \rightarrow N}^\theta(x_{N-1}) u'''(\eta_{N-1,2}^\theta)}{h_N} \right) d\theta \end{aligned}$$

So combine Lemma 3.10, Theorem 3.36, Theorem 3.39, Lemma 3.43 We have

LEMMA 3.44.

$$R_N \leq C(r-1)h^2|T - x_{N-1}|^{1-\alpha} + \begin{cases} Ch^2, & \alpha/2 - 2/r + 1 > 0 \\ Ch^2 \ln(N), & \alpha/2 - 2/r + 1 = 0 \\ Ch^{r\alpha/2+r}, & \alpha/2 - 2/r + 1 < 0 \end{cases}$$

and with Theorem 3.38 we prove the Theorem 3.3

4. Convergence analysis.

4.1. Properties of some Matrices. Review subsection 2.1, we have got (2.10).

DEFINITION 4.1. We call one matrix an M matrix, which means its entries are positive on major diagonal and nonpositive on others, and strictly diagonally dominant in rows.

Now we have

LEMMA 4.2. Matrix A defined by (2.12) where (2.13) is an M matrix. And there exists a constant $C_A = C(T, \alpha, r)$ such that

$$(4.1) \quad S_i := \sum_{j=1}^{2N-1} a_{ij} \geq C_A(x_i^{-\alpha} + (2T - x_i)^{-\alpha})$$

Proof. From (2.14), we have

$$(4.2) \quad \sum_{j=1}^{2N-1} \tilde{a}_{ij} = \frac{1}{\Gamma(4-\alpha)} \left(\frac{|x_i - x_0|^{3-\alpha} - |x_i - x_1|^{3-\alpha}}{h_1} + \frac{|x_{2N} - x_i|^{3-\alpha} - |x_{2N-1} - x_i|^{3-\alpha}}{h_{2N}} \right)$$

Let

$$(4.3) \quad g(x) = g_0(x) + g_{2N}(x)$$

where

$$g_0(x) := \frac{-\kappa_\alpha}{\Gamma(4-\alpha)} \frac{|x - x_0|^{3-\alpha} - |x - x_1|^{3-\alpha}}{h_1}$$

$$g_{2N}(x) := \frac{-\kappa_\alpha}{\Gamma(4-\alpha)} \frac{|x_{2N} - x|^{3-\alpha} - |x_{2N-1} - x|^{3-\alpha}}{h_{2N}}$$

Thus

$$-\kappa_\alpha \sum_{j=1}^{2N-1} \tilde{a}_{ij} = g(x_i)$$

Then

$$(4.4) \quad S_i := \sum_{j=1}^{2N-1} a_{ij}$$

$$= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right)$$

$$= D_h^2 g_0(x_i) + D_h^2 g_{2N}(x_i)$$

When $i = 1$

$$(4.5) \quad D_h^2 g_0(x_1) = \frac{2}{h_1 + h_2} \left(\frac{1}{h_2} g_0(x_2) - \left(\frac{1}{h_1} + \frac{1}{h_2} \right) g_0(x_1) + \frac{1}{h_1} g_0(x_0) \right)$$

$$= \frac{2\kappa_\alpha}{\Gamma(4-\alpha)} \frac{h_1^{3-\alpha} + h_2^{3-\alpha} + 2h_1^{2-\alpha}h_2 - (h_1 + h_2)^{3-\alpha}}{(h_1 + h_2)h_1h_2}$$

$$= \frac{2\kappa_\alpha}{\Gamma(4-\alpha)} \frac{h_1^{3-\alpha} + h_2^{3-\alpha} + 2h_1^{2-\alpha}h_2 - (h_1 + h_2)^{3-\alpha}}{(h_1 + h_2)h_1^{1-\alpha}h_2} h_1^{-\alpha}$$

$$= \frac{2\kappa_\alpha}{\Gamma(4-\alpha)} \frac{1 + (2^r - 1)^{3-\alpha} + 2(2^r - 1) - (2^r)^{3-\alpha}}{2^r(2^r - 1)} h_1^{-\alpha}$$

548 but

$$549 \quad (4.6) \quad 1 + (2^r - 1)^{3-\alpha} + 2(2^r - 1) - (2^r)^{3-\alpha} > 0$$

550 While for $i \geq 2$

$$\begin{aligned} D_h^2 g_0(x_i) &= g_0''(\xi), \quad \xi \in (x_{i-1}, x_{i+1}) \\ &= -\kappa_\alpha \frac{|\xi - x_0|^{1-\alpha} - |\xi - x_1|^{1-\alpha}}{\Gamma(2-\alpha)h_1} \\ 551 \quad (4.7) \quad &= \frac{\kappa_\alpha}{-\Gamma(1-\alpha)} |\xi - \eta|^{-\alpha}, \quad \eta \in [x_0, x_1] \\ &\geq \frac{\kappa_\alpha}{-\Gamma(1-\alpha)} x_{i+1}^{-\alpha} \geq \frac{\kappa_\alpha}{-\Gamma(1-\alpha)} 2^{-r\alpha} x_i^{-\alpha} \end{aligned}$$

552 So

$$553 \quad (4.8) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g_0(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g_0(x_i) + \frac{1}{h_i} g_0(x_{i-1}) \right) \geq C x_i^{-\alpha}$$

554 symmetricly,

$$555 \quad (4.9) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g_{2N}(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g_{2N}(x_i) + \frac{1}{h_i} g_{2N}(x_{i-1}) \right) \geq C(\alpha, r)(2T - x_i)^{-\alpha} \quad \square$$

556 Let

$$557 \quad (4.10) \quad g(x) = \begin{cases} x, & 0 < x \leq T \\ 2T - x, & T < x < 2T \end{cases}$$

558 And define

$$559 \quad (4.11) \quad G = \text{diag}(g(x_1), \dots, g(x_{2N-1}))$$

560 Then

561 LEMMA 4.3. *The matrix $B := AG$, the major diagonal is positive, and nonpositive*
 562 *on others. And there is a constant $C_{AG}, C = C(\alpha, r)$ such that*

$$563 \quad (4.12) \quad M_i := \sum_{j=1}^{2N-1} b_{ij} \geq -C_{AG}(x_i^{1-\alpha} + (2T - x_i)^{1-\alpha}) + C \begin{cases} |T - x_{i-1}|^{1-\alpha}, & i \leq N \\ |x_{i+1} - T|^{1-\alpha}, & i \geq N \end{cases}$$

Proof.

$$564 \quad b_{ij} = a_{ij}g(x_j) = -\kappa_\alpha \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} \tilde{a}_{i+1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{a}_{i,j} + \frac{1}{h_i} \tilde{a}_{i-1,j} \right) g(x_j)$$

565 Since

$$566 \quad (4.13) \quad g(x) \equiv \Pi_h g(x)$$

by ??, we have

$$\begin{aligned}
 \tilde{M}_i &:= \sum_{j=1}^{2N-1} \tilde{b}_{ij} = \sum_{j=1}^{2N-1} \tilde{a}_{ij} g(x_j) \\
 (4.14) \quad &= \int_0^{2T} \frac{|x_i - y|^{1-\alpha}}{\Gamma(2-\alpha)} \Pi_h g(y) dy = \int_0^{2T} \frac{|x_i - y|^{1-\alpha}}{\Gamma(2-\alpha)} g(y) dy \\
 &= \frac{-2}{\Gamma(4-\alpha)} |T - x_i|^{3-\alpha} + \frac{1}{\Gamma(4-\alpha)} (x_i^{3-\alpha} + (2T - x_i)^{3-\alpha}) \\
 &:= w(x_i) = p(x_i) + q(x_i)
 \end{aligned}$$

Thus,

$$\begin{aligned}
 M_i &:= \sum_{j=1}^{2N-1} b_{ij} = \sum_{j=1}^{2N-1} a_{ij} g(x_j) \\
 (4.15) \quad &= -\kappa_\alpha \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} \tilde{M}_{i+1} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{M}_i + \frac{1}{h_i} \tilde{M}_{i-1} \right) \\
 &= D_h^2(-\kappa_\alpha p)(x_i) - \kappa_\alpha D_h^2 q(x_i)
 \end{aligned}$$

for $1 \leq i \leq N-1$, by Lemma A.1

$$\begin{aligned}
 (4.16) \quad D_h^2(-\kappa_\alpha p)(x_i) &:= -\kappa_\alpha \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} p(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) p(x_i) + \frac{1}{h_i} p(x_{i-1}) \right) \\
 (4.17) \quad &= \frac{2\kappa_\alpha}{\Gamma(2-\alpha)} |T - \xi|^{1-\alpha} \quad \xi \in (x_{i-1}, x_{i+1}) \\
 &\geq \frac{2\kappa_\alpha}{\Gamma(2-\alpha)} |T - x_{i-1}|^{1-\alpha}
 \end{aligned}$$

$$\begin{aligned}
 (4.17) \quad D_h^2(-\kappa_\alpha p)(x_N) &:= -\kappa_\alpha \frac{2}{h_N + h_{N+1}} \left(\frac{1}{h_{N+1}} p(x_{N+1}) - \left(\frac{1}{h_N} + \frac{1}{h_{N+1}} \right) p(x_N) + \frac{1}{h_N} p(x_{N-1}) \right) \\
 (4.18) \quad &= \frac{4\kappa_\alpha}{\Gamma(4-\alpha) h_N^2} h_N^{3-\alpha} \\
 &= \frac{4\kappa_\alpha}{\Gamma(4-\alpha)} (T - x_{N-1})^{1-\alpha}
 \end{aligned}$$

Symmetricly for $i \geq N$, we get

$$(4.18) \quad D_h^2(-\kappa_\alpha p)(x_i) \geq \frac{2\kappa_\alpha}{\Gamma(2-\alpha)} \begin{cases} |T - x_{i-1}|^{1-\alpha}, & i \leq N \\ |x_{i+1} - T|^{1-\alpha}, & i \geq N \end{cases}$$

Similarly, we can get

$$\begin{aligned}
 (4.19) \quad D_h^2 q(x_i) &:= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} q(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) q(x_i) + \frac{1}{h_i} q(x_{i-1}) \right) \\
 &\leq \frac{2^{r(\alpha-1)+1}}{\Gamma(2-\alpha)} (x_i^{1-\alpha} + (2T - x_i)^{1-\alpha}), \quad i = 1, \dots, 2N-1
 \end{aligned}$$

So, we get the result.

Notice that

$$(4.20) \quad x_i^{-\alpha} \geq (2T)^{-1} x_i^{1-\alpha}$$

We can get

THEOREM 4.4. *There exists a real $\lambda = \lambda(T, \alpha, r) > 0$ and $C = C(T, \alpha, r) > 0$ such that $B := A(\lambda I + G)$ is an M matrix. And*

$$(4.21) \quad M_i := \sum_{j=1}^{2N-1} b_{ij} \geq C(x_i^{-\alpha} + (2T - x_i)^{-\alpha}) + C \begin{cases} |T - x_{i-1}|^{1-\alpha}, & i \leq N \\ |x_{i+1} - T|^{1-\alpha}, & i \geq N \end{cases}$$

Proof. By Lemma 4.2 with C_A and Lemma 4.3 with C_{AG} , it's sufficient to take $\lambda = (C + 2TC_{AG})/C_A$, then

$$(4.22) \quad M_i \geq C \left((x_i^{-\alpha} + (1 - x_i)^{-\alpha}) + \begin{cases} |T - x_{i-1}|^{1-\alpha}, & i \leq N \\ |x_{i+1} - T|^{1-\alpha}, & i \geq N \end{cases} \right) \quad \square$$

4.2. Proof of Theorem 2.6. For equation

$$(4.23) \quad AU = F \Leftrightarrow A(\lambda I + G)(\lambda I + G)^{-1}U = F \quad \text{i.e.} \quad B(\lambda I + G)^{-1}U = F$$

which means

$$(4.24) \quad \sum_{j=1}^{2N-1} b_{ij} \frac{\epsilon_j}{\lambda + g(x_j)} = -\tau_i$$

where $\epsilon_i = u(x_i) - u_i$.

And if

$$(4.25) \quad \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| = \max_{1 \leq i \leq 2N-1} \left| \frac{\epsilon_i}{\lambda + g(x_i)} \right|$$

Then, since $B = A(\lambda I + G)$ is an M matrix, it is Strictly diagonally dominant. Thus,

$$\begin{aligned} |\tau_{i_0}| &= \left| \sum_{j=1}^{2N-1} b_{i_0,j} \frac{\epsilon_j}{\lambda + g(x_j)} \right| \\ &\geq b_{i_0,i_0} \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| - \sum_{j \neq i_0} |b_{i_0,j}| \left| \frac{\epsilon_j}{\lambda + g(x_j)} \right| \\ &\geq b_{i_0,i_0} \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| - \sum_{j \neq i_0} |b_{i_0,j}| \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| \\ &= \sum_{j=1}^{2N-1} b_{i_0,j} \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| \\ &= M_{i_0} \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| \end{aligned} \quad (4.26)$$

By Theorem 2.5 and Theorem 4.4,

We know that there exists constants $C_1(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)}, \|f\|_{\beta}^{(\alpha/2)})$, and $C_2(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that

$$(4.27) \quad \left| \frac{\epsilon_i}{\lambda + g(x_i)} \right| \leq \left| \frac{\epsilon_{i_0}}{\lambda + g(x_{i_0})} \right| \leq C_1 h^{\min\{\frac{r\alpha}{2}, 2\}} + C_2(r-1)h^2$$

602 as $\lambda + g(x_i) \leq \lambda + T$

603 So, we can get

$$604 \quad (4.28) \quad |\epsilon_i| \leq C(\lambda + T)h^{\min\{\frac{\alpha}{2}, 2\}}$$

605 The convergency has been proved.

606 Remarks:

5. Experimental results.

5.1. $f \equiv 1$.

5.2. $f = x^\gamma, \gamma < 0$. Appendix A. Approximate of difference quotients.

LEMMA A.1. *If $g(x) \in C^2(\Omega)$, there exists $\xi \in (x_{i-1}, x_{i+1})$ such that*

$$(A.1) \quad D_h^2 g(x_i) := \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = g''(\xi), \quad \xi \in (x_{i-1}, x_{i+1})$$

$$(A.2) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \int_{x_{i-1}}^{x_i} g''(y)(y - x_{i-1}) dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} g''(y)(x_{i+1} - y) dy \right)$$

And if $g(x) \in C^4(\Omega)$, then

$$(A.3) \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = g''(x_i) + \frac{h_{i+1} - h_i}{3} g'''(x_i) + \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 g''''(\eta_1) + h_{i+1}^3 g''''(\eta_2))$$

where $\eta_1 \in [x_{i-1}, x_i], \eta_2 \in [x_i, x_{i+1}]$.

Proof.

$$g(x_{i-1}) = g(x_i) - (x_i - x_{i-1})g'(x_i) + \frac{(x_i - x_{i-1})^2}{2} g''(\xi_1), \quad \xi_1 \in (x_{i-1}, x_i)$$

$$g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i)g'(x_i) + \frac{(x_{i+1} - x_i)^2}{2} g''(\xi_2), \quad \xi_2 \in (x_i, x_{i+1})$$

Substitute them in the left side of (A.1), we have

$$\frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = \frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2)$$

Now, using **intermediate value theorem**, there exists $\xi \in [\xi_1, \xi_2]$ such that

$$\frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2) = g''(\xi)$$

For the second equation, similarly

$$g(x_{i-1}) = g(x_i) - (x_i - x_{i-1})g'(x_i) + \int_{x_{i-1}}^{x_i} g''(y)(y - x_{i-1}) dy \\ g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i)g'(x_i) + \int_{x_i}^{x_{i+1}} g''(y)(x_{i+1} - y) dy$$

And the last equation can be obtained by

$$\begin{aligned} g(x_{i-1}) &= g(x_i) - h_i g'(x_i) + \frac{h_i^2}{2} g''(x_i) - \frac{h_i^3}{3!} g'''(x_i) + \int_{x_{i-1}}^{x_i} g''''(y) \frac{(y - x_{i-1})^3}{3!} dy \\ g(x_{i+1}) &= g(x_i) + h_{i+1} g'(x_i) + \frac{h_{i+1}^2}{2} g''(x_i) + \frac{h_{i+1}^3}{3!} g'''(x_i) + \int_{x_i}^{x_{i+1}} g''''(y) \frac{(x_{i+1} - y)^3}{3!} dy \end{aligned}$$

Especially,

$$\begin{aligned} \int_{x_{i-1}}^{x_i} g''''(y) \frac{(y - x_{i-1})^3}{3!} dy &= \frac{h_i^4}{4!} g''''(\eta_1) \\ \int_{x_i}^{x_{i+1}} g''''(y) \frac{(x_{i+1} - y)^3}{3!} dy &= \frac{h_{i+1}^4}{4!} g''''(\eta_2) \end{aligned} \quad (A.4)$$

where $\eta_1 \in (x_{i-1}, x_i)$, $\eta_2 \in (x_i, x_{i+1})$. Subsitute them to the left side of (A.3), we can get the result. \square

LEMMA A.2. Denote $y_j^\theta = (1 - \theta)x_{j-1} + \theta x_j$, $\theta \in (0, 1)$,

$$u(y_j^\theta) - \Pi_h u(y_j^\theta) = -\frac{\theta(1 - \theta)}{2} h_j^2 u''(\xi), \quad \xi \in (x_{j-1}, x_j)$$

(A.6)

$$u(y_j^\theta) - \Pi_h u(y_j^\theta) = -\frac{\theta(1 - \theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1 - \theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1 - \theta)^2 u'''(\eta_2))$$

where $\eta_1 \in (x_{j-1}, y_j^\theta)$, $\eta_2 \in (y_j^\theta, x_j)$.

Proof. By Taylor expansion, we have

$$u(x_{j-1}) = u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(\xi_1), \quad \xi_1 \in (x_{j-1}, y_j^\theta)$$

$$u(x_j) = u(y_j^\theta) + (1 - \theta) h_j u'(y_j^\theta) + \frac{(1 - \theta)^2 h_j^2}{2!} u''(\xi_2), \quad \xi_2 \in (y_j^\theta, x_j)$$

Thus

$$\begin{aligned} u(y_j^\theta) - \Pi_h u(y_j^\theta) &= u(y_j^\theta) - (1 - \theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1 - \theta)}{2} h_j^2 (\theta u''(\xi_1) + (1 - \theta) u''(\xi_2)) \\ &= -\frac{\theta(1 - \theta)}{2} h_j^2 u''(\xi), \quad \xi \in [\xi_1, \xi_2] \end{aligned}$$

The second equation is similar,

$$\begin{aligned} u(x_{j-1}) &= u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(y_j^\theta) - \frac{\theta^3 h_j^3}{3!} u'''(\eta_1) \\ u(x_j) &= u(y_j^\theta) + (1 - \theta) h_j u'(y_j^\theta) + \frac{(1 - \theta)^2 h_j^2}{2!} u''(y_j^\theta) + \frac{(1 - \theta)^3 h_j^3}{3!} u'''(\eta_2) \end{aligned}$$

where $\eta_1 \in (x_{j-1}, y_j^\theta)$, $\eta_2 \in (y_j^\theta, x_j)$. Thus \square

$$\begin{aligned} u(y_j^\theta) - \Pi_h u(y_j^\theta) &= u(y_j^\theta) - (1 - \theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1 - \theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1 - \theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1 - \theta)^2 u'''(\eta_2)) \end{aligned}$$

LEMMA A.3. For $x \in [x_{j-1}, x_j]$

$$(A.7) \quad |u(x) - \Pi_h u(x)| = \left| \frac{x_j - x}{h_j} \int_{x_{j-1}}^x u'(y) dy - \frac{x - x_{j-1}}{h_j} \int_x^{x_j} u'(y) dy \right| \\ \leq \int_{x_{j-1}}^{x_j} |u'(y)| dy$$

If $x \in [0, x_1]$, with Corollary 2.4, we have

$$(A.8) \quad |u(x) - \Pi_h u(x)| \leq \int_0^{x_1} |u'(y)| dy \leq \int_0^{x_1} C y^{\alpha/2-1} dy \leq C \frac{2}{\alpha} x_1^{\alpha/2}$$

Similarly, if $x \in [x_{2N-1}, 1]$, we have

$$(A.9) \quad |u(x) - \Pi_h u(x)| \leq C \frac{2}{\alpha} (2T - x_{2N-1})^{\alpha/2} = C \frac{2}{\alpha} x_1^{\alpha/2}$$

LEMMA A.4.

$$(A.10) \quad b^{1-\theta} |a^\theta - b^\theta| \leq |a - b| \quad (\text{also } a^{1-\theta} |a^\theta - b^\theta| \leq |a - b|), \quad a, b \geq 0, \theta \in [0, 1]$$

Appendix B. Inequality. For convenience, we use the notation and \simeq . That $x_1 \simeq y_1$, means that $c_1 x_1 \leq y_1 \leq C_1 x_1$ for some constants c_1 and C_1 that are independent of mesh parameters.

LEMMA B.1.

$$(B.1) \quad h_i \simeq \begin{cases} h x_i^{1-1/r}, & 1 \leq i \leq N \\ h(2T - x_i)^{1-1/r}, & N < i \leq 2N - 1 \end{cases}$$

Since, $i^r - (i-1)^r \simeq i^{r-1}$, for $i \geq 1$

LEMMA B.2. There is a constant $C = 2^{|r-2|} r(r-1) T^{2/r}$ such that for all $i \in \{1, 2, \dots, 2N-1\}$

$$(B.2) \quad |h_{i+1} - h_i| \leq C h^2 \begin{cases} x_i^{1-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{1-2/r}, & N < i \leq 2N-1 \end{cases}$$

Proof.

$$h_{i+1} - h_i = \begin{cases} T \left(\left(\frac{i+1}{N} \right)^r - 2 \left(\frac{i}{N} \right)^r + \left(\frac{i-1}{N} \right)^r \right), & 1 \leq i \leq N-1 \\ 0, & i = N \\ -T \left(\left(\frac{2N-i-1}{N} \right)^r - 2 \left(\frac{2N-i}{N} \right)^r + \left(\frac{2N-i+1}{N} \right)^r \right), & N+1 \leq i \leq 2N-1 \end{cases}$$

For $i = 1$,

$$h_2 - h_1 = T(2^r - 2) \left(\frac{1}{N} \right)^r = (2^r - 2) T^{2/r} h^2 x_1^{1-2/r}$$

For $2 \leq i \leq N-1$, by Lemma A.1, we have

$$\begin{aligned} h_{i+1} - h_i &= r(r-1)T N^{-2}\eta^{r-2}, \quad \eta \in [\frac{i-1}{N}, \frac{i+1}{N}] \\ &= C(r-1)h^2x_i^{1-2/r} \end{aligned}$$

Summarizes the inequalities, we can get

$$(B.3) \quad |h_{i+1} - h_i| \leq 2^{|r-2|}r(r-1)T^{2/r}h^2 \begin{cases} x_i^{1-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{1-2/r}, & N < i \leq 2N-1 \end{cases} \quad \square$$

Appendix C. Proofs of some technical details.

Additional proof of Theorem 3.1. For $2 \leq i \leq N-1$,

$$\begin{aligned} &\frac{2}{h_i + h_{i+1}}(h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \\ &\leq C \frac{2}{h_i + h_{i+1}}(h_i^3 x_{i-1}^{-2-\alpha/2} + h_{i+1}^3 x_i^{-2-\alpha/2}) \\ &\leq 2C(h_i^2 x_{i-1}^{-2-\alpha/2} + h_{i+1}^2 x_i^{-2-\alpha/2}) \end{aligned}$$

There is a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{\alpha/2})$ such that

$$\frac{2}{h_i + h_{i+1}}(h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \leq Ch^2 x_i^{-\alpha/2-2/r}, \quad 2 \leq i \leq N-1$$

For $i = 1$, by (A.4)

$$\begin{aligned} &\frac{1}{4!} \frac{2}{h_1 + h_2}(h_1^3 f''(\eta_1) + h_2^3 f''(\eta_2)) \\ &= \frac{2}{h_1 + h_2} \left(\frac{1}{h_1} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy + \frac{1}{4!} h_2^3 f''(\eta_2) \right) \end{aligned}$$

We have proved above that

$$\frac{2}{h_1 + h_2} h_2^3 f''(\eta_2) \leq Ch^2 x_1^{-\alpha/2-2/r}$$

and we can get

$$\begin{aligned} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy &\leq C \frac{1}{3!} \int_0^{x_1} y^{1-\alpha/2} dy \\ &= C \frac{1}{3!(2-\alpha/2)} x_1^{2-\alpha/2} \end{aligned}$$

so

$$\frac{2}{h_1 + h_2} \frac{1}{h_1} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy = \frac{C2^{1-r}}{3!(2-\alpha/2)} x_1^{-\alpha/2} = \frac{C2^{1-r}}{3!(2-\alpha/2)} T^{2/r} h^2 x_1^{-\alpha/2-2/r}$$

And for $i = N$, we have

$$\begin{aligned} & \frac{2}{h_N + h_{N+1}} (h_N^3 f''(\eta_1) + h_{N+1}^3 f''(\eta_2)) \\ &= h_N^2 (f''(\eta_1) + f''(\eta_2)) \\ &\leq r^2 T^{2/r} h^2 x_N^{2-2/r} 2C x_{N-1}^{-2-\alpha/2} \\ &\leq 2r^2 T^{2/r} C 2^{-r(-2-\alpha/2)} h^2 x_N^{-\alpha/2-2/r} \end{aligned}$$

Finally, $N + 1 \leq i \leq 2N - 1$ is symmetric to the first half of the proof, so we can conclude that \square

$$\frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \leq Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T - x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N - 1 \end{cases}$$

LEMMA C.1. *By a standard error estimate for linear interpolation, and Corollary 2.4, There is a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ for $2 \leq j \leq N$,*

$$(C.1) \quad |u(y) - \Pi_h u(y)| \leq Ch^2 y^{\alpha/2-2/r}, \quad \text{for } y \in [x_{j-1}, x_j]$$

symmetricly, for $N < j \leq 2N - 1$, we have

$$(C.2) \quad |u(y) - \Pi_h u(y)| \leq Ch^2 (2T - y)^{\alpha/2-2/r}$$

LEMMA C.2. *There is a constant $C = C(\alpha, r)$ such that for all $1 \leq i < N/2$, $\max\{2i + 1, i + 3\} \leq j \leq 2N$, we have*

$$(C.3) \quad D_h^2 K_y(x_i) \leq C \frac{y^{-1-\alpha}}{\Gamma(-\alpha)}, \quad y \in [x_{j-1}, x_j]$$

Proof. Since $y \geq x_{j-1} > x_{i+1}$, by Lemma A.1, if $j - 1 > i + 1$ \square

$$\begin{aligned} D_h^2 K_y(x_i) &= K_y''(\xi) = \frac{|y - \xi|^{-1-\alpha}}{\Gamma(-\alpha)}, \quad \xi \in (x_{i-1}, x_{i+1}) \\ &\leq \frac{(y - x_{i+1})^{-1-\alpha}}{\Gamma(-\alpha)} \\ &\leq (1 - (\frac{2}{3})^r)^{-1-\alpha} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} \end{aligned}$$

LEMMA C.3. *There is a constant $C = C(\alpha, r)$ such that for all $3 \leq i \leq N, k = \lceil \frac{i}{2} \rceil$, $1 \leq j \leq k - 1$ and $y \in [x_{j-1}, x_j]$, we have*

$$(C.4) \quad D_h^2 K_y(x_i) \leq C \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)}$$

Proof. Since $y \leq x_j < x_{i-1}$, by Lemma A.1, \square

$$\begin{aligned} D_h^2 K_y(x_i) &= \frac{|\xi - y|^{-1-\alpha}}{\Gamma(-\alpha)}, \quad \xi \in (x_{i-1}, x_{i+1}) \\ &\leq \frac{(x_{i-1} - x_j)^{-1-\alpha}}{\Gamma(-\alpha)} \leq \frac{(x_{i-1} - x_{k-1})^{-1-\alpha}}{\Gamma(-\alpha)} \\ &\leq ((\frac{2}{3})^r - (\frac{1}{2})^r)^{-1-\alpha} \frac{x_i^{-1-\alpha}}{\Gamma(-\alpha)} \end{aligned}$$

705

706 LEMMA C.4. While $0 \leq i < N/2$, By Lemma A.3

$$\begin{aligned}
|T_{i1}| &\leq C \int_0^{x_1} x_1^{\alpha/2} \frac{|x_i - y|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\
&= C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2} |x_i^{2-\alpha} - |x_i - x_1|^{2-\alpha}| \\
&\leq C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2+2-\alpha} = C \frac{1}{\Gamma(3-\alpha)} x_1^{2-\alpha/2} \quad 0 < 2-\alpha < 1
\end{aligned}
\tag{C.5}$$

708 For $2 \leq j \leq N$, by Lemma A.2 and Corollary 2.4

$$\begin{aligned}
|T_{ij}| &\leq \frac{C}{4} \int_{x_{j-1}}^{x_j} h_j^2 x_{j-1}^{\alpha/2-2} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\
&\leq \frac{C}{4\Gamma(3-\alpha)} h_j^2 x_{j-1}^{\alpha/2-2} ||x_j - x_i|^{2-\alpha} - |x_{j-1} - x_i|^{2-\alpha}|
\end{aligned}
\tag{C.6}$$

710 LEMMA C.5. There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that

$$\sum_{j=1}^3 S_{1j} \leq C h^2 x_1^{-\alpha/2-2/r}
\tag{C.7}$$

712

$$\sum_{j=1}^4 S_{2j} \leq C h^2 x_2^{-\alpha/2-2/r}
\tag{C.8}$$

714

Proof.

$$S_{1j} = \frac{2}{x_2} \left(\frac{1}{x_1} T_{0j} - \left(\frac{1}{x_1} + \frac{1}{h_2} \right) T_{1j} + \frac{1}{h_2} T_{2j} \right)$$

716 So, by Lemma C.4

$$\begin{aligned}
S_{11} &\leq \frac{2}{x_2 x_1} 4 \frac{C}{\Gamma(3-\alpha)} x_1^{2-\alpha/2} \leq C x_1^{-\alpha/2} \\
S_{12} &\leq \frac{2}{x_2 x_1} \frac{C}{4\Gamma(3-\alpha)} h_2^2 x_1^{\alpha/2-2} (x_2^{2-\alpha} + 2h_2^{2-\alpha} + h_2^{2-\alpha}) \leq C x_1^{-\alpha/2} \\
S_{13} &\leq \frac{2}{x_2 x_1} \frac{C}{4\Gamma(3-\alpha)} h_3^2 x_2^{\alpha/2-2} (x_3^{2-\alpha} + 2x_3^{2-\alpha} + h_3^{2-\alpha}) \leq C x_1^{-\alpha/2}
\end{aligned}$$

722 But

$$x_1^{-\alpha/2} = T^{2/r} h^2 x_1^{-\alpha/2-2/r}$$

724 $i = 2$ is similar. □

725

LEMMA C.6. *There exists a constant $C = C(T, r, l)$ such that For $3 \leq i \leq N -$
 $1, \lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\},$
when $\xi \in (x_{i-1}, x_{i+1}),$*

$$(C.9) \quad (h_{j-i}^3(\xi))' \leq (r-1)Ch^2x_i^{1-2/r}h_j$$

$$(C.10) \quad (h_{j-i}^4(\xi))' \leq (r-1)Ch^2x_i^{1-2/r}h_j^2$$

Proof. From (3.34)

$$(C.11) \quad y'_{j-i}(x) = y_{j-i}^{1-1/r}(x)x^{1/r-1}$$

$$(C.12) \quad y''_{j-i}(x) = \frac{1-r}{r}y_{j-i}^{1-2/r}(x)x^{1/r-2}Z_{j-i}$$

For $\xi \in (x_{i-1}, x_{i+1})$ and $2 \leq k \leq j \leq \min\{2i-1, N-1\}$, using Lemma B.1

$$\xi \simeq x_i \simeq x_j$$

$$h_{j-i}(\xi) \simeq h_j \simeq hx_j^{1-1/r} \simeq hx_i^{1-1/r}$$

$$(C.13) \quad \begin{aligned} h'_{j-i}(\xi) &= y'_{j-i}(\xi) - y'_{j-i-1}(\xi) \\ &= \xi^{1/r-1}(y_{j-i}^{1-1/r}(\xi) - y_{j-i-1}^{1-1/r}(\xi)) \end{aligned}$$

Since

$$(C.14) \quad \begin{aligned} y_{j-i}^{1-1/r}(\xi) - y_{j-i-1}^{1-1/r}(\xi) &\leq x_{j+1}^{1-1/r} - x_{j-2}^{1-1/r} \\ &= T^{1-1/r}N^{1-r}((j+1)^{r-1} - (j-2)^{r-1}) \\ &\leq C(r-1)j^{r-2}N^{1-r} \\ &= C(r-1)hx_j^{1-2/r} \end{aligned}$$

Therefore,

$$(C.15) \quad h'_{j-i}(\xi) \leq Cx_i^{1/r-1}(r-1)hx_j^{1-2/r} \simeq (r-1)hx_i^{-1/r}$$

for $l = 3, 4$

$$(C.16) \quad \begin{aligned} (h_{j-i}^l(\xi))' &= lh_{j-i}^{l-1}(\xi)h'_{j-i}(\xi) \\ &\leq Ch_{j-i}^{l-1}(\xi)(r-1)hx_i^{-1/r} \\ &\simeq Ch_j^{l-2}hx_j^{1-1/r}(r-1)hx_i^{-1/r} \\ &\simeq C(r-1)h^2x_i^{1-2/r}h_j^{l-2} \end{aligned}$$

Meanwhile, we can get

$$(C.17) \quad h_{j-i}^3(\xi) \simeq h_j^3 \leq Ch^2x_i^{2-2/r}h_j$$

$$(C.18) \quad h_{j-i}^4(\xi) \simeq h_j^4 \leq Ch^2x_i^{2-2/r}h_j^2$$

□

749

750 **LEMMA C.7.** *There exists a constant $C = C(T, r, l)$ such that For $3 \leq i \leq N -$*
 751 *$1, \lceil \frac{i}{2} \rceil \leq j \leq \min\{2i, N\},$*
 752 *when $\xi \in (x_{i-1}, x_{i+1}),$*

$$753 \quad (C.19) \quad (h_{j-i}^3(\xi))'' \leq C(r-1)h^2x_i^{-2/r}h_j$$

Proof.

$$754 \quad (C.20) \quad (h_{j-i}^3(\xi))'' = 6h_{j-i}(\xi)(h'_{j-i}(\xi))^2 + 3h_{j-i}^2(\xi)h''_{j-i}(\xi)$$

755 By (C.15)

$$756 \quad (C.21) \quad h_{j-i}(\xi)(h'_{j-i}(\xi))^2 \leq Ch_j(r-1)^2h^2x_i^{-2/r}$$

757 For the second partial

$$\begin{aligned} h''_{j-i}(\xi) &= y''_{j-i}(\xi) - y''_{j-i-1}(\xi) \\ 758 \quad (C.22) \quad &= \frac{1-r}{r}\xi^{1/r-2}(y_{j-i}^{1-2/r}(\xi)Z_{j-i} - y_{j-i-1}^{1-2/r}(\xi)Z_{j-i-1}) \\ &= \frac{1-r}{r}\xi^{1/r-2}((y_{j-i}^{1-2/r}(\xi) - y_{j-i-1}^{1-2/r}(\xi))Z_{j-i} + y_{j-i-1}^{1-2/r}(\xi)Z_1) \end{aligned}$$

759 but

$$\begin{aligned} |y_{j-i}^{1-2/r}(\xi) - y_{j-i-1}^{1-2/r}(\xi)| &\leq |x_{j+1}^{1-2/r} - x_{j-2}^{1-2/r}| \\ 760 \quad (C.23) \quad &= T^{1-2/r}N^{2-r}|(j+1)^{r-2} - (j-2)^{r-2}| \\ &\leq C|r-2|N^{2-r}j^{r-3} \\ &= C|r-2|h x_j^{1-3/r} \end{aligned}$$

761 So we can get

$$\begin{aligned} 762 \quad (C.24) \quad |h''_{j-i}(\xi)| &\leq C(r-1)x_i^{1/r-2}(|r-2|h x_i^{1-3/r}x_i^{1/r} + x_i^{1-2/r}h) \\ &\leq C(r-1)h x_i^{-1-1/r} \end{aligned}$$

763 Summarizes, we have

$$764 \quad (C.25) \quad (h_{j-i}^3(\xi))'' \leq C(r-1)h^2x_i^{-2/r}h_j \quad \square$$

765 *proof of Lemma 3.18.* From (3.34)

$$766 \quad (C.26) \quad y'_{j-i}(x) = y_{j-i}^{1-1/r}(x)x^{1/r-1}$$

$$767 \quad (C.27) \quad y''_{j-i}(x) = \frac{1-r}{r}y_{j-i}^{1-2/r}(x)x^{1/r-2}Z_{j-i}$$

768 Since

$$769 \quad y_{j-i}^\theta(\xi) \simeq x_j \simeq x_i$$

770 We have known

$$771 \quad (C.28) \quad u''(y_{j-i}^\theta(\xi)) \leq C(y_{j-i}^\theta(\xi))^{\alpha/2-2} \simeq x_j^{\alpha/2-2} \simeq x_i^{\alpha/2-2}$$

772

$$\begin{aligned}
& (u''(y_{j-i}^\theta(\xi)))' = u'''(y_{j-i}^\theta(\xi))(y_{j-i}^\theta(\xi))' \\
& \leq Cx_i^{\alpha/2-3}\xi^{1/r-1}y_{j-i}^{1-1/r}(\xi) \\
& \simeq x_i^{\alpha/2-3}x_i^{1/r-1}x_i^{1-1/r} = Cx_i^{\alpha/2-3}
\end{aligned}
\tag{C.29}$$

774

$$\begin{aligned}
& (u''(y_{j-i}^\theta(\xi)))'' = u''''(y_{j-i}^\theta(\xi))(y_{j-i}^\theta(\xi))'^2 + u'''(y_{j-i}^\theta(\xi))y_{j-i}^{\theta''}(\xi) \\
& \leq Cx_i^{\alpha/2-4} + Cx_i^{\alpha/2-3}\frac{r-1}{r}x_i^{1-2/r}x_i^{1/r-2}Z_{|j-i|+1} \\
& \leq Cx_i^{\alpha/2-4} + C\frac{r-1}{r}x_i^{\alpha/2-3}x_i^{-1-1/r}x_i^{1/r} \\
& = Cx_i^{\alpha/2-4}
\end{aligned}
\tag{C.30} \quad \square$$

Proof of Lemma 3.19.

$$\begin{aligned}
& |y_{j-i}^\theta(\xi) - \xi| = |\theta(y_{j-i-1}(\xi) - \xi) + (1-\theta)(y_{j-i}(\xi) - \xi)| \\
& = \theta|y_{j-i-1}(\xi) - \xi| + (1-\theta)|y_{j-i}(\xi) - \xi|
\end{aligned}
\tag{C.31}$$

where $y_{j-i-1}(\xi) - \xi$ and $y_{j-i}(\xi) - \xi$ have the same sign (≥ 0 or ≤ 0), independent with ξ .

Since $|y_{j-i}(\xi) - \xi| = \text{sign}(j-i)(y_{j-i}(\xi) - \xi)$ is increasing with ξ ,

$$\left(\frac{i-1}{i}\right)^r |x_j - x_i| \leq |x_{j-1} - x_{i-1}| \leq |y_{j-i}(\xi) - \xi| \leq |x_{j+1} - x_{i+1}| \leq \left(\frac{i+1}{i}\right)^r |x_j - x_i|
\tag{C.32}$$

we have

$$|y_{j-i}(\xi) - \xi| \simeq |x_j - x_i| \tag{C.33}$$

Similarly, $|y_{j-1-i}(\xi) - \xi| \simeq |x_{j-1} - x_i|$. Thus, with (C.31), (C.33) and (3.32) we get

$$|y_{j-i}^\theta(\xi) - \xi| \simeq |y_j^\theta - x_i| \tag{C.34}$$

Next, since $|y_{j-i}^\theta(\xi) - \xi| = \text{sign}(j-i-1+\theta)(y_{j-i}^\theta(\xi) - \xi)$, so we can derivate it.

$$(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})' = (\alpha-1)|y_{j-i}^\theta(\xi) - \xi|^{-\alpha}(y_{j-i}^\theta(\xi))' - 1| \tag{C.35}$$

While, similar with (C.31), we have

$$(y_{j-i}^\theta(\xi))' - 1 = (1-\theta)|y_{j-i-1}'(\xi) - 1| + \theta|y_{j-i}'(\xi) - 1| \tag{C.36}$$

By Lemma A.4 and (C.33), we have

$$\begin{aligned}
& |y_{j-i}'(\xi) - 1| = \xi^{1/r-1}|y_{j-i}^{1-1/r}(\xi) - \xi^{1-1/r}| \\
& \leq \xi^{-1}|y_{j-i}(\xi) - \xi| \\
& \simeq x_i^{-1}|x_j - x_i|
\end{aligned}
\tag{C.37}$$

So similar with (C.34), we can get

$$|(y_{j-i}^\theta(\xi))' - 1| \leq Cx_i^{-1}|y_j^\theta - x_i| \tag{C.38}$$

Combine with (C.34), we get

$$(C.39) \quad |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'| \leq C|y_j^\theta - x_i|^{-\alpha} x_i^{-1} |y_j^\theta - x_i| = C|y_j^\theta - x_i|^{1-\alpha} x_i^{-1}$$

Finally, we have

$$(C.40) \quad (|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'' = \alpha(\alpha-1)|y_{j-i}^\theta(\xi) - \xi|^{-\alpha-1}((y_{j-i}^\theta(\xi))' - 1)^2 \\ + \text{sign}(j-i-1+\theta)(1-\alpha)|y_{j-i}^\theta(\xi) - \xi|^{-\alpha}(y_{j-i}^\theta(\xi))''$$

For

$$(C.41) \quad (y_{j-i}^\theta(\xi))'' = (1-\theta)y_{j-i-1}''(\xi) + \theta y_{j-i}''(\xi)$$

and

$$(C.42) \quad y_{j-i}''(\xi) = \frac{1-r}{r} y_{j-i}^{1-2/r}(x) x^{1/r-2} Z_{j-i} \\ \simeq \frac{1-r}{r} x_j^{1-2/r} x_i^{1/r-2} Z_{j-i}$$

while by Lemma A.4

$$(C.43) \quad |Z_{j-i}| = |x_j^{1/r} - x_i^{1/r}| \leq |x_j - x_i| x_i^{1/r-1}$$

we have

$$(C.44) \quad |y_{j-i}''(\xi)| \leq C(r-1)x_i^{-2}|x_j - x_i|$$

Therefore

$$(C.45) \quad |(y_{j-i}^\theta(\xi))''| \leq C(r-1)x_i^{-2}|y_j^\theta - x_i|$$

Then, combine with (C.38),

$$(C.46) \quad |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})''| \leq C|y_j^\theta - x_i|^{1-\alpha} x_i^{-2} \quad \square$$

proof of Lemma 3.21. For $\lceil \frac{i}{2} \rceil \leq j \leq \min\{2i-1, N-1\}$

$$(C.47) \quad \frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \\ = \frac{Q_{j-i}^\theta(x_{i+1}) - Q_{j-i}^\theta(x_i)}{h_{i+1}} u'''(\eta_{j+1}^\theta) + Q_{j-i}^\theta(x_i) \frac{u'''(\eta_{j+1}^\theta) - u'''(\eta_j^\theta)}{h_{i+1}}$$

Since mean value theorem

$$(C.48) \quad \frac{Q_{j-i}^\theta(x_{i+1}) - Q_{j-i}^\theta(x_i)}{h_{i+1}} = Q_{j-i}^{\theta \prime}(\xi), \quad \xi \in (x_i, x_{i+1})$$

From (3.41) and Leibniz rule, by Lemma C.6 and Lemma 3.19, we have

$$(C.49) \quad |Q_{j-i}^{\theta \prime}(\xi)| \leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{1-2/r} h_j^2$$

815 And by Definition 3.14 and Lemma B.1

$$816 \quad (C.50) \quad Q_{j-i}^\theta(x_i) = h_j^4 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} \simeq Ch^2 x_i^{2-2/r} \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} h_j^2$$

817 With $\eta_j^\theta \in (x_{j-1}, x_j)$

$$818 \quad u'''(\eta_{j+1}^\theta) \leq C(\eta_{j+1}^\theta)^{\alpha/2-3} \simeq x_j^{\alpha/2-3} \simeq x_i^{\alpha/2-3}$$

819 and

$$\begin{aligned} \frac{u'''(\eta_{j+1}^\theta) - u'''(\eta_j^\theta)}{h_{i+1}} &= u''''(\eta) \frac{\eta_{j+1}^\theta - \eta_j^\theta}{h_{i+1}} \\ 820 \quad &\leq C\eta^{\alpha/2-4} \frac{x_{j+1} - x_{j-1}}{h_{i+1}} = C\eta^{\alpha/2-4} \frac{h_{j+1} + h_j}{h_{i+1}} \\ &\simeq x_j^{\alpha/2-4} \simeq x_i^{\alpha/2-4} \end{aligned}$$

821 So we have

$$\begin{aligned} &\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \\ 822 \quad (C.51) \quad &\leq Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{1-2/r} h_j^2 x_i^{\alpha/2-3} + Ch^2 x_i^{2-2/r} \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} h_j^2 x_{j-1}^{\alpha/2-4} \\ &= Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j^2 \end{aligned}$$

823 while $h_j \simeq h_i$, substitute into the inequality above, we get the goal

$$\begin{aligned} &\frac{2}{h_i + h_{i+1}} \left(\frac{Q_{j-i}^\theta(x_{i+1})u'''(\eta_{j+1}^\theta) - Q_{j-i}^\theta(x_i)u'''(\eta_j^\theta)}{h_{i+1}} \right) \\ 824 \quad (C.52) \quad &\leq \frac{1}{h_i} Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j h_i \\ &= Ch^2 \frac{|y_j^\theta - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} x_i^{\alpha/2-2-2/r} h_j \end{aligned}$$

825 While, the later is similar. □

826

827 **LEMMA C.8.** *There exists a constant $C = C(T, r)$ such that For $N/2 \leq i \leq N-1$,*
 828 *$N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1$, $l = 3, 4$, $\xi \in (x_{i-1}, x_{i+1})$, we have*

$$829 \quad (C.53) \quad h_{j-i}^l(\xi) \leq Ch_j^l \leq Ch^2 h_j^{l-2}$$

$$830 \quad (C.54) \quad (h_{j-i-1}^l(\xi))' \leq C(r-1)h^2 h_j^{l-2}$$

$$831 \quad (C.55) \quad (h_{j-i}^3(\xi))'' \leq C(r-1)h^2 h_j$$

Proof.

$$\begin{aligned} 832 \quad (C.56) \quad &(h_{j-i}(\xi))' = y_{j-i}'(\xi) - y_{j-i-1}'(\xi) \\ &= \xi^{1/r-1}((2T - y_{j-i}(\xi))^{1-1/r} - (2T - y_{j-i-1}(\xi))^{1-1/r}) \leq 0 \end{aligned}$$

833 Thus,

$$834 \quad (C.57) \quad Ch_j \leq h_{j+1} \leq h_{j-i}(\xi) \leq h_{j-i}(x_{i-1}) = h_{j-1} \leq Ch_j$$

835 So as $4^{-r}T \leq 2T - x_j \leq T, 2^{-r}T \leq x_i \leq T$, we have

$$836 \quad (C.58) \quad h_{j-i}^l(\xi) \leq Ch_j^l \leq Ch^2(2T - x_j)^{2-2/r} h_j^{l-2} \leq Ch^2 h_j^{l-2}$$

837 Since

$$\begin{aligned} & |(2T - y_{j-i}(\xi))^{1-1/r} - (2T - y_{j-i-1}(\xi))^{1-1/r}| \\ 838 \quad (C.59) \quad &= |(Z_{2N-(j-i)} - \xi^{1/r})^{r-1} - (Z_{2N-(j-1-i)} - \xi^{1/r})^{r-1}| \\ &= (r-1)Z_1(Z_{2N-(j-i-\gamma)} - \xi^{1/r})^{r-2} \quad \gamma \in [0, 1] \\ &\leq C(r-1)h(2T - x_j)^{1-2/r} \end{aligned}$$

839 we have

$$840 \quad (C.60) \quad |(h_{j-i}(\xi))'| \leq C(r-1)h(2T - x_j)^{1-2/r} x_i^{1/r-1}$$

841 And

$$\begin{aligned} & (h_{j-i}^l(\xi))' = lh_{j-i}^{l-1}(\xi)h_{j-i}'(\xi) \\ 842 \quad (C.61) \quad &\leq C(r-1)h_j^{l-1} h(2T - x_j)^{1-2/r} x_i^{1/r-1} \\ &\leq C(r-1)h^2 h_j^{l-2} (2T - x_j)^{2-3/r} x_i^{1-1/r} \\ &\leq C(r-1)h^2 h_j^{l-2} \end{aligned}$$

(C.62)

$$\begin{aligned} (h_{j-i}^3(\xi))'' &= 6h_{j-i}(\xi)(y_{j-i}'(\xi) - y_{j-i-1}'(\xi))^2 + 3h_{j-i}^2(\xi)(y_{j-i}''(\xi) - y_{j-i-1}''(\xi)) \\ &\leq C(r-1)h_j h^2 + Ch_j^2 \frac{1-r}{r} \xi^{1/r-2} ((2T - y_{j-i}(\xi))^{1-2/r} Z_{2N-(j-i)} - (2T - y_{j-i-1}(\xi))^{1-2/r} Z_{2N-(j-1-i)}) \\ 843 \quad &\leq C(r-1)h_j h^2 + C(r-1)h_j^2 (C(r-2)h(2T - x_j)^{1-3/r} Z_{2N-(j-i)} + Z_1(2T - x_{j-1})^{1-2/r}) \\ &\leq C(r-1)h_j h^2 + C(r-1)h_j^2 h = Ch^2 h_j \end{aligned} \quad \square$$

844

845 **LEMMA C.9.** *There exists a constant $C = C(T, \alpha, r, \|u\|_{\beta+\alpha}^{(-\alpha/2)})$ such that For*
846 *$N/2 \leq i \leq N-1, N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1, \xi \in (x_{i-1}, x_{i+1})$, we have*

$$847 \quad (C.63) \quad u''(y_{j-i}^\theta(\xi)) \leq C$$

$$848 \quad (C.64) \quad (u''(y_{j-i}^\theta(\xi)))' \leq C$$

$$849 \quad (C.65) \quad (u''(y_{j-i}^\theta(\xi)))'' \leq C$$

Proof.

$$850 \quad (C.66) \quad x_{j-2} \leq y_{j-i}^\theta(\xi) \leq x_{j+1} \Rightarrow 4^{-r}T \leq 2T - y_{j-i}^\theta(\xi) \leq T$$

851 Thus, for $l = 2, 3, 4$,

$$852 \quad (C.67) \quad u^{(l)}(y_{j-i}^\theta(\xi)) \leq C(2T - y_{j-i}^\theta(\xi))^{\alpha/2-l} \leq C$$

and

$$\begin{aligned} (y_{j-i}^\theta(\xi))' &= \theta y_{j-1-i}'(\xi) + (1-\theta)y_{j-i-1}'(\xi) \\ &= \xi^{1/r-1}(\theta(2T - y_{j-1-i}(\xi))^{1-1/r} + (1-\theta)(2T - y_{j-i-1}(\xi))^{1-1/r}) \\ &\leq C(2T - x_{j-2})^{1-1/r} \leq C \end{aligned}$$

With

$$Z_{2N-j-i} \leq 2T^{1/r}$$

$$\begin{aligned} (C.70) \quad (y_{j-i}^\theta(\xi))'' &= \theta y_{j-1-i}''(\xi) + (1-\theta)y_{j-i-1}''(\xi) \\ &= \frac{1-r}{r} \xi^{1/r-2} (\theta(2T - y_{j-i-1}(\xi))^{1-2/r} Z_{2N-(j-i-1)} + (1-\theta)(2T - y_{j-i}(\xi))^{1-2/r} Z_{2N-(j-i)}) \\ &\leq C(r-1) \end{aligned}$$

Therefore,

$$\begin{aligned} (C.71) \quad (u''(y_{j-i}^\theta(\xi)))' &= u'''(y_{j-i}^\theta(\xi))(y_{j-i}^\theta(\xi))' \\ &\leq C \end{aligned}$$

$$\begin{aligned} (C.72) \quad (u''(y_{j-i}^\theta(\xi)))'' &= u'''(y_{j-i}^\theta(\xi))(y_{j-i}^\theta(\xi))'^2 + u''''(y_{j-i}^\theta(\xi))y_{j-i}^\theta(\xi)'' \\ &\leq C + C(r-1) = C \quad \square \end{aligned}$$

LEMMA C.10. *There exists a constant $C = C(T, \alpha, r)$ such that For $N/2 \leq i \leq N-1$, $N+2 \leq j \leq 2N - \lceil \frac{N}{2} \rceil + 1$, $\xi \in (x_{i-1}, x_{i+1})$*

$$(C.73) \quad |y_{j-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_j^\theta - x_i|^{1-\alpha}$$

$$(C.74) \quad |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'| \leq C|y_j^\theta - x_i|^{-\alpha}(|2T - x_i - y_j^\theta| + h_N)$$

$$(C.75) \quad |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})''| \leq C(r-1)|y_j^\theta - x_i|^{-\alpha} + C|y_j^\theta - x_i|^{-1-\alpha}(|2T - x_i - y_j^\theta| + h_N)^2$$

Proof. Since $y_{j-i-1}(\xi) > x_{j-2} \geq x_N > \xi$

$$(C.76) \quad y_{j-i}^\theta(\xi) - \xi = (1-\theta)(y_{j-1-i}(\xi) - \xi) + \theta(y_{j-i}(\xi) - \xi) > 0$$

$$\begin{aligned} (C.77) \quad (y_{j-i}(\xi) - \xi)'' &= y_{j-i}''(\xi) \\ &= \frac{1-r}{r} \xi^{1/r-2} (2T - y_{j-i}(\xi))^{1-2/r} Z_{2N-(j-i)} \leq 0 \end{aligned}$$

It's concave, so

$$(C.78) \quad y_{j-i}(\xi) - \xi \geq \min_{\xi \in \{x_{i-1}, x_{i+1}\}} y_{j-i}(\xi) - \xi = \min\{x_{j+1} - x_{i+1}, x_{j-1} - x_{i-1}\} \geq C(x_j - x_i)$$

With (C.76), we have

$$(C.79) \quad |y_{j-i}^\theta(\xi) - \xi|^{1-\alpha} \leq C|y_j^\theta - x_i|^{1-\alpha}$$

By Lemma A.4

$$(C.80) \quad \begin{aligned} |y_{j-i}'(\xi) - 1| &= \xi^{1/r-1} |(2T - y_{j-i}(\xi))^{1-1/r} - \xi^{1-1/r}| \\ &\leq \xi^{-1} |2T - y_{j-i}(\xi) - \xi| \end{aligned}$$

879

$$(C.81) \quad \begin{aligned} |2T - \xi - y_{j-i}(\xi)| &\leq |2T - x_i - x_j| + |x_i - \xi| + |x_j - y_{j-i}(\xi)| \\ &\leq |2T - x_i - x_j| + h_{i+1} + h_j \\ &\leq C(|2T - x_i - x_j| + h_N) \end{aligned}$$

With $\xi \simeq x_i \simeq 1$,

$$(C.82) \quad |y_{j-i}'(\xi) - 1| \leq C(|2T - x_i - x_j| + h_N)$$

Thus,

$$(C.83) \quad \begin{aligned} |(y_{j-i}^\theta(\xi))' - 1| &\leq (1 - \theta)|y_{j-i-1}'(\xi) - 1| + \theta|y_{j-i}'(\xi) - 1| \\ &\leq C((1 - \theta)|2T - x_i - x_{j-1}| + \theta|2T - x_i - x_j| + h_N) \\ &= C(|2T - x_i - y_j^\theta| + h_N) \end{aligned}$$

So

$$(C.84) \quad \begin{aligned} |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})'| &= |1 - \alpha||y_{j-i}^\theta(\xi) - \xi|^{-\alpha} |(y_{j-i}^\theta(\xi))' - 1| \\ &\leq C|y_j^\theta - x_i|^{-\alpha} (|2T - x_i - y_j^\theta| + h_N) \end{aligned}$$

887

(C.85)

$$(C.85) \quad \begin{aligned} |(|y_{j-i}^\theta(\xi) - \xi|^{1-\alpha})''| &\leq |1 - \alpha||y_{j-i}^\theta(\xi) - \xi|^{-\alpha} |(y_{j-i}^\theta(\xi) - \xi)''| + \alpha(\alpha - 1)|y_{j-i}^\theta(\xi) - \xi|^{-1-\alpha} (y_{j-i}^\theta(\xi) - 1)^2 \\ &\leq C(r - 1)|y_j^\theta - x_i|^{-\alpha} + C|y_j^\theta - x_i|^{-1-\alpha} (|2T - x_i - y_j^\theta| + h_N)^2 \end{aligned}$$

Acknowledgments. We would like to acknowledge the assistance of volunteers in putting together this example manuscript and supplement.

891

REFERENCES

- 892 [1] D. GILBARG AND N. S. TRUDINGER, *Poisson's Equation and the Newtonian Potential*,
893 Springer Berlin Heidelberg, Berlin, Heidelberg, 1977, pp. 50–67, [https://doi.org/10.1007/](https://doi.org/10.1007/978-3-642-96379-7_4)
894 [978-3-642-96379-7_4](https://doi.org/10.1007/978-3-642-96379-7_4), https://doi.org/10.1007/978-3-642-96379-7_4.
895 [2] X. ROS-OTON AND J. SERRA, *The dirichlet problem for the fractional laplacian: Regular-*
896 *ity up to the boundary*, Journal de Mathématiques Pures et Appliquées, 101 (2014),
897 pp. 275–302, <https://doi.org/https://doi.org/10.1016/j.matpur.2013.06.003>, [https://www.](https://www.sciencedirect.com/science/article/pii/S0021782413000895)
898 [sciencedirect.com/science/article/pii/S0021782413000895](https://www.sciencedirect.com/science/article/pii/S0021782413000895).