Assembly and network stability of planktonic microorganisms under the salinity

gradient: a case study from Lena River estuary to Laptev Sea, Arctic

Qian Liu^{a,b}, Yan Li^a, Hualong Wang^{a,*}, Guipeng Yang^{b,*}, Jinjun Kan^e, Mengyao Yang^a, Xiaowen Yu^a, Cui

Guo^a, Min Wang^{a,c}, Wei Wang^d, Qingli Zhang^d, Jiancheng Zhu^d, Xianyong Zhao^d, Yong Jiang^{a,c,*}

^aCollege of Marine Life Science & Institute of Evolution and Marine Biodiversity, Ocean University of

China, Qingdao 266003, China

^bKey Laboratory of Marine Chemistry Theory and Technology, Frontiers Science Center for Deep

Ocean Multispheres and Earth System, Ministry of Education, Ocean University of China, Qingdao,

China.

^cKey Lab of Polar Oceanography and Global Ocean Change, Ocean University of China,

Qingdao266003, China

^dYellow Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, Qingdao 266071,

China

^eMicrobiology Division, Stroud Water Research Center, Avondale, PA 19311, USA

* Co-corresponding author

E-mail address: yongjiang@ouc.edu.cn (Y. Jiang), wanghualong@ouc.edu.cn

(H. Wang), gpyang@mail.ouc.edu.cn (G. Yang)

ORCID: https://orcid.org/0000-0001-6055-488X




Figure S1. The rarefaction curves of prokaryotes (A) and microeukaryotes (B).




Figure S2. Bray-Curtis dissimilarity of prokaryotes (A) and microeukaryotes (B).




Figure S3. Univariate Spearman correlations between biotic (abundant eukaryotes, prokaryotes) and abiotic data (temperature and salinity).