

Dimension Convergence Model: A Resolution-Dependent Framework for Perceived Dimensionality under Finite Observation

Abstract

Classical geometry treats dimension as an intrinsic property of objects. However, empirical observation suggests that perceived dimensionality depends on observational resolution: distant objects appear point-like, thin structures appear one-dimensional, and planar surfaces appear two-dimensional despite existing in three-dimensional space.

This work proposes the Dimension Convergence Model, in which dimensionality is treated as an observer-dependent, resolution-constrained quantity. A distinction is introduced between ontological dimension and perceived dimension, and a formal framework is constructed based on finite-resolution measurement and observable information.

Unlike classical fractal dimension, which is typically defined in asymptotic scaling regimes, this model focuses on finite-resolution observational conditions, incorporating anisotropic object structure and detectability constraints. The framework provides a structured interpretation of dimensional reduction and highlights its relevance in both physical observation and complex system modeling.

1 Introduction

Dimension is traditionally defined as the minimum number of coordinates required to specify a point in space:

- 0D — point
- 1D — line
- 2D — surface
- 3D — volume

This definition assumes that dimension is an intrinsic property of objects.

However, empirical observations suggest that perceived dimensionality depends on observational resolution:

- distant stars appear as point sources
- thin wires appear one-dimensional
- surfaces appear two-dimensional

This work proposes a framework in which dimension is treated as an observable quantity under finite resolution.

2 Ontological and Perceived Dimension

We distinguish two notions:

2.1 Ontological Dimension

The intrinsic dimensionality of an object in physical space:

$$D_{ont} = 3$$

2.2 Perceived Dimension

The effective dimensionality observable under finite resolution:

- distant star $\rightarrow D_p \approx 0$
- thin wire $\rightarrow D_p \approx 1$
- planar surface $\rightarrow D_p \approx 2$

$$0 \leq D_p \leq D_{ont}$$

3 Measurement Framework

3.1 Multi-Channel Resolution

An observer is modeled as having multiple measurement channels:

$$\{R_1, R_2, \dots, R_n\}$$

Each channel has resolution R_i and weight w_i .

We define effective resolution:

$$R_{\text{eff}} = \left(\sum_{i=1}^n w_i R_i^{-p} \right)^{-1/p}$$

This formulation corresponds to a standard L_p -norm aggregation and is adopted here as a convenient representation of multi-channel resolution rather than a novel construction.

3.2 Anisotropic Object Structure

Objects are characterized by directional scales:

$$\mathbf{S} = (S_1, S_2, \dots, S_{D_{ont}})$$

This allows representation of anisotropic structures such as wires and surfaces.

3.3 Observable Structure Count

We define:

$$N_{\text{geom}} = \prod_{i=1}^{D_{\text{ont}}} \max\left(1, \frac{S_i}{R_{\text{eff}}}\right)$$

Here N_{geom} is interpreted as a continuous measure of observable structural capacity rather than a discrete count, allowing smooth transitions under finite resolution.

3.4 Detectability Constraint

We introduce a detectability factor:

$$f(R_{\text{eff}}) = \frac{1}{1 + (R_{\text{eff}}/R_c)^\alpha}$$

The effective observable structure becomes:

$$N_{\text{eff}} = N_{\text{geom}} \cdot f(R_{\text{eff}})$$

4 Resolution-Dependent Dimension

We define perceived dimension as:

$$D_p = \begin{cases} 0, & R_{\text{eff}} \geq S_{\text{max}} \text{ or } N_{\text{eff}} < 1 \\ \frac{\log N_{\text{eff}}}{\log(S_{\text{max}}/R_{\text{eff}})}, & \text{otherwise} \end{cases}$$

where $S_{\text{max}} = \max(S_i)$.

5 Limiting Behavior

- As $R_{\text{eff}} \rightarrow 0$: $D_p \rightarrow D_{\text{ont}}$
- As $R_{\text{eff}} \rightarrow \infty$: $D_p \rightarrow 0$

6 Examples

6.1 Thin Wire

If $S_1 \gg S_2, S_3$ and $R_{\text{eff}} > S_2, S_3$:

$$D_p \approx 1$$

6.2 Planar Surface

If $S_1, S_2 \gg S_3$ and $R_{\text{eff}} > S_3$:

$$D_p \approx 2$$

6.3 Sphere

If $S_1 = S_2 = S_3$, the object remains effectively three-dimensional until resolution approaches object scale, after which:

$$D_p \rightarrow 0$$

7 Relation to Existing Frameworks

This work relates to fractal dimension, coarse-graining, and information theory. It does not replace these frameworks but provides a finite-resolution, anisotropic extension emphasizing observer-dependent dimensional perception. .

8 Implications

8.1 Physical Observation

Dimensionality depends on measurement resolution.

8.2 Measurement Systems

Different observers may assign different dimensions.

8.3 Complex Systems

In complex systems, the effective dimensionality of a system is often significantly higher than the dimensionality of the models used to describe it. Real-world systems such as healthcare, economic networks, and social structures involve many interacting variables, producing high informational dimensionality.

However, decision-making frameworks frequently operate on reduced representations, such as single metrics, binary classifications, or simplified models. Within the present framework, this can be interpreted as a consequence of finite observational resolution: only a limited subset of the system's structure is accessible, resulting in a reduced perceived dimension.

This mismatch between the intrinsic complexity of a system and the dimensionality of its representation may lead to information loss, suboptimal decisions, and structural inefficiencies. The Dimension Convergence Model provides a conceptual lens for understanding how such dimensional mismatch arises from resolution constraints rather than purely from modeling choices.

9 Limitations

- Simplified detectability model
- Parameters R_c and α require empirical grounding
- Framework is interpretive and operational rather than fundamental

10 Conclusion

The Dimension Convergence Model interprets dimensionality as a resolution-dependent observable. By incorporating anisotropic structure and detectability constraints, the model provides a structured framework for understanding dimensional perception under finite observation.

This framework also implies that dimensionality is not only subject to collapse under reduced resolution, but can emerge as resolution increases. As observational capacity improves, previously unresolvable structure becomes distinguishable, allowing perceived dimension to increase toward its ontological limit. In this sense, dimensionality can be understood as a bidirectional function of accessible information.

References

- [1] B. B. Mandelbrot, *The Fractal Geometry of Nature*, W. H. Freeman, 1982.
- [2] B. B. Mandelbrot, "How Long Is the Coast of Britain? Statistical Self-Similarity and Fractional Dimension," *Science*, 1967.
- [3] C. E. Shannon, "A Mathematical Theory of Communication," *Bell System Technical Journal*, 1948.
- [4] K. G. Wilson, "The Renormalization Group and Critical Phenomena," *Reviews of Modern Physics*, 1975.