

Ancillary qubits

Quantum Computing Minicourse ICTP-SAIFR

Stefano Carrazza[‡] and Matteo Robbiati[†]

8 April 2024


[‡] Associate Professor & Researcher, University of Milan and INFN Milan, Italy.

[†] PhD candidate, University of Milan, Italy and CERN, Switzerland.



Ancillary qubits

Ancilla qubits are extra qubits, which help a qubit system in some computations. For example:



The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

3. Let's consider now a controlled Z operation.

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

3. Let's consider now a controlled Z operation.
4. What happen if we now apply the CZ using x as control and y as target?

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

3. Let's consider now a controlled Z operation.
4. What happen if we now apply the CZ using x as control and y as target?
5. we could expect something happen on the target qubit! Not really:

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

3. Let's consider now a controlled Z operation.
4. What happen if we now apply the CZ using x as control and y as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) =$$

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

3. Let's consider now a controlled Z operation.
4. What happen if we now apply the CZ using x as control and y as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} =$$

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

3. Let's consider now a controlled Z operation.
4. What happen if we now apply the CZ using x as control and y as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} = \frac{|01\rangle - |11\rangle}{\sqrt{2}} =$$

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

3. Let's consider now a controlled Z operation.
4. What happen if we now apply the CZ using x as control and y as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} = \frac{|01\rangle - |11\rangle}{\sqrt{2}} = \frac{|0\rangle - |1\rangle}{\sqrt{2}} \otimes |1\rangle.$$

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

3. Let's consider now a controlled Z operation.
4. What happen if we now apply the CZ using x as control and y as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} = \frac{|01\rangle - |11\rangle}{\sqrt{2}} = \frac{|0\rangle - |1\rangle}{\sqrt{2}} \otimes |1\rangle.$$

6. adding an extra H to the control qubit, we will detect the state manipulation!

The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit x and an ancilla y .
2. Suppose we prepare x into a superposed state and y into the excited state:

$$|x\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |y\rangle = |1\rangle.$$

3. Let's consider now a controlled Z operation.
4. What happens if we now apply the CZ using x as control and y as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} = \frac{|01\rangle - |11\rangle}{\sqrt{2}} = \frac{|0\rangle - |1\rangle}{\sqrt{2}} \otimes |1\rangle.$$

6. adding an extra H to the control qubit, we will detect the state manipulation!

Important

This happens if the ancilla is prepared into a state which is eigenvector of the controlled operation.

Is the phase kickback useful?

Some of the most powerful and famous quantum computing algorithms make use of the phase kickback



Figure 1: Deutsch-Josza




Figure 2: Grover




Figure 3: Shor




Figure 4: HHL

All of them are proved to outperform any classical algorithm, leading to a theoretical **quantum advantage**!

It can be useful to define the phase kickback using the concept of quantum **oracle**.


1. Considering a system of qubits of state $|x\rangle$ and a set of extra qubits of state $|y\rangle$;

It can be useful to define the phase kickback using the concept of quantum **oracle**.

1. Considering a system of qubits of state $|x\rangle$ and a set of extra qubits of state $|y\rangle$;
2. in reversible computation x is called **input register** and y **output register**. In this course x will always be composed of our system of qubits and y will be ancillary qubits;


It can be useful to define the phase kickback using the concept of quantum **oracle**.

1. Considering a system of qubits of state $|x\rangle$ and a set of extra qubits of state $|y\rangle$;
2. in reversible computation x is called **input register** and y **output register**. In this course x will always be composed of our system of qubits and y will be ancillary qubits;
3. considering a function $f(x)$, acting on y , we define oracle U_f , a **black box** operator whose action on $|y\rangle$ is:



It can be useful to define the phase kickback using the concept of quantum **oracle**.


1. Considering a system of qubits of state $|x\rangle$ and a set of extra qubits of state $|y\rangle$;
2. in reversible computation x is called **input register** and y **output register**. In this course x will always be composed of our system of qubits and y will be ancillary qubits;
3. considering a function $f(x)$, acting on y , we define oracle U_f , a **black box** operator whose action on $|y\rangle$ is:



In the explicit case of the phase kickback, we can use U_f to write the action: $U_f |x\rangle |- \rangle = (-1)^x |x\rangle |- \rangle$.

It can be useful to define the phase kickback using the concept of quantum **oracle**.

1. Considering a system of qubits of state $|x\rangle$ and a set of extra qubits of state $|y\rangle$;
2. in reversible computation x is called **input register** and y **output register**. In this course x will always be composed of our system of qubits and y will be ancillary qubits;
3. considering a function $f(x)$, acting on y , we define oracle U_f , a **black box** operator whose action on $|y\rangle$ is:



In the explicit case of the phase kickback, we can use U_f to write the action: $U_f |x\rangle |-\rangle = (-1)^x |x\rangle |-\rangle$.

Extending it to a multi-dimensional case, we can write: $U_f |x\rangle |-\rangle = (-1)^{f(x)} |x\rangle |-\rangle$, whith $f(x) \in \{0, 1\}$.

Let's code!

