

# Ancillary qubits

Quantum Computing Minicourse ICTP-SAIFR

---

Stefano Carrazza<sup>‡</sup> and Matteo Robbiati<sup>†</sup>

8 April 2024


<sup>‡</sup> Associate Professor & Researcher, University of Milan and INFN Milan, Italy.

<sup>†</sup> PhD candidate, University of Milan, Italy and CERN, Switzerland.



## Ancillary qubits

Ancilla qubits are extra qubits, which help a qubit system in some computations. For example:



## The phase kickback

---

One clever trick we can implement using ancillas is the **phase kickback**.

## The phase kickback

---

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .

## The phase kickback

---

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

## The phase kickback

---

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

3. Let's consider now a controlled  $Z$  operation.

## The phase kickback

---

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

3. Let's consider now a controlled  $Z$  operation.
4. What happens if we now apply the  $CZ$  using  $q$  as control and  $a$  as target?

## The phase kickback

---

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

3. Let's consider now a controlled  $Z$  operation.
4. What happen if we now apply the  $CZ$  using  $q$  as control and  $a$  as target?
5. we could expect something happen on the target qubit! Not really:

## The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

3. Let's consider now a controlled  $Z$  operation.
4. What happen if we now apply the  $CZ$  using  $q$  as control and  $a$  as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) =$$

## The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

3. Let's consider now a controlled  $Z$  operation.
4. What happen if we now apply the  $CZ$  using  $q$  as control and  $a$  as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} =$$

## The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

3. Let's consider now a controlled  $Z$  operation.
4. What happen if we now apply the  $CZ$  using  $q$  as control and  $a$  as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} = \frac{|01\rangle - |11\rangle}{\sqrt{2}} =$$

## The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

3. Let's consider now a controlled  $Z$  operation.
4. What happen if we now apply the  $CZ$  using  $q$  as control and  $a$  as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} = \frac{|01\rangle - |11\rangle}{\sqrt{2}} = \frac{|0\rangle - |1\rangle}{\sqrt{2}} \otimes |1\rangle.$$

## The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

3. Let's consider now a controlled  $Z$  operation.
4. What happen if we now apply the  $CZ$  using  $q$  as control and  $a$  as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} = \frac{|01\rangle - |11\rangle}{\sqrt{2}} = \frac{|0\rangle - |1\rangle}{\sqrt{2}} \otimes |1\rangle.$$

6. acting on  $a$  we got a phase kickback on the system!

## The phase kickback

One clever trick we can implement using ancillas is the **phase kickback**.

1. Take into account two qubits: a system qubit  $q$  and an ancilla  $a$ .
2. Suppose we prepare  $q$  into a superposed state and  $a$  into the excited state:

$$|q\rangle = H|0\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |a\rangle = |1\rangle.$$

3. Let's consider now a controlled  $Z$  operation.
4. What happens if we now apply the  $CZ$  using  $q$  as control and  $a$  as target?
5. we could expect something happen on the target qubit! Not really:

$$CZ\left(\frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes |1\rangle\right) = CZ\left(\frac{|01\rangle + |11\rangle}{\sqrt{2}}\right) = \frac{|01\rangle + Z|11\rangle}{\sqrt{2}} = \frac{|01\rangle - |11\rangle}{\sqrt{2}} = \frac{|0\rangle - |1\rangle}{\sqrt{2}} \otimes |1\rangle.$$

6. acting on  $a$  we got a phase kickback on the system!

### Important

This happens if the ancilla is prepared into a state which is eigenvector of the controlled operation.

# Is the phase kickback useful?

Some of the most powerful and famous quantum computing algorithms make use of the phase kickback



Figure 1: Deutsch-Josza




Figure 2: Grover




Figure 3: Shor




Figure 4: HHL

All of them are proved to outperform any classical algorithm, leading to a theoretical **quantum advantage**!