



北京大学

本科生毕业论文

题目：镍铁层状双氢氧化物高活性析
氧电催化剂 (流水线演示)

姓 名：演示作者
学 号：2600000000
院 系：示例学院
专 业：材料化学
研究方向：电催化
导 师：示例导师教授

☒ 学术学位 ☐ 专业学位

二〇二六年七月

摘要

DEMO ONLY - generated by the paperflow pipeline with an illustrative synthetic dataset; not real experimental results and not for submission.

The oxygen evolution reaction (OER) limits the efficiency of alkaline water electrolysis, and inexpensive catalysts that rival noble-metal oxides are therefore in continuous demand [2]. In this demonstration study we examine, on an illustrative example dataset, how iron incorporation transforms nickel hydroxide into a high-activity NiFe layered double hydroxide (NiFe-LDH) electrocatalyst. Powder X-ray diffraction confirms a phase-pure hydrotalcite-type structure, while Ni 2p X-ray photoelectron spectroscopy resolves coexisting Ni²⁺ and Ni³⁺ states consistent with Fe-modulated electronic structure [1]. In 1 M KOH, the illustrative polarization data give an overpotential of 273 mV at 10 mA cm⁻² for NiFe-LDH, surpassing both the RuO₂ benchmark (324 mV) and Fe-free Ni(OH)₂ (466 mV), and the accompanying Tafel slope of 38 mV per decade indicates fast electron-transfer kinetics [3]. These trends reproduce the well-established activity ordering of the NiFe (oxy)hydroxide family and illustrate an end-to-end, citation-verified manuscript-generation workflow in which every quantitative claim is traceable either to the declared example dataset or to literature that survived authenticity checking. The workflow, rather than the chemistry, is the deliverable of this document, and all numbers herein are illustrative rather than experimental.

关键词：析氧反应、镍铁层状双氢氧化物、电催化

Iron Incorporation Turns Nickel Hydroxide into a High-Activity Oxygen Evolution Electrocatalyst: An Illustrative NiFe-LDH Case Study

Paperflow Demo (Materials Chemistry)

Directed by Prof. Demo

ABSTRACT

(English abstract omitted in this demo.)

KEYWORDS: OER, NiFe-LDH, electrocatalysis

目录

第一章	Introduction	1
第二章	Results and Discussion	2
第三章	Conclusion	4

第一章 Introduction

Electrochemical water splitting stores renewable electricity as hydrogen, but its overall efficiency is throttled by the sluggish four-electron oxygen evolution reaction at the anode. Benchmarking studies have standardized the comparison of OER catalysts through the overpotential required to reach 10 mA cm^{-2} , a metric tied to the photocurrent of an ideal solar-driven device [2]. Within design-principle frameworks that correlate activity with electronic-structure descriptors, transition-metal oxides and (oxy)hydroxides have emerged as the most promising earth-abundant candidates in alkaline media [4, 5]. Among them, nickel-iron systems occupy a special position: seminal work demonstrated that even incidental iron impurities dramatically enhance the apparent activity of nickel (oxy)hydroxide, identifying Fe incorporation as the true origin of its celebrated performance [1]. Subsequent studies of NiFe layered double hydroxides consolidated this family as the state of the art for non-acidic OER, with overpotentials frequently below 300 mV [3]. The present manuscript revisits this well-charted chemistry for a different purpose. It serves as an end-to-end demonstration of paperflow, a pipeline that drafts, cites, verifies, and assembles a manuscript automatically, using an illustrative NiFe-LDH dataset. Our aim is to show that each stage, from outline to citation-authenticity gating to document assembly, can be orchestrated reproducibly while keeping every factual claim anchored to verified sources.

第二章 Results and Discussion

Figure 1 summarizes the illustrative structural and electrochemical dataset. The diffraction pattern of NiFe-LDH in Figure 1a reproduces the characteristic (003), (006), (012), and (015) reflections of a hydrotalcite-type layered structure and matches the LDH reference card, whereas the Fe-free control exhibits the distinct beta-Ni(OH)₂ pattern; no impurity phases are apparent in either trace. The Ni 2p spectrum in Figure 1b is deconvoluted into Ni²⁺ and Ni³⁺ components together with their shake-up satellites, and the appreciable Ni³⁺ fraction is consistent with the Fe-induced modulation of nickel electronic structure that has been linked mechanistically to enhanced OER activity in mixed Ni-Fe (oxy)hydroxides [1]. Turning to catalysis, the polarization curves in Figure 1c, acquired on the illustrative dataset in 1 M KOH, place the three electrodes in the order NiFe-LDH > RuO₂ > Ni(OH)₂, with overpotentials at 10 mA cm⁻² of 273, 324, and 466 mV, respectively, following the benchmarking convention of the field [2]. The Tafel analysis in Figure 1d yields slopes of 38, 42, and 65 mV per decade for the same ordering, indicating that Fe incorporation accelerates the rate-determining electron-transfer step rather than merely increasing the electrochemically active area. Both the absolute overpotential range and the roughly one-hundred-millivolt advantage over the nickel-only control agree with the consensus picture assembled for the NiFe (oxy)hydroxide family [3], lending face validity to the synthetic dataset even though it is illustrative rather than experimental.




图 2.1 Structural and electrochemical characterization of NiFe-LDH (illustrative synthetic data): (a) XRD patterns with the LDH reference; (b) Ni 2p XPS with peak deconvolution; (c) OER polarization curves with the η_{10} annotation; (d) Tafel analysis.

第三章 Conclusion

This demonstration assembled a complete, citation-verified manuscript around an illustrative NiFe-LDH oxygen-evolution dataset. The narrative reproduces the accepted understanding of the system: a phase-pure layered double hydroxide, mixed Ni²⁺/Ni³⁺ surface states, and an alkaline OER activity ranking of NiFe-LDH ahead of RuO₂ and far ahead of Fe-free Ni(OH)₂, in line with the benchmarking literature and mechanistic studies of iron incorporation [2, 1, 3]. More importantly for the purpose of this document, every reference cited here passed an automated authenticity gate that queries Crossref and Semantic Scholar and drops any candidate whose DOI or title cannot be confirmed; one deliberately fabricated reference planted in the candidate list was rejected by this gate and therefore does not appear in the bibliography. The pipeline thus enforces, by construction, the separation between verifiable literature claims and the clearly labeled illustrative dataset. Future work on the pipeline itself includes journal-specific formatting profiles, richer figure-placement logic, and a reviewer-simulation stage, while any scientific use of the chemistry discussed here would of course require real experimental data in place of the example dataset used throughout this demonstration.

参考文献

- [1] Lena Trotochaud, Samantha L. Young, James K. Ranney, Shannon W. Boettcher. Nickel–Iron Oxyhydroxide Oxygen-Evolution Electrocatalysts: The Role of Intentional and Incidental Iron Incorporation. *Journal of the American Chemical Society*, 2014, 136: 6744–6753. DOI: 10.1021/ja502379c.
- [2] Charles C. L. McCrory, Suho Jung, Jonas C. Peters, Thomas F. Jaramillo. Benchmarking Heterogeneous Electrocatalysts for the Oxygen Evolution Reaction. *Journal of the American Chemical Society*, 2013, 135: 16977–16987. DOI: 10.1021/ja407115p.
- [3] Fabio Dionigi, Peter Strasser. NiFe-Based (Oxy)hydroxide Catalysts for Oxygen Evolution Reaction in Non-Acidic Electrolytes. *Advanced Energy Materials*, 2016, 6. DOI: 10.1002/aenm.201600621.
- [4] Jin Suntivich, Kevin J. May, Hubert A. Gasteiger, John B. Goodenough, Yang Shao-Horn. A Perovskite Oxide Optimized for Oxygen Evolution Catalysis from Molecular Orbital Principles. *Science*, 2011, 334: 1383–1385. DOI: 10.1126/science.1212858.
- [5] Alexis Grimaud, Kevin J. May, Christopher E. Carlton, Yueh-Lin Lee, Marcel Risch, Wesley T. Hong, Jigang Zhou, Yang Shao-Horn. Double perovskites as a family of highly active catalysts for oxygen evolution in alkaline solution. *Nature Communications*, 2013, 4. DOI: 10.1038/ncomms3439.