


推荐系统

Example: Predicting movie ratings

→ User rates movies using one to five stars

Movie	Alice (1)	Bob (2)	Carol (3)	Dave (4)
Love at last	5	5	0	0
Romance forever	5	?	?	0
Cute puppies of love	?	4	0	?
Nonstop car chases	0	0	5	4
Swords vs. karate	0	0	5	?



→ n_u = no. users
 → n_m = no. movies
 → $r(i, j) = 1$ if user j has rated movie i
 → $y^{(i, j)}$ = rating given by user j to movie i (defined only if $r(i, j) = 1$)
 0, ..., 5

$n_u = 4$ $n_m = 5$

Content-based recommender systems

Movie	Alice (1)	Bob (2)	Carol (3)	Dave (4)	x_1 (romance)	x_2 (action)
Love at last 1	5	5	0	0	0.9	0
Romance forever 2	5	?	?	0	1.0	0.01
Cute puppies of love 3	?	4	0	?	0.99	0
Nonstop car chases 4	0	0	5	4	0.1	1.0
Swords vs. karate 5	0	0	5	?	0	0.9

→ For each user j , learn a parameter $\theta^{(j)} \in \mathbb{R}^3$. Predict user j as rating movie i with $(\theta^{(j)})^T x^{(i)}$ stars.

$x^{(3)} = \begin{bmatrix} 1 \\ 0.99 \\ 0 \end{bmatrix} \leftrightarrow \theta^{(1)} = \begin{bmatrix} 0 \\ 5 \\ 0 \end{bmatrix}$ $(\theta^{(1)})^T x^{(3)} = 5 \times 0.99 = 4.95$

Problem formulation

→ $r(i, j) = 1$ if user j has rated movie i (0 otherwise)

→ $y^{(i,j)}$ = rating by user j on movie i (if defined)

→ $\theta^{(j)}$ = parameter vector for user j

→ $x^{(i)}$ = feature vector for movie i

→ For user j , movie i , predicted rating: $(\theta^{(j)})^T x^{(i)}$

$$\theta^{(j)} \in \mathbb{R}^{n+1}$$

→ $m^{(j)}$ = no. of movies rated by user j

To learn $\theta^{(j)}$:

$$\min_{\theta^{(j)}} \frac{1}{2m^{(j)}} \sum_{i:r(i,j)=1} \left((\theta^{(j)})^T x^{(i)} - y^{(i,j)} \right)^2 + \frac{\lambda}{2m^{(j)}} \sum_{k=1}^n (\theta_k^{(j)})^2$$

Optimization objective:

To learn $\theta^{(j)}$ (parameter for user j):

$$\rightarrow \min_{\theta^{(j)}} \frac{1}{2} \sum_{i:r(i,j)=1} \left((\theta^{(j)})^T x^{(i)} - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{k=1}^n (\theta_k^{(j)})^2$$

To learn $\theta^{(1)}, \theta^{(2)}, \dots, \theta^{(n_u)}$:

$$\min_{\theta^{(1)}, \dots, \theta^{(n_u)}} \frac{1}{2} \sum_{j=1}^{n_u} \sum_{i:r(i,j)=1} \left((\theta^{(j)})^T x^{(i)} - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{j=1}^{n_u} \sum_{k=1}^n (\theta_k^{(j)})^2$$

Optimization algorithm:

$$\min_{\theta^{(1)}, \dots, \theta^{(n_u)}} \frac{1}{2} \sum_{j=1}^{n_u} \sum_{i:r(i,j)=1} \left((\theta^{(j)})^T x^{(i)} - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{j=1}^{n_u} \sum_{k=1}^n (\theta_k^{(j)})^2$$

$\mathcal{J}(\theta^{(1)}, \dots, \theta^{(n_u)})$

Gradient descent update:

$$\theta_k^{(j)} := \theta_k^{(j)} - \alpha \sum_{i:r(i,j)=1} ((\theta^{(j)})^T x^{(i)} - y^{(i,j)}) x_k^{(i)} \quad (\text{for } k = 0)$$

$$\theta_k^{(j)} := \theta_k^{(j)} - \alpha \left(\sum_{i:r(i,j)=1} ((\theta^{(j)})^T x^{(i)} - y^{(i,j)}) x_k^{(i)} + \lambda \theta_k^{(j)} \right) \quad (\text{for } k \neq 0)$$

Problem motivation

Movie	Alice (1) $\theta^{(1)}$	Bob (2) $\theta^{(2)}$	Carol (3) $\theta^{(3)}$	Dave (4) $\theta^{(4)}$	x_1 (romance)	x_2 (action)
$x^{(1)}$ Love at last	5	5	0	0	1.0	0.0
Romance forever	5	?	?	0	?	?
Cute puppies of love	?	4	0	?	?	?
Nonstop car chases	0	0	5	4	?	?
Swords vs. karate	0	0	5	?	?	?

$\theta^{(1)} = \begin{bmatrix} 0 \\ 5 \\ 0 \end{bmatrix}, \theta^{(2)} = \begin{bmatrix} 0 \\ 5 \\ 0 \end{bmatrix}, \theta^{(3)} = \begin{bmatrix} 0 \\ 0 \\ 5 \end{bmatrix}, \theta^{(4)} = \begin{bmatrix} 0 \\ 0 \\ 5 \end{bmatrix}$

$x^{(1)}$
 $(\theta^{(1)})^T x^{(1)} \approx 5$
 $(\theta^{(2)})^T x^{(1)} \approx 5$
 $(\theta^{(3)})^T x^{(1)} \approx 0$

Optimization algorithm

Given $\theta^{(1)}, \dots, \theta^{(n_u)}$, to learn $x^{(i)}$:

$$\min_{x^{(i)}} \frac{1}{2} \sum_{j:r(i,j)=1} ((\theta^{(j)})^T x^{(i)} - y^{(i,j)})^2 + \frac{\lambda}{2} \sum_{k=1}^n (x_k^{(i)})^2$$

Given $\theta^{(1)}, \dots, \theta^{(n_u)}$, to learn $x^{(1)}, \dots, x^{(n_m)}$:

$$\min_{x^{(1)}, \dots, x^{(n_m)}} \frac{1}{2} \sum_{i=1}^{n_m} \sum_{j:r(i,j)=1} ((\theta^{(j)})^T x^{(i)} - y^{(i,j)})^2 + \frac{\lambda}{2} \sum_{i=1}^{n_m} \sum_{k=1}^n (x_k^{(i)})^2$$

Collaborative filtering

Given $x^{(1)}, \dots, x^{(n_m)}$ (and movie ratings),
can estimate $\theta^{(1)}, \dots, \theta^{(n_u)}$

Given $\theta^{(1)}, \dots, \theta^{(n_u)}$,
can estimate $x^{(1)}, \dots, x^{(n_m)}$

Guess $\theta \rightarrow x \rightarrow \theta \rightarrow x \rightarrow \theta \rightarrow x \rightarrow \dots$

Collaborative filtering optimization objective

→ Given $x^{(1)}, \dots, x^{(n_m)}$, estimate $\theta^{(1)}, \dots, \theta^{(n_u)}$:

$$\min_{\theta^{(1)}, \dots, \theta^{(n_u)}} \left[\frac{1}{2} \sum_{j=1}^{n_u} \sum_{i:r(i,j)=1} ((\theta^{(j)})^T x^{(i)} - y^{(i,j)})^2 + \frac{\lambda}{2} \sum_{j=1}^{n_u} \sum_{k=1}^n (\theta_k^{(j)})^2 \right]$$

→ Given $\theta^{(1)}, \dots, \theta^{(n_u)}$, estimate $x^{(1)}, \dots, x^{(n_m)}$:

$$\min_{x^{(1)}, \dots, x^{(n_m)}} \left[\frac{1}{2} \sum_{i=1}^{n_m} \sum_{j:r(i,j)=1} ((\theta^{(j)})^T x^{(i)} - y^{(i,j)})^2 + \frac{\lambda}{2} \sum_{i=1}^{n_m} \sum_{k=1}^n (x_k^{(i)})^2 \right]$$

Minimizing $x^{(1)}, \dots, x^{(n_m)}$ and $\theta^{(1)}, \dots, \theta^{(n_u)}$ simultaneously:

$$J(x^{(1)}, \dots, x^{(n_m)}, \theta^{(1)}, \dots, \theta^{(n_u)}) = \frac{1}{2} \sum_{(i,j):r(i,j)=1} ((\theta^{(j)})^T x^{(i)} - y^{(i,j)})^2 + \frac{\lambda}{2} \sum_{i=1}^{n_m} \sum_{k=1}^n (x_k^{(i)})^2 + \frac{\lambda}{2} \sum_{j=1}^{n_u} \sum_{k=1}^n (\theta_k^{(j)})^2$$

$$\min_{\substack{x^{(1)}, \dots, x^{(n_m)} \\ \theta^{(1)}, \dots, \theta^{(n_u)}}} J(x^{(1)}, \dots, x^{(n_m)}, \theta^{(1)}, \dots, \theta^{(n_u)})$$

Andrew Ng

Collaborative filtering algorithm

1. Initialize $x^{(1)}, \dots, x^{(n_m)}, \theta^{(1)}, \dots, \theta^{(n_u)}$ to small random values.
2. Minimize $J(x^{(1)}, \dots, x^{(n_m)}, \theta^{(1)}, \dots, \theta^{(n_u)})$ using gradient descent (or an advanced optimization algorithm). E.g. for every $j = 1, \dots, n_u, i = 1, \dots, n_m$:

$$x_k^{(i)} := x_k^{(i)} - \alpha \left(\sum_{j:r(i,j)=1} ((\theta^{(j)})^T x^{(i)} - y^{(i,j)}) \theta_k^{(j)} + \lambda x_k^{(i)} \right)$$

$$\theta_k^{(j)} := \theta_k^{(j)} - \alpha \left(\sum_{i:r(i,j)=1} ((\theta^{(j)})^T x^{(i)} - y^{(i,j)}) x_k^{(i)} + \lambda \theta_k^{(j)} \right)$$

3. For a user with parameters θ and a movie with (learned) features x , predict a star rating of $\theta^T x$.

向量化

Collaborative filtering

$X \Theta^T \leftarrow (\Theta^{(j)})^T (x^{(i)})$

$Y = \begin{bmatrix} 5 & 5 & 0 & 0 \\ 5 & ? & ? & 0 \\ ? & 4 & 0 & ? \\ 0 & 0 & 5 & 4 \\ 0 & 0 & 5 & 0 \end{bmatrix}$

Predicted ratings:

$$\begin{bmatrix} (\theta^{(1)})^T (x^{(1)}) & (\theta^{(2)})^T (x^{(1)}) & \dots & (\theta^{(n_u)})^T (x^{(1)}) \\ (\theta^{(1)})^T (x^{(2)}) & (\theta^{(2)})^T (x^{(2)}) & \dots & (\theta^{(n_u)})^T (x^{(2)}) \\ \vdots & \vdots & \ddots & \vdots \\ (\theta^{(1)})^T (x^{(n_m)}) & (\theta^{(2)})^T (x^{(n_m)}) & \dots & (\theta^{(n_u)})^T (x^{(n_m)}) \end{bmatrix}$$

$\rightarrow X = \begin{bmatrix} -(x^{(1)})^T \\ -(x^{(2)})^T \\ \vdots \\ -(x^{(n_m)})^T \end{bmatrix}$

$\rightarrow \Theta = \begin{bmatrix} -(\theta^{(1)})^T \\ -(\theta^{(2)})^T \\ \vdots \\ -(\theta^{(n_u)})^T \end{bmatrix}$

推荐相关相似电影

Finding related movies

For each product i , we learn a feature vector $x^{(i)} \in \mathbb{R}^n$.

$\rightarrow x_1 = \text{romance}, x_2 = \text{action}, x_3 = \text{comedy}, x_4 = \dots$

How to find movies j related to movie i ?

small $\|x^{(i)} - x^{(j)}\| \rightarrow$ movie j and i are "similar"

5 most similar movies to movie i :

Find the 5 movies j with the smallest $\|x^{(i)} - x^{(j)}\|$.

均值归一化 让 算法运行的更好

Mean Normalization:

$$Y = \begin{bmatrix} 5 & 5 & 0 & 0 & ? \\ 5 & ? & ? & 0 & ? \\ ? & 4 & 0 & ? & ? \\ 0 & 0 & 5 & 4 & ? \\ 0 & 0 & 5 & 0 & ? \end{bmatrix} \quad \mu = \begin{bmatrix} 2.5 \\ 2.5 \\ 2 \\ 2.25 \\ 1.25 \end{bmatrix} \rightarrow \underline{Y} = \begin{bmatrix} 2.5 & 2.5 & -2.5 & -2.5 & ? \\ 2.5 & ? & ? & -2.5 & ? \\ ? & 2 & -2 & ? & ? \\ -2.25 & -2.25 & 2.75 & 1.75 & ? \\ -1.25 & -1.25 & 3.75 & -1.25 & ? \end{bmatrix}$$

For user j , on movie i predict:

$$\rightarrow (\theta^{(j)})^T (x^{(i)}) + \mu_i$$

learn $\underline{\theta^{(j)}}, \underline{x^{(i)}}$

User 5 (Eve):

$$\theta^{(5)} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$(\theta^{(5)})^T (x^{(i)}) + \mu_i$$