


The three bars AB, BC, and CD are each pinned at their ends as shown. Bar AB has length 2 meters; bar BC has length 4 meters, and bar CD has length 3 meters. Bar AB has an angular velocity of $\omega_{AB} = +4 \text{ s}^{-1}$, and an angular acceleration of $\alpha_{AB} = -3 \text{ s}^{-2}$.

- Find the magnitude of acceleration, $|a_B|$, of point B.
- Find the angular velocity, ω_{BC} , of bar BC.
- Find the horizontal component of acceleration of point C.



(From April '98 exam)

$$(a) \quad \vec{a}_B = \vec{a}_A + (\vec{a}_{B/A})_n + (\vec{a}_{B/A})_t$$


$$a_n = \omega^2 r_{AB} = v_B^2 / r_{AB} = \frac{8^2}{2} = 32 \text{ m/s}^2.$$

$$a_t = \alpha r_{AB} = (3)(2) = 6 \text{ m/s}^2$$

But a_n & a_t are perpendicular to each other

$$\begin{aligned} \Rightarrow |\vec{a}| &= \sqrt{a_n^2 + a_t^2} \\ &= \sqrt{32^2 + 6^2} = 32.56 \text{ m/s}^2 \end{aligned}$$


(b) center of zero velocity of BC is at "Z"



$$\tan 30^\circ = \frac{l}{4} \Rightarrow l = 2.31 \text{ meters}$$

$$V_B = \omega_{BC} l \Rightarrow \omega_{BC} = \frac{V_B}{l} = \frac{8}{2.31} = 3.464 \text{ s}^{-1}$$

or draw a vector triangle: $\vec{V}_C = \vec{V}_B + \vec{V}_{C/B}$



(c)

$$V_{C/B} = \omega_{BC} r_{BC} = (3.464)(4) = 13.856$$

$$\& V_B = 8 \text{ m/s.}$$

By Pythagoras,

$$V_C = \sqrt{13.856^2 + 8^2} = 16 \text{ m/s.}$$

$$\Rightarrow (a_c)_{\text{horizontal}} = \frac{V_C^2}{r_{CD}} = \frac{16^2}{3} = 85.3 \text{ m/s}^2 \uparrow$$