

Axi-Higgs
Cosmology

Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook

Axi-Higgs Cosmology

Yu-Cheng QIU

邱渔骋

Collaborators: Leo WH Fung, Lingfeng Li, Hoang Nhan Luu, Tao Liu, S.-H. Henry Tye

Tsung-Dao Lee Institute, Shanghai Jiao Tong University

May 25, 2023

2102.11257 & 2105.01631

I

MOTIVATION

- ▶ Hubble tension: $H_{0,P18} = 67.36 \pm 0.54$ km/s/Mpc [Aghanim et al., 2022] vs. $H_{0,\text{late}} = 73.3 \pm 0.8$ km/s/Mpc [Verde et al., 2019] from $z \lesssim 2$.
 $H_{0,\text{late}} = 66.6^{+4.4}_{-3.3}$ km/s/Mpc from Refsdal's reappearance. (2305.06367)
- ▶ ${}^7\text{Li}$ Problem in BBN: the abundance ratio ${}^7\text{Li}/\text{H} \times 10^{10} : 1.6 \pm 0.3$ (observed) vs. 5.6 ± 0.3 (theoretical) [Zyla et al. 2020, Pitrou et al., 2018, Iliadis and Coc, 2020].

- ▶ **Hubble tension:** $H_{0,P18} = 67.36 \pm 0.54$ km/s/Mpc [Aghanim et al., 2022] vs. $H_{0,\text{late}} = 73.3 \pm 0.8$ km/s/Mpc [Verde et al., 2019] from $z \lesssim 2$.
 $H_{0,\text{late}} = 66.6^{+4.4}_{-3.3}$ km/s/Mpc from Refsdal's reappearance. (2305.06367)
- ▶ **${}^7\text{Li}$ Problem** in BBN: the abundance ratio ${}^7\text{Li}/\text{H} \times 10^{10} : 1.6 \pm 0.3$ (observed) vs. 5.6 ± 0.3 (theoretical) [Zyla et al. 2020, Pitrou et al., 2018, Iliadis and Coc, 2020].
- ▶ The weak lensing measurement of S_8 together with the clustering parameter σ_8 [Troxel et al., 2018] yields a value smaller $S_{8,\text{DES}} = 0.773^{+0.026}_{-0.020}$ than given by the CMB- Λ CDM value, $S_{8,\text{CMB}} = 0.832 \pm 0.013$.
- ▶ Isotropic cosmic birefringence angle based on the cross-power (parity-violating) C_l^{EB} data in CMB [Minami and Komatsu, 2020], deviate from 0 by $\sim 2.4\sigma$.
Later improved result: deviate from 0 by $\sim 3.6\sigma$ [Eskilt and Komatsu, 2022].

Tensions in Λ CDM





Figure: Valentino et al. 2021

Figure: Zyla et al. 2020




Figure: Handley and Lemos 2019




Figure: Minami and Komatsu 2020



II

SOLUTIONS

A Higher Higgs VEV during the BBN epoch.

- ▶ Fermi constant $G_F \propto v^{-2}$. A smaller G_F leads to an earlier freeze out of the $n \rightleftharpoons p$ and a longer n lifetime.

A larger n density than that in the standard BBN.

- ▶ Electro mass $m_e \propto v$. A larger m_e will reduce the rate of $n \rightleftharpoons p$ and delay neutron decay.

A Higher Higgs VEV during the BBN epoch.

- ▶ Fermi constant $G_F \propto v^{-2}$. A smaller G_F leads to an earlier freeze out of the $n \rightleftharpoons p$ and a longer n lifetime.

A larger n density than that in the standard BBN.

- ▶ Electro mass $m_e \propto v$. A larger m_e will reduce the rate of $n \rightleftharpoons p$ and delay neutron decay.
- ▶ Mass difference $\Delta m_q = m_d - m_u \propto v$, which contributes to Δm_{np} and impact $n \rightleftharpoons p$ and neutron decay oppositely relative to G_F and m_e .
- ▶ Averaged light quark mass, which contributes to pion mass m_π . A larger pion mass makes nuclei less tightly bound. The nuclear-reaction rates thus may change substantially.

An upward variance of ν will

- ▶ reduce the primordial mass fraction of Helium-4, Y_p ,
- ▶ raise the Deuterium primordial abundance D/H relative to Hydrogen.

An upward variance of ν will

- ▶ reduce the primordial mass fraction of Helium-4, Y_p ,
- ▶ raise the Deuterium primordial abundance D/H relative to Hydrogen.

The current experimental bounds on Y_p and D/H are still compatible for $\delta\nu_{\text{BBN}} \sim \mathcal{O}(1\%)$ and $\delta\eta \sim \mathcal{O}(1\%)$ (**CMB baryon-to-photon ratio**).

Following this, to addressing the ${}^7\text{Li}$ problem, one needs [[Pitrou et al. 2018](#)]

$$\delta\nu_{\text{BBN}} = (1.1 \pm 0.1)\% , \quad \delta\eta = (1.7 \pm 1.3)\% .$$

Focus on the model $\Lambda\text{CDM}+m_e$. [Hart and Chluba, 2019]

$$\delta m_e \approx \delta v_{\text{rec}} \sim 1\%.$$

- ▶ Thompson scattering cross-section, $\sigma_T \propto m_e^{-2}$.
- ▶ The atomic energy levels, $E_i \propto m_e$.
- ▶ ...

Focus on the model $\Lambda\text{CDM}+m_e$. [Hart and Chluba, 2019]

$$\delta m_e \approx \delta v_{\text{rec}} \sim 1\%.$$

- ▶ Thompson scattering cross-section, $\sigma_T \propto m_e^{-2}$.
- ▶ The atomic energy levels, $E_i \propto m_e$.
- ▶ ...

\implies Shift up the redshift of the rec z_* , and the baryon drag redshift z_d .

Focus on the model $\Lambda\text{CDM}+m_e$. [Hart and Chluba, 2019]

$$\delta m_e \approx \delta v_{\text{rec}} \sim 1\%.$$

- ▶ Thompson scattering cross-section, $\sigma_T \propto m_e^{-2}$.
- ▶ The atomic energy levels, $E_i \propto m_e$.
- ▶ ...

\Rightarrow Shift up the redshift of the rec z_* , and the baryon drag redshift z_d .

\Rightarrow Sound horizon at rec decrease.

Focus on the model $\Lambda\text{CDM}+m_e$. [Hart and Chluba, 2019]

$$\delta m_e \approx \delta v_{\text{rec}} \sim 1\%.$$

- ▶ Thompson scattering cross-section, $\sigma_T \propto m_e^{-2}$.
- ▶ The atomic energy levels, $E_i \propto m_e$.
- ▶ ...

\Rightarrow Shift up the redshift of the rec z_* , and the baryon drag redshift z_d .

\Rightarrow Sound horizon at rec decrease.

\Rightarrow To keep angular sound horizon at rec unchanged, H_0 increases.




Figure: BM₁: $\delta v = 1.1\%$. BM₂: $\delta v = 1.0\%$. BM₃: $\delta v = 0\%$. BM_{BBN}: $\delta v = \delta v_{\text{BBN}}$.

S_8/σ_8 'Tension'

Axi-Higgs
Cosmology

Yu-Cheng QIU


Motivation

Solutions

A(-xion) Solution

Summary and
Outlook

Introduce the ultra-light axion-like particle, ω_a .



Isotropic Cosmic Birefringence

Axi-Higgs
Cosmology

Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook

Consider the axion-photon coupling $aF\tilde{F}$. [Carroll and Field, 1990]

$$\mathcal{L} \supset \frac{1}{32\pi^2} \frac{a}{f_a} F_{\mu\nu} \tilde{F}^{\mu\nu}$$

$$\Rightarrow \beta_{\text{obs}} \sim \frac{1}{16\pi^2} \frac{a_{\text{ini}}}{f_a}$$

$$\Rightarrow \frac{a_{\text{ini}}}{f_a} \approx 1.0 \pm 0.3$$




Figure: T. Fujita et. al., 2020




Figure: Credit: Y. Minami.



III

A(-XION) SOLUTION

- ▶ An up-shift of electron mass at the recombination era $\delta m_e \sim 1\%$ could resolve Hubble tension. [Hart & Chluba (2020)]
- ▶ ^7Li problem could be solved by lifting the Higgs VEV $\delta v \sim 1\%$. [2102.11257 and refs therein]

- ▶ An up-shift of electron mass at the recombination era $\delta m_e \sim 1\%$ could resolve Hubble tension. [Hart & Chluba (2020)]
- ▶ ^7Li problem could be solved by lifting the Higgs VEV $\delta v \sim 1\%$. [2102.11257 and refs therein]
- ▶ Ultra light axion could be used to explain ICB. [Minami & Komatsu (2020)]
- ▶ Introducing the ultra-light axion may be helpful suppressing the S_8/σ_8 . [KiDS-450 (2017), Handley & Lemos (2019)]

- ▶ An up-shift of electron mass at the recombination era $\delta m_e \sim 1\%$ could resolve Hubble tension. [Hart & Chluba (2020)]
- ▶ ^7Li problem could be solved by lifting the Higgs VEV $\delta v \sim 1\%$. [2102.11257 and refs therein]
- ▶ Ultra light axion could be used to explain ICB. [Minami & Komatsu (2020)]
- ▶ Introducing the ultra-light axion may be helpful suppressing the S_8/σ_8 . [KiDS-450 (2017), Handley & Lemos (2019)]

A model with evolving Higgs VEV and the axion?

- ▶ An up-shift of electron mass at the recombination era $\delta m_e \sim 1\%$ could resolve Hubble tension. [Hart & Chluba (2020)]
- ▶ ^7Li problem could be solved by lifting the Higgs VEV $\delta v \sim 1\%$. [2102.11257 and refs therein]
- ▶ Ultra light axion could be used to explain ICB. [Minami & Komatsu (2020)]
- ▶ Introducing the ultra-light axion may be helpful suppressing the S_8/σ_8 . [KiDS-450 (2017), Handley & Lemos (2019)]

A model with evolving Higgs VEV and the axion?

Axi-Higgs is constructed by introducing coupling between ultra-light axion(s) and the Higgs.



Axi-Higgs Potential

Axi-Higgs
Cosmology

Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook

Scalar potential is $V = V_a + V_\phi$, where

$$V_a = m_a^2 f_a^2 \left(1 - \cos \frac{a}{f_a} \right) \approx \frac{1}{2} m_a^2 a^2 - \frac{1}{24} \frac{m_a^2}{f_a^2} a^4 + \dots ,$$

$$V_\phi = \left| m_s^2 F(a) - \kappa \phi^\dagger \phi \right|^2 , \quad F(a) = 1 + \frac{C a^2}{M_{\text{Pl}}^2} .$$

Scalar potential is $V = V_a + V_\phi$, where

$$V_a = m_a^2 f_a^2 \left(1 - \cos \frac{a}{f_a} \right) \approx \frac{1}{2} m_a^2 a^2 - \frac{1}{24} \frac{m_a^2}{f_a^2} a^4 + \dots ,$$

$$V_\phi = \left| m_s^2 F(a) - \kappa \phi^\dagger \phi \right|^2 , \quad F(a) = 1 + \frac{C a^2}{M_{\text{Pl}}^2} .$$

Neglect three Goldstone directions and let $\phi^\dagger \phi \rightarrow v^2/2$, then

$$V \approx \frac{1}{2} m_a^2 a^2 + |B(a, v)|^2 , \quad B = m_s^2 \left(1 + \frac{C a^2}{M_{\text{Pl}}^2} \right) - \frac{1}{2} \kappa v^2 .$$

Treating $a(t)$ as a background field, the Higgs VEV is given by minimize $\langle V_\phi \rangle$, which is

$$\langle \phi^\dagger \phi \rangle \equiv \frac{v^2}{2} = \frac{m_s^2}{\kappa} F(a)$$

Evolving Higgs VEV

Axi-Higgs
Cosmology

Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook

▶ $m_a \sim 10^{-29} \text{ eV} \implies a(t)$ evolves in the cosmic time scale.

▶ Higgs VEV $v(a(t))$ also evolves in the cosmic time scale,

$$\delta v(t) = \frac{v(t) - v_0}{v_0} = [F(a(t))]^{1/2} - 1 \simeq \frac{Ca(t)^2}{2M_{\text{Pl}}^2},$$

where $v_0 = \sqrt{2}m_s/\sqrt{\kappa} = 246 \text{ GeV}$.

▶ $a(t)$ is determined by KG equation in the FLRW background,

$$\ddot{a} + 3H(t)\dot{a} + \frac{\partial V_a}{\partial a} = 0.$$

Evolving Higgs VEV is driven by misalignment mechanism of the ultra-light axion.

One may worry about the back reaction from the Higgs to the axion.

One may worry about the back reaction from the Higgs to the axion.
Consider the coupled EoMs for $a(t)$ and $v(t)$,

$$\ddot{a} + 3H\dot{a} + \left[m_a^2 + \frac{4Cm_s^2}{M_{\text{Pl}}^2} B(a, v) \right] a \approx 0$$
$$\ddot{v} + (3H + \Gamma_\phi) \dot{v} - 2\kappa B(a, v)v = 0 ,$$

where $\Gamma_\phi \sim 4 \text{ MeV}$ is effective Higgs field dissipation.

One may worry about the back reaction from the Higgs to the axion.

Consider the coupled EoMs for $a(t)$ and $v(t)$,

$$\ddot{a} + 3H\dot{a} + \left[m_a^2 + \frac{4Cm_s^2}{M_{\text{Pl}}^2} B(a, v) \right] a \approx 0$$
$$\ddot{v} + (3H + \Gamma_\phi) \dot{v} - 2\kappa B(a, v)v = 0 ,$$

where $\Gamma_\phi \sim 4 \text{ MeV}$ is effective Higgs field dissipation.

- ▶ At first sight, $m_s^2 B / M_{\text{Pl}}^2 \sim 10^{50} m_a^2$ and second term in potential driven force dominated over m_a . Fortunately, this is not the case.
- ▶ Thanks to the presence of Γ_ϕ , Higgs field profile got damped quickly to the value where $B(a, v) = 0$.

The coupled system will evolve along the valley in a - ϕ configuration space.

Axi-Higgs
Cosmology


Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook





Evolving Higgs VEV

Axi-Higgs
Cosmology

Yu-Cheng QIU

Motivation


Solutions

A(-xion) Solution

Summary and
Outlook

$$\begin{aligned}\delta v(t) &= \frac{Ca(t)^2}{2M_{\text{Pl}}^2} \\ &= \delta v_{\text{ini}} \frac{a(t)^2}{a_{\text{ini}}^2}\end{aligned}$$

Set $\delta v_{\text{ini}} = 1\%$.



- ▶ One has two bounds on the axion mass.
 1. Keep $\delta v \geq 1\%$ until $t \gtrsim t_{\text{rec}}$.
 2. $d(\delta v)/dt \lesssim 10^{-16} \text{ yr}^{-1}$ required by experimental bound.

This leads to

$$1.0 \lesssim \frac{m_a}{10^{-29} \text{ eV}} \lesssim 3.3, \quad 68\% \text{ C.L.}$$

- ▶ ICB determines $a_{\text{ini}}/f_a \approx 1$.
- ▶ Suppose a takes up x fraction of today's matter energy density and one could constrain a_{ini} .

Numerically, parameters are

$$a_{\text{ini}} \approx 3.7 \times 10^{17} \text{ GeV} \left(\frac{x}{0.01} \right)^{1/2} \left(\frac{\xi}{1.5} \right)^{-1},$$

$$f_a \approx 3.8 \times 10^{17} \text{ GeV} \left(\frac{x}{0.01} \right)^{1/2} \left(\frac{\xi}{1.5} \right)^{-1},$$

$$C \approx 0.84 \left(\frac{\delta v_{\text{ini}}}{0.01} \right) \left(\frac{x}{0.01} \right)^{-1} \left(\frac{\xi}{1.5} \right)^2,$$

where $x = \omega_a / (\omega_a + \omega_b + \omega_c)$,

and ξ is the numerical factor from equation $\xi H(z_a) = m_a$.

Be aware that m_a does not come in above parameters.

There are four parameters in the single axion axi-Higgs model,

$$m_a, \delta v_{\text{ini}}, a_{\text{ini}}, f_a$$

which are all relatively well-constrained.

- ▶ $\delta v_{\text{BBN}} \approx \delta v_{\text{rec}} > \delta v_0 = 0 \implies m_a \approx 10^{-30} \text{ eV} - 10^{-29} \text{ eV}$
- ▶ ${}^7\text{Li}$ and H_0 puzzle $\implies \delta v_{\text{ini}} \approx 1\%$
- ▶ H_0 tension & S_8/σ_8 tension $\implies a_{\text{ini}} \approx 10^{17} \text{ GeV}$
- ▶ CMB Birefringence $\implies f_a \approx a_{\text{ini}} \approx 10^{17} \text{ GeV}$

Axi-Higgs Parameters

Axi-Higgs
Cosmology


Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook



Two-axion Axi-Higgs

Axi-Higgs
Cosmology

Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook

- ▶ As the evolution of single-axion, one would expect that $\delta v_{\text{BBN}} > \delta v_{\text{rec}}$.
- ▶ To better resolve Hubble tension, one would require $\delta v_{\text{rec}} > 1\%$.
Meanwhile, BBN analysis prefers $\delta v_{\text{BBN}} < 1.1\%$.

Two-axion Axi-Higgs

Axi-Higgs
Cosmology

Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook

- ▶ As the evolution of single-axion, one would expect that $\delta v_{\text{BBN}} > \delta v_{\text{rec}}$.
- ▶ To better resolve Hubble tension, one would require $\delta v_{\text{rec}} > 1\%$. Meanwhile, BBN analysis prefers $\delta v_{\text{BBN}} < 1.1\%$.
- ▶ In fuzzy dark matter scenario, an axion with mass $\sim 10^{-22}$ eV as the cold dark matter can resolve a number of problems in the weakly interacting massive particle (WIMP) model.
- ▶ A second axion could naturally appear in function F that actually responsible for Higgs VEV.

Two-axion Axi-Higgs

Axi-Higgs
Cosmology

Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook

- ▶ As the evolution of single-axion, one would expect that $\delta v_{\text{BBN}} > \delta v_{\text{rec}}$.
- ▶ To better resolve Hubble tension, one would require $\delta v_{\text{rec}} > 1\%$. Meanwhile, BBN analysis prefers $\delta v_{\text{BBN}} < 1.1\%$.
- ▶ In fuzzy dark matter scenario, an axion with mass $\sim 10^{-22}$ eV as the cold dark matter can resolve a number of problems in the weakly interacting massive particle (WIMP) model.
- ▶ A second axion could naturally appear in function F that actually responsible for Higgs VEV.

Consider two axions with mass $m_1 = 10^{-29}$ eV and $m_2 = 10^{-22}$ eV. Neglecting interaction between a_1 and a_2 , function F is given by

$$F(a_1, a_2) = 1 + \frac{C_1 a_1^2}{M_{\text{Pl}}^2} + \frac{C_2 a_2^2}{M_{\text{Pl}}^2}.$$

We expect small deviation of Higgs VEV, which could be approximated by

$$\delta v(t) = F^{1/2} - 1 \approx \frac{C_1 a_1^2}{2M_{\text{Pl}}^2} + \frac{C_2 a_2^2}{2M_{\text{Pl}}^2}.$$

Two-axion Axi-Higgs

Axi-Higgs
Cosmology

Yu-Cheng QIU

Motivation

Solutions

A(-xion) Solution

Summary and
Outlook

- ▶ $a_{1,\text{ini}}$ the same as single-axion case. Consider $a_{2,\text{ini}}$ consists of most of the dark matter and it starts to roll down at $z_2 \sim 2 \times 10^6$, one has

$$a_{1,\text{ini}} \approx 3.7 \times 10^{17} \text{ GeV} , \quad a_{2,\text{ini}} \approx 1.5 \times 10^{17} \text{ GeV} .$$

- ▶ $C_{1,2}$ are determined by

$$\frac{C_1 a_{1,\text{ini}}^2}{2M_{\text{Pl}}^2} + \frac{C_2 a_{2,\text{ini}}^2}{2M_{\text{Pl}}^2} = \delta v_{\text{BBN}} , \quad \frac{C_1 a_1^2(t_{\text{rec}})}{2M_{\text{Pl}}^2} = \delta v_{\text{rec}} .$$

For $\delta v_{\text{BBN}} = 1\%$, $\delta v_{\text{rec}} = 2\%$ and relation $a_1(t_{\text{rec}}) \approx 0.99 a_{1,\text{ini}}$,

$$C_1 \approx 1.7 , \quad C_2 \approx -5.1 .$$




Figure: Drift rate of ν .




Figure: $\delta\nu$ from Quasar Spectrum.



IV

SUMMARY

- ▶ Axi-Higgs model introduces an evolving Higgs VEV using $|\cdots|^2$.
- ▶ This form $|\cdots|^2$ ensures the cosmic evolution of v and protects the axion evolution from the Higgs back-reaction.

- ▶ Axi-Higgs model introduces an evolving Higgs VEV using $|\dots|^2$.
- ▶ This form $|\dots|^2$ ensures the cosmic evolution of v and protects the axion evolution from the Higgs back-reaction.
- ▶ A slightly higher v in the early universe helps resolve Hubble tension and ${}^7\text{Li}$ problem.
- ▶ The introduction of axion helps explain the ICB and suppress S_8/σ_8 .

- ▶ Axi-Higgs model introduces an evolving Higgs VEV using $|\dots|^2$.
- ▶ This form $|\dots|^2$ ensures the cosmic evolution of v and protects the axion evolution from the Higgs back-reaction.
- ▶ A slightly higher v in the early universe helps resolve Hubble tension and ${}^7\text{Li}$ problem.
- ▶ The introduction of axion helps explain the ICB and suppress S_8/σ_8 .
- ▶ The parameters are tightly constrained and could be tested in the near future.

Thank you for your attention.

- Consider $X^2 = 0$ with $X = x + \sqrt{2}\theta G + \theta^2 F^X$.

$$x^2, \quad xG_\alpha = 0, \quad 2xF^X - GG = 0 \quad \implies \quad X = \frac{GG}{2F^X} + \sqrt{2}\theta G + \theta^2 F^X.$$

- For $X^2 = XQ = 0$ with $Q = q + \sqrt{2}\theta\phi + \theta^2 F^Q$,

$$x = \frac{GG}{2F^X}, \quad xq = 0, \quad qG_\alpha + x\psi_\alpha = 0, \quad qF^X + xF^Q - G\psi = 0$$

$$\implies \quad q = \frac{1}{F^X} \left(\psi - \frac{F^Q G}{2F^X} \right) G$$

- $\bar{D}_{\dot{\alpha}}(X\bar{Z}) = X^2 = 0$ with $Z = z + \sqrt{2}\theta\chi + \theta^2 F^Z$ gives

$$\chi = i\partial_\mu z \sigma^\mu \frac{\bar{G}}{\bar{F}^X}$$

$$F^Z = -\partial_\mu \left(\frac{\bar{G}}{\bar{F}^X} \right) \bar{\sigma}^\nu \sigma^\mu \frac{\bar{G}}{\bar{F}^X} \partial_\nu z + \frac{\bar{G}\bar{G}}{2(\bar{F}^X)^2} \partial^2 z$$

The axi-Higgs mass matrix is given by

$$\mathbf{M} = \begin{pmatrix} m_a^2 + \frac{8m_s^4 a^2}{M_{\text{Pl}}^4} & -\frac{2\sqrt{2}m_s^2 a v}{M_{\text{Pl}}^2} \\ -\frac{2\sqrt{2}m_s^2 a v}{M_{\text{Pl}}^2} & v^2 \end{pmatrix}, \quad \lim_{m_a \rightarrow 0} \det \mathbf{M} = 0$$

Diagonalize \mathbf{M} , one has

$$\left(m_\phi^{\text{phys}}\right)^2 \approx 4m_s^2 \left(1 + \frac{a^2}{M_{\text{Pl}}^2}\right) + \mathcal{O}(m_a^2)$$

$$\left(m_a^{\text{phys}}\right)^2 \approx m_a^2 + \mathcal{O}(m_a^4)$$

- Above the scale $\sqrt{m_a f_a}$, the shift symmetry of axion is restored.

$$\Delta m_a^2 \sim \frac{1}{\pi^2} \left(\frac{m_s^2}{M_{\text{Pl}}^2}\right) \left(\frac{m_a^2 f_a^2}{m_\phi^2}\right) \lesssim m_a^2$$

Spontaneous SUSY-breaking by $\overline{D3}$ -brane could be described by a nonlinear supergravity model. [Kallosh & Wrase (2014)]

$$K = X^\dagger X + \dots, \quad W = MX + \dots, \quad \underbrace{X^2 = 0}_{\text{nilpotent condition}}$$

The nilpotent condition projects out the scalar part of X ,

$$X = \frac{GG}{2F^X} + \sqrt{2}\theta G + \theta^2 F^X.$$

The X contributes to the scalar potential as

$$V_X = |M|^2$$

Nilpotent superfield X could be used as projector to eliminate d.o.f.s in other superfields. [Lindstorm & Rocek (1979), Komargodski & Seiberg (2009), Dall'Agata & Farakos (2016)]

- ▶ $XQ = 0$ projects out scalar part of Q
- ▶ $XQ = \text{chiral}$ projects out fermionic d.o.f. of Q .
- ▶ ...

Higgs d.o.f are projected out to properly explain the cosmological constant problem. [Li, Qiu and Tye 2010.10089]

$$\begin{aligned} H_u H_d &\rightarrow \phi^\dagger \phi \\ \langle V_\mu \rangle + \langle V_D \rangle &\rightarrow 0 \end{aligned}$$

The axi-Higgs coupling

Axi-Higgs
Cosmology

Yu-Cheng QIU

Recall superpotential $W \supset X (\tilde{m}_s^2 + \tilde{\gamma} H_u H_d)$, where parameter \tilde{m}_s and $\tilde{\gamma}$ is in principle determined by geometric sector (U_i, S) , which intrinsically include axion-like fields. Thus, it is natural to introduce

$$V_X \rightarrow \left| m_s^2 G(a) - \kappa K(a) \phi^\dagger \phi \right|^2 = \left| K(a) \left[m_s^2 F(a) - \kappa \phi^\dagger \phi \right] \right|^2,$$

where $G(a=0) = K(a=0) = 1$,

$$G(a) = 1 + \frac{ga^2}{M_{\text{Pl}}^2}, \quad K(a) = 1 + \frac{ka^2}{M_{\text{Pl}}^2}, \quad F(a) = \frac{G(a)}{K(a)} \approx 1 + \frac{Ca^2}{M_{\text{Pl}}^2},$$

and $C = g - k$ is a constant whose positivity is undetermined.

$K(a)$ is not important. Let $K(a) = 1$.