

# Idiopathic pulmonary fibrosis: a study using volumetric imaging and functional data in a computational lung model




Yuwen Zhang, Alys Clark, Haribalan Kumar, Merryn Tawhai, Auckland Bioengineering Institute, The University of Auckland, New Zealand  
David Milne, Margaret Wilsher, Auckland City Hospital, Auckland, New Zealand  
Brian Bartholmai, Department of Radiology, Mayo Clinic, Minnesota, US

## Introduction

Idiopathic pulmonary fibrosis (IPF) is an aggressive idiopathic interstitial pneumonia, and often occurs in elderly adults. In IPF, fibrosis typically develops preferentially in posterior-basal lung regions, and often co-exists with emphysema. Currently it is not clear how - or whether - the spatial distribution of tissue abnormalities in IPF (including classifications of tissue type) correlate with pulmonary function tests (PFTs) and their change over time. This work aims to develop a new quantitative tool that integrates data from volumetric imaging, PFTs, and computational models for lung function, to understand differences between IPF and normal older lungs.

## Tissue classification and quantification





- Data**- Clinical IPF CT data from Auckland City Hospital 8 patients, 4 of them contain more than one time point scan for each
- Pulmonary parenchyma classification** - CALIPER (Computer-Aided Lung Information for Pathology Evaluation and Ratings) classification of abnormalities based on signature mapping techniques



3D grid density visualization of tissue patterns

- Fibrosis has a consistently higher tissue density (0.34/0.41 for reticular/ground-glass) compared to normal tissue (0.28) over time. In contrast, emphysema has lower density (0.08)
- Fibrosis presents predominantly in lower lobes (72%, 58%, 65% for honeycomb, reticular, ground-glass). But emphysema appears predominantly in upper lobes (73%).
- The distribution of fibrosis is basal, peripheral and pathy. The disease mostly occurs in small airways which causes peribronchiolar airway constriction.

## Analysis 1




- Mode 1 of the SSM is significantly different between IPF and normal subjects and correlates with percent of fibrosis ( $p < 0.01$ ).
- The most significant variation in shape (mode 1 of the SSM) relates to the anteroposterior diameter of the lung, and the ratio of apical and basal diameters
- There is a significant difference of right lower lobe volume and right middle lobe volume between normal old and IPF lungs ( $p < 0.001$ ,  $p < 0.001$  respectively).




## Analysis 2

### Patient-specific computational modelling of lung function


#### Step1: Airway and vessel tree generation with disease labeling




#### Step 2: Ventilation and perfusion simulation



- Patient-specific perfusion
- Reduce vessel radius of fibrosis labelled regions



#### Step 3: V/Q distribution (gas transfer modeling)



Old normal PaO2 89.02mm Hg  
CT based PaO2 is 78.11mm Hg  
Model base PaO2 is 72.91mm Hg

## Summary

- We classified the pulmonary parenchyma representing IPF features and performed quantitative analysis of IPF lungs.
- Statistical shape analysis suggests quantifiable differences from normal in lung shape are present in IPF.
- V/Q mismatch (impaired gas exchange) is present in 'normal' tissue as well as regions that are classified as abnormal.

**Acknowledgements** – Clinical data for this study was provided by Drs ML Wilsher and DG Milne, Auckland City Hospital; CALIPER analysis was conducted by Dr B Bartholmai, The Mayo Clinic.