

# Homework 10 of Introduction to Analysis(II)

AM15 黃琦翔 111652028

April 23, 2024

1. (a)  $f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0} 1 + 2h \sin\left(\frac{1}{h}\right) = 1.$
- (b)  $\lim_{x \rightarrow 0} f(x) = \lim_{h \rightarrow \infty} \frac{1}{h} + 2 \frac{\sin(h)}{h^2} = 0.$  Thus,  $f$  is continuous on 0. And for any  $x$  close to 0,  $|f(x+h) - f(x)| \leq |h| + 2|(x+h)^2| + 2|x^2| \rightarrow 0$  as  $h, x \rightarrow 0$ ,  $f$  is continuous on a small interval  $I_1$ .

Since  $f'(x) = 1 + 2(2x \sin(\frac{1}{x}) - \cos(\frac{1}{x}))$ , there exists  $x$  in any interval contains 0 s.t.  $f'(x) < 0$ .

Therefore,  $f$  is neither increasing nor decreasing and not one to one in any interval contains 0.

Thus,  $f$  is not invertible near 0.

- (c) This is not contradict to inverse function theorem.

2.