

# Homework 13 of Introduction to Analysis(II)

AM15 黃琦翔 111652028

May 21, 2024

1. (a) Since a rectangle  $R$  is compact, for all  $\varepsilon > 0$ , we can find finite open cubes  $C_i$  s.t.  $\cup C_i \supseteq R$  and  $\text{Vol}(\cup C_i) < \text{Vol}(R) + \varepsilon$ . Therefore, for  $R_j$ , we can find  $\text{Vol}(\cup C_{i,j}) < \text{Vol}(R_j) + \frac{\varepsilon}{2^j}$ . And the total volume of  $\cup_{i,j} C_{i,j} < \text{Vol}(\cup R_j) + \varepsilon < 2\varepsilon$ . Thus, we can find a countable set of open cubes covers a measure zero set.  
  
(b) Since for any  $E_i$  is measure zero, we can find  $\{R_{i,j}\}_j$  s.t.  $\sum_j |R_{i,j}| < \frac{\varepsilon}{2^i}$ . Then, let  $R_{1,1} = R_1$ ,  $R_{1,2} = R_2$ ,  $R_{2,1} = R_3$ ,  $R_{1,3} = R_4$ ,  $R_{2,2} = R_5$  etc. And we can get  $\{R_k\}_k$  contains  $\bigcup_i E_i$  and  $\sum_k |R_k| < \varepsilon(\frac{1}{2} + \frac{1}{4} + \dots) = \varepsilon$ . Therefore, union of countable measure zero sets is measure zero.  
  
(c) Since  $A$  is compact and (a), we can find finite open cubes which covers  $A$  and volume of their union is less than  $\varepsilon$ . Thus, take union of closure on each open cube and we can find a rectangle covers  $A$  with volume less than  $\varepsilon$ . Therefore, volume of  $A$  is 0.
2. For any grid  $g$ ,  $U(1, g) = L(1, g)$ . Then, 1 is integrable. And for any  $\varepsilon > 0$ , there exists rectangles  $R_i$  s.t.  $|\sum |R_i| - \text{Vol}(E)| < \varepsilon$ . Thus,  $\int_E 1 dE \leq U(1, g) = \sum 1 \cdot |R_i| < \text{Vol}(E) + \varepsilon$ . Also,  $\int_E 1 dE \geq \text{Vol}(E) - \varepsilon$ . Therefore,  $\int_E 1 dE = \text{Vol}(E)$ .
3. (a) If  $x$  is rational,  $G(x, t) = \int_0^t 1 dy = t$ . If  $x$  is irrational,  $G(x, t) = \int_0^t 2y dy = t^2$ .  
Since  $t \in [0, 1]$ ,  $t^2 \leq t$ . Then, (U)  $\int_0^1 G(x, t) dx = t$  and (L)  $\int_0^1 G(x, t) dx = t^2$ .  
Therefore,  $\int_0^1 (\int_0^1 f(x, y) dy) dx = \int_0^1 G(x, 1) dx = \int_0^1 1 dx = 1$ .  
(b) As  $y \rightarrow 0$ , (U)  $\int_0^1 f(x, y) dx = 1$  but (L)  $\int_0^1 f(x, y) dx = 2y \rightarrow 0$ . Thus,  $\int_Q f(X) dx$  doesn't exist.