

# Homework 12 of Introduction to Analysis(II)

AM15 黃琦翔 111652028

May 9, 2024

1. (a) Since  $E$  is Jordan region,  $\text{Vol}(\partial E) = 0$ .

(b) Since  $\text{cl}(E) = \text{int}(E) \cup \partial E$  and  $\text{int}(E) \subseteq E \subseteq \text{cl}(E)$ ,

$$\text{Vol}(\text{cl}(E)) \leq \text{Vol}(\text{int}(E)) + \text{Vol}(\partial E) = \text{Vol}(\text{int}(E)) \leq \text{Vol}(E) \leq \text{Vol}(\text{cl}(E)).$$

Therefore,  $\text{Vol}(\text{cl}(E)) = \text{Vol}(\text{int}(E)) = E$ .

(c)

( $\implies$ ) From (b), we know  $\text{Vol}(\text{int}(E)) = \text{Vol}(E) > 0$ , then we can find a set of rectangles  $R_n$  s.t.

$$\sum |R_n| > 0 \text{ and } \cup R_n \subseteq \text{int}(E). \text{ Therefore, } \text{int}(E) \neq \emptyset.$$

( $\impliedby$ ) Since  $\text{int}(E)$  is non-empty, for any  $x_0 \in \text{int}(E)$ , there exists  $\varepsilon > 0$  s.t.  $D(x_0, \varepsilon) \subseteq \text{int}(E)$ . Then,

we can find a small rectangle  $R$  with each length is  $\frac{\varepsilon}{2}$  and  $R$  is contained in  $D(x_0, \varepsilon)$ . Thus,

$$\text{Vol}(\text{int}(E)) > \left(\frac{\varepsilon}{2}\right)^2 > 0.$$

(d) Since  $f$  is continuous, for any  $x_0 \in [a, b]$ , we can find a sequence  $x_k \rightarrow x_0$  s.t.  $f(x_k) \rightarrow f(x_0)$ . That

is,  $A = \{(x, f(x)) \mid x \in [a, b]\}$  is closed. And since  $\partial A \subseteq A$ ,  $\text{Vol}(\partial A) \leq \text{Vol}(A)$ .

Since  $f$  is continuous on  $[a, b]$  is compact, for all  $\varepsilon > 0$ , we can find  $\delta > 0$  s.t.  $|x - y| < \delta$  implies

That  $|f(x) - f(y)| < \frac{\varepsilon}{b-a}$ . Then, we can find a finite increasing sequence  $\{x_i \mid x_i \in [a, b]\}_{i=1}^N$  s.t.

$[a, b] \subseteq D(x_i, \frac{\delta}{2})$ . Therefore, for any  $y = f(x)$ ,  $y \in D(f(x_i), \frac{\varepsilon}{b-a})$  for some  $i$ .

Then, take  $u_0 = a$ ,  $u_i \in D(x_i, \frac{\delta}{2}) \cap D(x_{i+1}, \frac{\delta}{2})$ ,  $u_N = b$ , and we can get  $[a, b] = \cup [u_i, u_{i+1}]$ . Thus,

$A \subseteq \bigcup_{i=0}^N [u_i, u_{i+1}] \times D(\xi_i, \frac{\varepsilon}{b-a})$  for some  $\xi_i \in \{f(x) \mid x \in [u_i, u_{i+1}]\}$  with the sum of the rectangles is  $\frac{\varepsilon}{b-a} \cdot (b-a) = \varepsilon$ . Hence,  $\text{Vol}(A) = 0$  and  $\text{Vol}(\partial A) = 0$ .

(e) Yes. Since  $f$  is integrable, for any  $\varepsilon > 0$ , we can find a partition  $P$  s.t.  $U(f, P) - L(f, P) < \varepsilon$ .

Thus,