Homework 2 of Introduction to Analysis(II)

AM15 黃琦翔 111652028

March 1, 2024

1. Suppose $f_k(x) = \sum_{n=1}^{k} \frac{x}{n^\alpha(1+nx^2)}$ and $E_I = [-L, L]$ for $L \in \mathbb{N}$. Then, we want to proof that for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ s.t. $|f_k(x) - f_l(x)| < \varepsilon$ for all $k, l > N$ and all $x \in I_L$.

   First, suppose that $l > k > N$, then

$$
\begin{aligned}
|f_k(x) - f_l(x)| &= \sum_{n=k}^{l} \frac{x}{n^\alpha(1+nx^2)} \\
&\leq \sum_{n=k}^{l} \frac{x}{n^{\alpha+1}x^2} \\
&= \sum_{n=k}^{l} \frac{1}{n^{\alpha+1}x} \\
&=
\end{aligned}
$$

2. Since $f_k \to f$ uniformly and f_k are continuous, f is continuous. Then, for any $\varepsilon > 0$, we have $\delta > 0$ s.t. if $|y - y'| < \delta$ then $|f(y) - f(y')| < \frac{\varepsilon}{2}$ for all $y, y' \in \mathbb{R}$. Since $x_k \to x$, there exists $N_1 \in \mathbb{N}$ s.t. $|x_k - x| < \delta$ for all $k > N_1$. Also we have $N_2 \in \mathbb{N}$ s.t. $|f_k(x) - f(x)| < \frac{\varepsilon}{2}$ for all $k > N_2$.

   Then, take $N = \max\{N_1, N_2\}$, we can get $|f_k(x_k) - f(x)| \leq |f_k(x_k) - f(x_k)| + |f(x_k) - f(x)| = \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$ for all $k > N$. Thus, $\lim_{k \to \infty} f_k(x_k) = f(x)$.

3. Since f_n are continuous and converges uniformly, f is continuous(also integrable on $[0,1]$). And since f is continuous, we can find the maximun of $|f|$ on $[0,1]$ which is called as M. For any $\varepsilon > 0$, there exists $N_1 \in \mathbb{N}$ s.t. $|f_k(x) - f(x)| < \frac{\varepsilon}{4}$ for all $k > N_1$. And there exists $N_2 \in \mathbb{N}$ s.t. $N_2 > \frac{M}{2\varepsilon}$. Then, take

$n > N = \max N_1, N_2$

$$\begin{aligned}\left|\int_0^{1-\frac{1}{n}} f_n(x)dx - \int_0^1 f(x)dx\right| &\leq \left|\int_0^{1-\dfrac{1}{n}} f_n(x) - f(x)dx\right| + \left|\int_{1-\dfrac{1}{n}}^1 f(x)\right| \\ &\leq \int_0^1 |f_k(x) - f(x)|dx + \frac{1}{n}\cdot M \\ &\leq 1\cdot 2\frac{\varepsilon}{4} + \frac{\varepsilon}{2} \\ &= \varepsilon\end{aligned}$$

Thus, $\lim_{n\to\infty}\int_0^{1-\frac{1}{n}} f_n(x)dx = \int_0^1 f(x)dx.$

4.