

Homework 4 of Introduction to Analysis(II)

AM15 黃琦翔 111652028

March 14, 2024

1. Since B is equicontinuous, for any $f \in B$ and $\varepsilon > 0$, there exists $\delta > 0$ s.t. for any $x, y \in A$, $|x - y| < \delta$ implies that $|f(x) - f(y)| < \frac{\varepsilon}{2}$.

2. First, we want to show that B is closed. For any sequence $f_k \in B$ which converges to f ,

Then, since $|f'(x)| \leq 1$ for all $f \in B$ and $x \in (0, 1)$. For any $\varepsilon > 0$, we take $\delta < \varepsilon$, for any $x, y \in [0, 1]$ s.t. $|x - y| < \varepsilon$, $|f(x) - f(y)| \leq 1 \cdot |x - y| = \delta < \varepsilon$. Thus, B is equicontinuous.

Last, for any $x \in [0, 1]$, we want to proof B_x is compact. For $x = 0$, $B_0 = \{0\}$ obviously compact. For $x > 0$, since $|f'| \leq 1$, we can easily get $B_x = [-x, x]$ by $f(x) = ax$ for all $|a| \leq 1$. Thus, B is pointwise compact.

Therefore, by Arzela-Ascoli Theorem, B is compact.

3.