

Exercises(15)

December 19, 2023

1. Let f be continuous at each point (x, y) of a rectangle

$$Q = \{(x, y) : a \leq x \leq b, c \leq y \leq d\}.$$

Let F be the function defined on $[c, d]$ by the equation

$$F(y) = \int_a^b f(x, y) dx.$$

- (a) Prove that F is continuous on $[c, d]$. In other words, if $y_0 \in [c, d]$, we have

$$\lim_{y \rightarrow y_0} \int_a^b f(x, y) dx = \int_a^b \lim_{y \rightarrow y_0} f(x, y) dx = \int_a^b f(x, y_0) dx.$$

- (b) If the partial derivative $\frac{\partial f}{\partial y}$ is continuous on Q , show that for each $y \in (c, d)$,

$$F'(y) = \int_a^b \frac{\partial f}{\partial y}(x, y) dx.$$

Note: The partial derivative $\frac{\partial f}{\partial y}$ is defined by

$$\frac{\partial f}{\partial y}(x, y) = \lim_{h \rightarrow 0} \frac{f(x, y+h) - f(x, y)}{h}.$$

2. Define

$$f(x) = \left(\int_0^x e^{-t^2} \right)^2, \quad g(x) = \int_0^1 \frac{e^{-x^2(t^2+1)}}{t^2+1} dt.$$

- (a) Show that $g'(x) + f'(x) = 0$ for all x and deduce that $g(x) + f(x) = \pi/4$.
(b) Use (a) to prove that

$$\lim_{x \rightarrow \infty} \int_0^x e^{-t^2} dt = \frac{\sqrt{\pi}}{2}.$$

3. (a) Suppose that f is improperly integrable on $[0, \infty)$. Prove that if $L = \lim_{x \rightarrow \infty} f(x)$ exists, then $L = 0$.
(b) Let

$$f(x) = \begin{cases} 1 & n \leq x < n + 2^{-n}, n \in \mathbb{N} \\ 0 & \text{otherwise.} \end{cases}$$

Prove that f is improperly integrable on $[0, \infty)$ but $\lim_{x \rightarrow \infty} f(x)$ does not exist.