

# Homework 6 of Introduction to Analysis (I), Honor Class

AM15 黃琦翔 111652028

October 20, 2023

1. First, we show that  $S$  is closed and bounded. There exists  $D(2,2) \subseteq S \implies S$  is bounded. And for  $x \in \mathbb{Q} \setminus S$ , if  $x \leq \sqrt{2}$ ,  $D(x, d(x, \sqrt{2})) \subseteq \mathbb{Q} \setminus S$ . If  $x \geq \pi$ ,  $D(x, \pi) \subseteq \mathbb{Q} \setminus S \implies \mathbb{Q} \setminus S$  is open. Then,  $S$  is closed.

$G_i = (\sqrt{2} - \frac{1}{n}, \pi - \frac{1}{n}) \cap \mathbb{Q}$ , then  $\bigcap_{i=1}^{\infty} G_i$  is an open cover of  $S$  but doesn't have finite subcover. Thus,  $S$  is not compact.

2.  $\bigcap_{n=1}^{\infty} V_n = \emptyset$ .

3. Since  $x, y \in K$ , then assume there doesn't exist a set in  $G$  s.t. it contains both of  $x, y$ , then  $G$  doesn't contain one of  $x, y$  or both neither. Thus,  $G$  doesn't cover  $K$ , contradict to  $G$  is a open cover. Thus, there exists a set in  $G$  containing both  $x, y$ .

4. Since  $A$  is totally bounded, for all  $\varepsilon > 0$ , exists  $\{x_i\}_{i=1}^N$  is a finite set s.t.  $A \subseteq \bigcup_{i=1}^N D(x_i, \varepsilon)$ .

And  $M$  is complete  $\implies$  every Cauchy sequence in  $M$  converge to a point in  $M$ .

Then, for any sequence  $\{y_i\}$  in  $\bar{A}$ , if there is not any subsequence converge to  $\bar{A}$ , there exists  $\varepsilon' > 0$  s.t.  $|y_m - y_n| > \varepsilon'$  for all  $m, n > N \in \mathbb{N}$ .  $\bigcup D(y_i, \varepsilon')$  will be a infinite open cover of  $A$ , which contradicts to there has a finite  $N$  s.t.  $\bigcup D(x_i, \varepsilon')$  is cover of  $A$ .

Thus, every sequence in  $\bar{A}$  have subsequence converge to a point in  $\bar{A} \implies \bar{A}$  is sequence compact.

Then, by BWT,  $\bar{A}$  is compact.