

Homework 4 of Introduction to Analysis (I), Honor Class

AM15 黃琦翔 111652028

October 5, 2023

1. Since $A \subseteq \mathbb{R}^n$ is open, then for any point $x \in A$, exists some $\varepsilon > 0$ s.t. $B(x, \varepsilon) \subset A$.
2. (a) i. x is an accumulation point, for all $\varepsilon > 0$, $D(x, \varepsilon) \setminus \{x\} \cap A \neq \emptyset \implies D(x, \varepsilon) \cap A \neq \emptyset$. Thus, x is a limit point.
ii. Let $M = \mathbb{R}$, $d(x, y) = |x - y|$, $A = \{n \mid n \in \mathbb{N}\}$, then 1 is a limit point of A since $B(0, \varepsilon) \cap A \neq \emptyset$ for all $\varepsilon > 0$.
But for $0 < \varepsilon < 1$, $B(1, \varepsilon) \setminus \{1\} \cap A = \emptyset$. Thus, limit point is not accumulation all the time.
- (b) Assume there exists a limit point x of A is not in A . Then, $x \in A^c$. We want to show that A is not close.

For A^c , since all neighborhood U of x has intersection with A , then $U \not\subset A^c$. Thus, A^c is not open.

Therefore, if A is close, A contains all its limit points.

3. (a) Since $x \in A'_1$, for all $\varepsilon > 0$, $(D(x, \varepsilon)/\{x\}) \cap A_1 \neq \emptyset$. Then, let $y \in (D(x, \varepsilon)/\{x\}) \cap A_1$, $d(x, y) < \varepsilon$ for all $\varepsilon > 0 \implies \inf\{d(x, y) \mid y \in A_1\} = 0$.

(b) (\subseteq) For $x \in B'_n$, for any $\varepsilon > 0$, $(D(x, \varepsilon)/\{x\}) \cap B_n = (D(x, \varepsilon)/\{x\}) \cap (\cup A_i)$
 $= \cup((D(x, \varepsilon)/\{x\}) \cap A_i) \neq \emptyset$. Thus, $(D(x, \varepsilon)/\{x\}) \cap A_i \neq \emptyset$ for some $i \implies x \in \cup A'_i$.

(\supseteq) If $x \in A'_i$, for any $\varepsilon > 0$, $(D(x, \varepsilon)/\{x\}) \cap A_i \neq \emptyset \implies \emptyset \neq (D(x, \varepsilon)/\{x\}) \cap (\cup A_i)$
 $= (D(x, \varepsilon)/\{x\}) \cap B_n$.

Thus, $B'_n = \cup A'_i$.

4. False.

Let $A_i = (1/i, 1)$. Then, $B = (0, 1)$. And for 0, $0 \in B'$ but not in $\cup A'_i$.