Homework 10 of Introduction to Analysis (I), Honor Class

AM15 黃琦翔 111652028

November 16, 2023

- 1. (a)
 - (\Longrightarrow) Since f is continuous, then for $x \in \bar{A}$, we can find a $\{x_i \mid x_i \in A\}$ converges to x. Thus, for all $\varepsilon > 0$, exists $\delta > 0$ s.t. $d(x_i, x) < \delta \implies \rho(f(x_i), f(x)) < \varepsilon$. Then, for $y \in f(\bar{A})$, $D(y, \varepsilon) \cap f(A) \neq \emptyset$ for all $\varepsilon > 0$. Thus, $y \in \overline{f(A)}$.
 - $(\longleftarrow) \ \ \text{For} \ V \ \text{closed in} \ T \text{, then} \ f(\text{cl}(f^{-1}(V))) \subseteq \text{cl}(V) = V. \ \text{Applied} \ f^{-1} \ \text{on the both side, cl}(f^{-1}(V)) \subseteq f^{-1}(\text{cl}(V)) = f^{-1}(V). \ \ \text{Thus, } f^{-1}(V) \ \text{ is closed in} \ V \ \text{ is closed implies} \ f \ \text{ is continuous}.$
 - (b) Since f(p) = g(p) for all $p \in S$ and $\bar{E} = S$. For any $x \in S$, $x \in \bar{E}$, then we can find a sequence $\{x_i\} \subseteq E$ converges to x.

Since
$$f(x_i) = g(x_i)$$
 for all $i \in \mathbb{N}$, $|f(x) - g(x)| \le |f(x) - f(x_i)| + |f(x_i) - g(x)| = |f(x) - f(x_i)| + |g(x) - g(x_i)| \to 0$ as $i \to \infty$. Then, $f(x) = g(x)$ for all $x \in S$.

2. Since B is compact, f(B) is compact. Then, we want to show for an sequence y_k converges to $y \in f(B)$, we can get $f^{-1}(y_k)$ converges to $f^{-1}(y)$. Since f is one-to-one, we can find $x_k = f^{-1}(y_k) \in B$ and $x = f^{-1}(y)$. Then, assume x_k is not converges to x, $|x_k - x| > \delta$ for some $\delta > 0$ and $k \in \mathbb{N}$. Then, $f(D(x,\delta) \cap B) \not\subseteq D(f(x),\varepsilon)$ since f is one-to-one. Thus, f is not continuous.(contradiction) Therefore, f^{-1} is continuous.

3. Since f is strictly increasing, f is one-to-one. By question 2, f^{-1} is continuous, too.

Then, we want to check f^{-1} is strictly increasing. For $f(x_1) < f(x_2)$ with $x_1, x_2 \in \mathbb{R}$. Assume $x_1 > x_2$. Then, $f(x_1) > f(x_2)$.(contradiction)

Thus, $f(x_1) < f(x_2) \implies f^{-1}(f(x_1)) = x_1 < x_2 = f^{-1}(f(x_2))$. Therefore, f^{-1} is continuous and strictly increasing.