

$$\begin{array}{c}
\begin{array}{ccccc}
& & R\chi_E^M & & \\
& & \downarrow & & \\
& & R^2G(SM, E) & \xrightarrow{R\nu_{M,E}} & RG(M, TE) \\
& \nearrow R^2G(\alpha_M^S, E) & \searrow R^2G(S\alpha_M^S, E) & & \downarrow \text{nat } \nu \\
R^2G(M, E) & & R^2G(S^2M, E) & & RG(\alpha_M^S, TE) \\
& \searrow R^2G(\alpha_M^S, E) & \nearrow R^2G(\mu_M^S, E) & \searrow R\nu_{SM,E} & \downarrow \\
& R^2G(SM, E) & & RG(SM, TE) & \\
& \downarrow \text{nat } \mu & & \downarrow \nu_{M,TE} & \\
& R^2G(SM, E) & & G(M, T^2E) & \\
& \downarrow \mu_{G(SM,E)}^R & & \downarrow G(M, \mu_E^T) & \\
RG(M, E) & \xrightarrow{RG(\alpha_M^S, E)} & RG(SM, E) & \xrightarrow{\nu_{M,E}} & G(M, TE) \xleftarrow{G(M, \mu_E^T)} G(M, T^2E)
\end{array}
\end{array}$$

(28)

$\mu_{G(M,E)}^R$ (vertical arrow from $R^2G(M, E)$ to $RG(M, E)$)
 χ_E^M (curved arrow from $RG(M, E)$ to $G(M, TE)$)
 χ_{TE}^M (curved arrow from $RG(M, TE)$ to $G(M, T^2E)$)