

# STATS 210P

## Lecture 4

Sevan Koko Gulesserian

University of California, Irvine

# Simple Linear Regression: Model Specification

To recap the set up:

The simple linear regression model for the population is:

$$Y = \beta_0 + \beta_1 X + \varepsilon.$$

where the  $\varepsilon$ 's are independent and  $\varepsilon \sim \text{Normal}(0, \sigma_\varepsilon^2)$  ( $\sim$  notation meant "follows").

$\sigma_\varepsilon$  = standard deviation of the errors = standard deviation of the Y values at each X value.

- The above is meant to imply that  $Y|X \sim N(\beta_0 + \beta_1 X, \sigma_\varepsilon^2)$
- Remember that the notation  $Y|X$  means Y given X.
- At the unit level this population model is:  $Y_i = \beta_0 + \beta_1 X_i + \varepsilon_i$ , where the  $\varepsilon_i$ 's are independent and  $\varepsilon_i \sim \text{Normal}(0, \sigma_\varepsilon^2)$ .
- The sample model is as follows:  $\hat{Y} = \hat{\beta}_0 + \hat{\beta}_1 X$ .

# Simple Linear Regression: Model Specification

- Would like to obtain estimates of  $\beta_0$  and  $\beta_1$  based on observed data.
- The estimates will be noted as  $\hat{\beta}_0$  and  $\hat{\beta}_1$ , and will be used to estimate  $\hat{Y}_i$  (the predicted value of  $Y_i$  given a  $X_i$  value).
- Will do this by minimizing the residuals, noted as  $r_i$  or  $e_i$ , across all observation.

$$\bullet \quad r_i = Y_i - \hat{Y}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i)$$

- Note we will set  $\bar{X}$  and  $\bar{Y}$  to be used as the sample means of the X's and the Y's respectively (for example  $\bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i$ ).

# Simple Linear Regression: Slope Inference

The estimates of  $\beta_0$  and  $\beta_1$  minimizes the following function (also called the objective function):

$$f(\hat{\beta}_0, \hat{\beta}_1) = \sum_{i=1}^n r_i^2 = \sum_{i=1}^n (Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i))^2$$

The solutions for  $\hat{\beta}_0$  and  $\hat{\beta}_1$  are as follows:

- $\hat{\beta}_1 = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sum_{i=1}^n (X_i - \bar{X})^2}$ , note that this is the sample covariance between  $X$  and  $Y$  divided by the sample variance of  $X$ .
- $\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{X}$ , note that is a function of the slope estimate.

(How we get these is by minimizing  $f(\hat{\beta}_0, \hat{\beta}_1)$ . That is to say take the partial derivative of the function with respect to  $\hat{\beta}_1$ , set it to 0, and solve for  $\hat{\beta}_1$ . And then do the same for  $\hat{\beta}_0$ . And of course do a second derivative test.)

# Simple Linear Regression: Slope Inference

Side note.

- The solutions for  $\beta_0$  and  $\beta_1$  are the same had we done a maximum likelihood approach.
- In the maximum likelihood approach, we assume the  $Y_i$ 's are independent from a normal distribution with mean/expectation  $\beta_0 + \beta_1 X_i$  and variance  $\sigma_\epsilon^2$ .
- We construct the likelihood using the form of the normal/Gaussian distribution density, and then maximize it with respect to  $\beta_0$  and  $\beta_1$ .

# Simple Linear Regression: Slope Inference

Estimates of  $\beta_0$  and  $\beta_1$ .

- Our primary concern is inference on the slope parameter  $\beta_1$  (as this determines the relationship between X and Y).
- The estimate,  $\hat{\beta}_1$  is function of the Y's, which are random variables (also note that the estimate  $\hat{\beta}_1$  will vary from sample to sample).
- And so  $\hat{\beta}_1$  has an approximate Normal distribution, with expectation equal to the true  $\beta_1$  and variance equal to 
$$\frac{\sigma_\epsilon^2}{\sum_{i=1}^n (X_i - \bar{X})^2}.$$
- Note we do not know the true  $\sigma_\epsilon^2$  so we need to replace it with an estimate  $\hat{\sigma}_\epsilon^2$ .
- The idea is that  $\hat{\beta}_1$  will vary from sample to sample based on the population (and thus will have a distribution with expectation and a variance).

# Simple Linear Regression: Slope Inference

As a result, we can conduct hypothesis tests on  $\beta_1$  based on the estimate  $\hat{\beta}_1$ .

Will use the form of the solution of  $\hat{\beta}_1$  to derive the needed quantities to create a test statistic, and determine its distribution.

We are mainly interested in  $\beta_1$ . Just like with the example of  $\rho$ , we are usually interested in if the parameter values as compared to 0 (since  $\beta_1 = 0$  means  $X$  has no association with  $Y$ ).

In simple linear regression, testing  $\beta_1$  being equal to 0, greater than 0, or less than 0 is the same as testing if the correlation coefficient between  $X$  and  $Y$  is equal to 0, greater than 0, or less than 0 respectively.

# Simple Linear Regression: Slope Inference

Just like in previous set of slides, with inference on the parameters  $\beta$  follow the same hypothesis test procedure.

First state the null and alternative hypothesis. Either one of the following:

- $H_0 : \beta = a$  (or  $H_0 : \beta \leq a$ ) and  $H_a : \beta > a$
- $H_0 : \beta = a$  (or  $H_0 : \beta \geq a$ ) and  $H_a : \beta < a$
- $H_0 : \beta = a$  and  $H_a : \beta \neq a$

**Often the interest is when  $a = 0$ .**

The R output will automatically test the scenario of  $H_0 : \beta = 0$  and  $H_a : \beta \neq 0$



# Simple Linear Regression: Slope Inference

For a constant  $a$  that is specified in the null and alternative hypothesis (usually  $a=0$ ), create a test statistic  $t^*$  as follows:

- $$t^* = \frac{\hat{\beta}_1 - a}{\text{se}(\hat{\beta}_1)} = (\hat{\beta}_1 - a) / \left( \frac{\sqrt{MSE}}{\sqrt{\sum_{i=1}^n (X_i - \bar{X})^2}} \right)$$
- The denominator term above in the big parenthesis' is called "Std. Error" in R .
- Variance is  $\text{var}(\hat{\beta}_1) = \frac{\sigma_\varepsilon^2}{\sum_{i=1}^n (X_i - \bar{X})^2}$  and under the null hypothesis we have expectation being  $E(\hat{\beta}_1) = a$ .
- And since we do not know the true  $\sigma_\varepsilon^2$  value, we replace it with a sample based estimate  $\hat{\sigma}_\varepsilon^2$  (called MSE, mean squared error).
- $t^*$  follows a  $t$  distribution with  $n - 2$  degrees of freedom.
- Note that  $MSE = \hat{\sigma}_\varepsilon^2$ .
- In the R output, we have all the needed information to test the scenario of  $H_0 : \beta = 0$  and  $H_a : \beta \neq 0$ , that is to say we have the test statistic value and the 2-sided p-value for this test given to us by default.




# Simple Linear Regression: Slope Inference

Now, with  $t^*$  calculated and known to follow  $t$  distribution with  $n - 2$  degrees of freedom, and with  $H_0$  and  $H_a$  set, can compute  $p$ -value.

For  $H_a$  ***less than***, the  $p$ -value is the area below  $t$ , even if  $t$  is positive.

For  $H_a$  ***greater than***, the  $p$ -value is the area above  $t$ , even if  $t$  is negative.

For  $H_a$  ***two-sided***,  $p$ -value is  $2 \times$  area above  $|t|$ .

| Statement of $H_a$           | $p$ -Value Area                                | $t$ -Curve Region                                                                   |
|------------------------------|------------------------------------------------|-------------------------------------------------------------------------------------|
| $\mu < \mu_0$ (less than)    | Area to the left of $t$<br>(even if $t > 0$ )  |   |
| $\mu > \mu_0$ (greater than) | Area to the right of $t$<br>(even if $t < 0$ ) |   |
| $\mu \neq \mu_0$ (not equal) | $2 \times$ area to the right of $ t $          |  |

# Simple Linear Regression: Slope Inference

A significance level,  $\alpha$ , is specified and set. Based on the computed p-value, a conclusion can be made.

- If p-value is less than  $\alpha$ , reject the null and conclude evidence for the alternative. We have statistically significant results.
  - Reject the null that  $\beta = a$  and conclude evidence that  $\beta \neq a$  (or  $\beta < a$  or  $\beta > a$ ).
  - When the alternative is  $H_a : \beta_1 \neq 0$ , then rejecting the null and going with the alternative means that we have evidence that  $X$  has an association with  $Y$ .
- If p-value is greater than  $\alpha$ , fail to reject the null and don't conclude evidence for the alternative. We don't have statistically significant results.
  - Fail to reject the the null hypothesis.
  - When we fail to reject the null (that is to say we go with the null)  $H_0 : \beta_1 = 0$ , this means that we do not have evidence that  $X$  has an association with  $Y$ .

# Simple Linear Regression: Fitting the Model

Example: Return to the skin cancer data set. Say  $Y$ =Mortality and  $X$ =Longitude.

Output from R is as follows.

```
lm(formula = Mort ~ Long, data = skincancer)
```

Residuals:

| Min     | 1Q      | Median | 3Q     | Max    |
|---------|---------|--------|--------|--------|
| -63.898 | -25.995 | -5.952 | 21.856 | 78.444 |

Coefficients:

|             | Estimate | Std. Error | t value | Pr(> t )    |
|-------------|----------|------------|---------|-------------|
| (Intercept) | 182.7696 | 29.8893    | 6.115   | 1.8e-07 *** |
| Long        | -0.3287  | 0.3245     | -1.013  | 0.316       |

---

Residual standard error: 33.42 on 47 degrees of freedom

Multiple R-squared: 0.02137, Adjusted R-squared: 0.0005491

F-statistic: 1.026 on 1 and 47 DF, p-value: 0.3162

# Simple Linear Regression: Slope Inference

Want to test  $H_0 : \beta_1 = 0$  against  $H_a : \beta_1 \neq 0$ .

- This is to say that we want to investigate if longitude is able to explain (or predict) mortality.
- With  $H_a : \beta_1 \neq 0$ , we are not concerned with the direction of the association, just whether there is a significant one.
- The R output gives all the needed pieces.
  - The output gives the estimate of  $\beta$ , the standard error of the estimate, the computed  $t^*$  statistics (assuming  $a = 0$ ), and the two sided p-value ( $H_a : \beta_1 \neq 0$ ).

# Simple Linear Regression: Slope Inference

The p-value of R is denoted as  $\Pr(>|t|)$ .

This is the two sided p-value. Use this to test  $H_a : \beta_1 \neq 0$ . Say this 2-sided p-value is  $p$ .

- If testing  $H_a : \beta_1 < 0$ .
  - If  $t^*$  (t-value in R) is negative, then take the p-value given by R ( $\Pr(> |t|)$ ), and divide it by two ( $\frac{p}{2}$ ).
  - If  $t^*$  is positive, then take the p-value given by R, and divide it by two and subtract it from one ( $1 - \frac{p}{2}$ ).
- If testing  $H_a : \beta_1 > 0$ .
  - If  $t^*$  is positive, then take the p-value given by R ( $\Pr(> |t|)$ ), and divide it by two ( $\frac{p}{2}$ ).
  - If  $t^*$  is negative, then take the p-value given by R, and divide it by two and subtract it from one ( $1 - \frac{p}{2}$ ).

# Simple Linear Regression: Slope Inference

Assume  $\alpha = 0.05$  (a 5% significance level).

- In the skin cancer longitude example,  $t^* = -1.013$ .
- The two-sided p-value is 0.316 (this is already in the output, but can also get it as follows:  $2 * (1 - pt(1.013, 47))$ ).
- P-value is greater than significance level ( $0.316 > 0.05$ ).
- Fail to reject the null.
- Conclude that we don't have evidence that  $\beta_1 \neq 0$ . There is no evidence that longitude predicts mortality rates (at a 5% significance level).

# Simple Linear Regression: Fitting the Model

Example: Using the same dataset, set  $Y$ =Mortality and  $X$ =Latitude.

```
lm(formula = Mort ~ Lat, data = skincancer)
```

Residuals:

|  | Min     | 1Q      | Median | 3Q     | Max    |
|--|---------|---------|--------|--------|--------|
|  | -38.972 | -13.185 | 0.972  | 12.006 | 43.938 |

Coefficients:

|             | Estimate | Std. Error | t value | Pr(> t )     |
|-------------|----------|------------|---------|--------------|
| (Intercept) | 389.1894 | 23.8123    | 16.34   | < 2e-16 ***  |
| Lat         | -5.9776  | 0.5984     | -9.99   | 3.31e-13 *** |
| ---         |          |            |         |              |

Residual standard error: 19.12 on 47 degrees of freedom  
Multiple R-squared: 0.6798, Adjusted R-squared: 0.673  
F-statistic: 99.8 on 1 and 47 DF, p-value: 3.309e-13



# Simple Linear Regression: Slope Inference

Assume  $\alpha = 0.05$  (a 5% significance level). Want to test  $H_0 : \beta_1 = 0$  against  $H_a : \beta_1 \neq 0$ .

- This is to say that we want to investigate if latitude is able to explain (or predict) mortality.
- The test statistic  $t^*$  is -9.99.
- Two sided p-value is approximately 0 ( $3.31 * 10^{-13}$ ).
- Reject the null hypothesis and conclude evidence that  $\beta_1 \neq 0$ .  
Conclude evidence that latitude does predict mortality rates (at a 5% significance level).

# Simple Linear Regression: Slope Inference

Note that we could have tested alternatives of the form  $H_a : \beta_1 > 0$  or  $H_a : \beta_1 < 0$  with just the R output as well.

- Can use the test statistic given,  $t^*$ , and degrees of freedom ( $n - 2$ ) to compute p-value (slide 10).
- Or can use p-value given (two sided) and convert it to the needed one sided p-value for the alternative hypothesis (slide 14).
- In the skin cancer data with  $X$ =latitude, the p-value for the test of  $H_a : \beta_1 < 0$  is also 0 (divide 2-sided p-value by 2).
- Conclude evidence that latitude does predict expected mortality rates (at a 5% significance level) and that the association is negative (as latitude increases, predicted/expected mortality decreases).

# Simple Linear Regression: Slope Inference

Getting p-values in R using `pnorm` and `pt` functions.

- To get the area below a test statistic  $t^*$  using a standard normal distribution, that will be `pnorm( $t^*$ )`.
- Thus if you need the area above, it is `1-pnorm( $t^*$ )`.
- And so the two sided p-value would be `2*(1-pnorm( $t^*$ ))`.
- To use the t-distribution with  $n-k$  (in our case  $k=2$  to be  $n-2$  degrees of freedom) to get area below, we would do `pt( $t^*$ ,  $n-k$ )` and to get area above would do `1-pt(| $t^*$ |,  $n-k$ )`.
- And so two sided p-value using t-distribution with  $n-k$  degrees of freedom would be `2*(1-pt(| $t^*$ |,  $n-k$ ))`.

# Simple Linear Regression: Fitting the Model

Example: Using the med school data set. Fitting a model to  $Y =$  MCAT scores and  $X =$  GPA.

```
lm(formula = MCAT ~ GPA, data = mcat)
```

Residuals:

| Min      | 1Q      | Median  | 3Q     | Max    |
|----------|---------|---------|--------|--------|
| -11.4148 | -2.5168 | -0.1519 | 2.6653 | 8.6616 |

Coefficients:

|             | Estimate | Std. Error | t value | Pr(> t )     |
|-------------|----------|------------|---------|--------------|
| (Intercept) | 3.923    | 6.922      | 0.567   | 0.573        |
| GPA         | 9.104    | 1.942      | 4.688   | 1.97e-05 *** |
| ---         |          |            |         |              |

Residual standard error: 4.088 on 53 degrees of freedom  
Multiple R-squared: 0.2931, Adjusted R-squared: 0.2798  
F-statistic: 21.98 on 1 and 53 DF, p-value: 1.969e-05

# Simple Linear Regression: Slope Inference

Say we want to test if GPA is predictive of MCAT scores.

$$H_a : \beta_1 \neq 0.$$

- The test statistics is 4.688 and the two sided p-value is 0.000196.
- Assuming  $\alpha = 0.05$ , reject the null hypothesis. Conclude evidence that GPA is predictive of MCAT scores.
- In this example, the p-value for the test of  $H_a : \beta_1 > 0$  is also approximately 0 (half of 0.000196).
- Conclude evidence that GPA does predict MCAT scores (and more precisely GPA is positively related to MCAT scores).

# Simple Linear Regression: Slope Inference

Just as with testing hypothesis, can use the information obtained to create *confidence intervals*.

Can create confidence intervals for  $\beta_0$  and  $\beta_1$  based on a specified confidence level.

Confidence is defined to be  $1-\alpha$ , where  $\alpha$  is the significance level.

We read an interval  $(a, b)$  based on a specified significance level as follows: We are  $(1 - \alpha) * 100\%$  confident the parameter of interest is in the interval  $(a, b)$ , for example when  $\alpha = 0.05$ , confidence level is 95% .

- Confidence is defined as follows: If we were to repeat this procedure a very large (infinite) number of times (obtain data and calculate interval), then  $(1 - \alpha) * 100\%$  of the intervals computed will contain the true parameter value.

# Simple Linear Regression: Estimation and Prediction

Remember that a confidence interval was of the form:

$$\text{estimate} \pm \text{multiplier} * \text{s.e.}$$

- For our interests for now, estimate is the single point estimate of the unknown parameter  $\beta_1$ .
- Multiplier was the  $(1 - \alpha)$  critical value from the distribution used to obtain the interval.
  - For example using a standard normal distribution, this is the  $z^*$  value that has area below being  $1 - \alpha/2$ .
  - In R this is  $z^* = \text{qnorm}(1 - \frac{\alpha}{2})$ .
- The true multiplier is from a  $t$  distribution with d.f. =  $n - 2$  and area below is  $1 - \frac{\alpha}{2}$ , but since we will have more than enough data, the  $t$  multiplier will be approximately equal to the  $z^*$  multiplier (for the  $t$  multiplier, we would do  $\text{qt}(1 - \frac{\alpha}{2}, n - 2)$  in R).
- s.e. is the standard error (or standard deviation) of the estimate.

# Simple Linear Regression: Estimation and Prediction

- All needed pieces are given in R.
- Use `confint(model)` in R to obtain confidence intervals for  $\beta_0$  and  $\beta_1$ .
- By default, this will give 95% confidence intervals for each parameter.
- Can change the confidence level using an option: For example a 90% confident level, `confint(model, level=0.90)` .



# Simple Linear Regression: Estimation and Prediction

Example using the skin cancer dataset.

Let response  $Y$ =Mortality and  $X$ =Longitude.

Say the model in R is as follows:

- `model.long = lm(Mort~Long, data=skincancer)`

# Simple Linear Regression: Estimation and Prediction

Using `confint(model.long)`, the output from R is:

|             | 2.5 %       | 97.5 %      |
|-------------|-------------|-------------|
| (Intercept) | 122.6400848 | 242.8991374 |
| Long        | -0.9814471  | 0.3240217   |

This is read as follows:

- The 95% confidence interval for the intercept  $\beta_0$  is (122.6, 242.9)
- The 95% confidence interval for the slope  $\beta_1$  (longitude) is (-0.98, 0.32).
  - Note that 0 is in the interval.

# Simple Linear Regression: Estimation and Prediction

Another example using the skin cancer dataset.

Let response  $Y$ =Mortality and now let  $X$ =Latitude.

Say the model in R is as follows:

- `model.lat = lm(Mort ~ Lat, data=skincancer)`

# Simple Linear Regression: Estimation and Prediction

Using `confint(model.lat)`, the output from R is:

|             | 2.5 %      | 97.5 %     |
|-------------|------------|------------|
| (Intercept) | 341.285151 | 437.093552 |
| Lat         | -7.181404  | -4.773867  |

This is read as follows:

- The 95% confidence interval for the intercept  $\beta_0$  is (341.28, 437.09)
- The 95% confidence interval for the slope  $\beta_1$  (latitude) is (-7.18, -4.77).
  - Note that 0 is not in the interval, and the entire interval is below 0.

# Simple Linear Regression: Estimation and Prediction

Can also get intervals for different confidence levels. For example a 99% confidence level ( $\alpha = 0.01$ ).

Using `confint(model.lat, level=0.99)`, the output from R is:

|             | 0.5 %      | 99.5 %     |
|-------------|------------|------------|
| (Intercept) | 325.263865 | 453.114837 |
| Lat         | -7.583998  | -4.371274  |

This is read as follows:

- The 99% confidence interval for the intercept  $\beta_0$  is (325.26, 453.11).
- The 99% confidence interval for the slope  $\beta_1$  (latitude) is (-7.58, -4.37).
  - Note that 0 is not in the interval, and the entire interval is below 0.

# Simple Linear Regression: Estimation and Prediction

Note that as the confidence level increases, the width of the interval  $(b - a)$  gets larger (where  $(a,b)$  is our confidence interval).

This is a result of the multiplier in the formula for confidence intervals increases as confidence level increases.

This is to say the higher the confidence you want to attribute to the interval, the wider its going to be (and so the lower the confidence, the narrower the interval).

# Simple Linear Regression: Estimation and Prediction

Another way the confidence interval width is affected is the sample size.

Remember the multiplier is multiplied with the standard error (s.e.) of the estimate of the parameter (here  $\hat{\beta}_1$ ).

The larger the sample size, the lower the s.e. Therefore, as sample size increases, the interval width (at a given confidence level) will tend to decrease.

Hypothesis testing and confidence interval.

- Say we are given a 95% confidence interval,  $(l,u)$ , where  $l$ =lower number and  $u$ =upper number of the interval, for the unknown parameter  $\theta$ .
- Then we are 95% confident that the unknown parameter  $\theta$  is in the interval  $(l,u)$ .
- Thus any value in  $(l,u)$  we are 95% confident that that value can be the true value of  $\theta$ .



Hypothesis testing and confidence interval.

- The interval  $(l,u)$  are all the values which we will fail to reject the null hypothesis of  $\theta = \theta_0$  (for  $\theta_0$  in  $(l,u)$ ).
- For example let the interval be  $(5,10)$ .
- Then we would fail to reject  $\theta = \theta_0$  for all values of  $\theta_0$  in  $(5,10)$ .
- We will reject the null for all values of  $\theta_0$  outside of  $(5,10)$ .

# Simple Linear Regression: Estimation and Prediction

Remember that simple linear regression model for the population is:

$$Y = \beta_0 + \beta_1 X + \varepsilon.$$

where the  $\varepsilon$ 's are independent and  $\varepsilon \sim \text{Normal}(0, \sigma_\varepsilon^2)$

This implies that  $Y|X \sim N(\beta_0 + \beta_1 X, \sigma_\varepsilon^2)$

- $\mu_Y$  can be thought of as the true mean/expectation of  $Y$  at a given value of  $X$ .
- Can think of it as  $E(Y) = \mu_Y = \beta_0 + \beta_1 X$
- It is the true population mean of the  $Y$  values at a given value of  $X$ .

# Simple Linear Regression: Estimation and Prediction

- Can estimate  $\mu_Y$  by  $\hat{\mu}_Y$ .
- $\hat{\mu}_Y = \hat{\beta}_0 + \hat{\beta}_1 X$
- Can incorporate uncertainty about the estimate into a confidence interval instead of a single estimate.
- Want a confidence interval for  $\mu_Y$  at a certain value of  $X$ , say  $X = X_p$ .

# Simple Linear Regression: Estimation and Prediction

- The form of the confidence interval is like before:

$$\text{estimate} \pm \text{multiplier} * \text{s.e.}$$

- The estimate is  $\hat{\mu}_Y = \hat{\beta}_0 + \hat{\beta}_1 X$ .
- The multiplier is from a  $t$  distribution with  $n - 2$  degrees of freedom (with area below being  $1 - \frac{\alpha}{2}$ , but again can approximate this using standard normal distribution to get multiplier when we have large datasets).
- The standard error of the estimate is as follows:

$$\text{s.e.}(\hat{\mu}_Y) = \sqrt{MSE \left( \frac{1}{n} + \frac{(X_p - \bar{X})^2}{\sum_{i=1}^n (X_i - \bar{X})^2} \right)}$$

- The confidence interval will be:

$$\hat{\mu}_Y \pm t * \sqrt{MSE \left( \frac{1}{n} + \frac{(X_p - \bar{X})^2}{\sum_{i=1}^n (X_i - \bar{X})^2} \right)}$$

- Will read this interval,  $(a, b)$ , as that we are  $(1 - \alpha) * 100\%$  confident that the interval  $(a, b)$  contains the true value of  $\mu_Y$ .
- Remember that  $MSE$  was the estimate of  $\sigma_\varepsilon^2$ .

# Simple Linear Regression: Estimation and Prediction

To do this in R, use the following function:

```
predict(model, list(Newdata=X), interval= "c")
```

Example: Take the medical school data.  $Y$ =MCAT is the response and the explanatory variable is  $X$ =GPA.

Say you want a 95% confidence interval for the mean value of the response when  $GPA=4.0$  (that is to say  $X_p = 4.0$ ).

The command in R would be:

```
predict(model, list(GPA=4), interval= "c").
```

|   | fit      | lwr      | upr      |
|---|----------|----------|----------|
| 1 | 40.33983 | 38.27828 | 42.40138 |

# Simple Linear Regression: Estimation and Prediction

| fit      | lwr      | upr      |
|----------|----------|----------|
| 40.33983 | 38.27828 | 42.40138 |

How to read the output:

fit is the value of the prediction when GPA=4.0

lwr is the lower bound of the confidence interval.

upr is the upper bound of the confidence interval.

- The 95% confidence interval for the true mean of the response variable (MCAT scores) when GPA=4.0 is (38.28, 42.40).

# Simple Linear Regression: Estimation and Prediction

As before, can change the confidence level, say to 99%.

The command in R would be:

```
predict(model, list(GPA=4), interval= "c",  
level=0.99).
```

| fit      | lwr      | upr      |
|----------|----------|----------|
| 40.33983 | 37.80685 | 42.87281 |

How to read the output:

- The 99% confidence interval for the mean of the response variable (MCAT scores) when GPA=4.0 is (37.81, 42.87).
- Note how the interval got wider compared to the 95% confidence interval.



# Simple Linear Regression: Estimation and Prediction

Another example using the skin cancer dataset. Say  $Y$ =Mortality and  $X$ =Latitude.

We want a 95% confidence interval for the mean of mortality when latitude is 40.

```
predict(model.skin, list(Lat=40), interval="c")
```

|  | fit      | lwr      | upr      |
|--|----------|----------|----------|
|  | 150.0839 | 144.5617 | 155.6061 |

How to read the output:

- The predicted mortality rate when  $\text{Lat}=40$  is 150.08
- The 95% confidence interval for the mean of the response variable of Mortality rate when  $\text{Lat}=40$  is (144.56, 155.60).

# Simple Linear Regression: Estimation and Prediction

Up to now, we have created a confidence interval for  $\mu_Y$ , the true mean/expectation of the responses at a given value of  $X = X_p$ .

Now say we are interested in a confidence interval for the individual prediction  $\hat{Y}_p$ , the predicted value of  $Y$  given  $X = X_p$ .

Note that  $\hat{Y}_p = \hat{\beta}_0 + \hat{\beta}_1 X_p$ .

Just like with the mean response  $\mu_Y$ , we can create a confidence interval about  $Y_p$ .

# Simple Linear Regression: Estimation and Prediction

Note that  $\hat{Y} = \hat{\mu}_Y$  for a given value of  $X = X_p$ .

The confidence interval for  $Y_p$  will be the same as that for  $\mu_Y$ , with the exception of the standard error.

Remember the standard error in the confidence interval for  $\mu_Y$  is:

$$\sqrt{MSE \left( \frac{1}{n} + \frac{(X_p - \bar{X})^2}{\sum (X_i - \bar{X})^2} \right)}$$

The standard error for the confidence interval for  $Y_p$  is now:

$$\sqrt{MSE \left( 1 + \frac{1}{n} + \frac{(X_p - \bar{X})^2}{\sum (X_i - \bar{X})^2} \right)}$$

**These equations for the standard errors only apply to simple linear regression.**

# Simple Linear Regression: Estimation and Prediction

Note that the standard error for the individual prediction has an extra *MSE* term compared to that of the mean response.

This is because when it comes to the individual response  $Y_p$ , we have to account for the variation of  $\mu_Y$  as well as the error term ( $\varepsilon$ ).

As a result, for a given level of confidence, the confidence interval for  $Y_p$  will always be wider than that of  $\mu_Y$ .

Intervals for  $Y_p$  are called *prediction intervals*.

# Simple Linear Regression: Estimation and Prediction

To do this in R, use the function

```
predict(model, list(Newdata=X), interval= "p")
```

Note that before we had `interval="c"`, now it is `interval="p"`

Example: Take the medical school data.  $Y$ =MCAT is the response and the explanatory variable is  $X$ =GPA.

Say you want a 95% confidence interval for the value of the response when  $GPA=4.0$  (that is to say  $X_p = 4.0$ . The command in R would be:

```
predict(model, list(GPA=4), interval= "p").
```

|  | fit      | lwr      | upr      |
|--|----------|----------|----------|
|  | 40.33983 | 31.88554 | 48.79413 |

# Simple Linear Regression: Estimation and Prediction

- We interpret this as follows: We are 95% confident the individual response  $Y$  (MCAT score) at  $X = X_p$  (GPA=4.0) will be in the interval (31.88, 48.79).
- Note how the prediction interval is wider than the confidence interval.
- The 95% confidence interval for the mean response was (38.28, 42.40).
- The 95% confidence interval for the individual response is (31.88, 48.79)

# Simple Linear Regression: Estimation and Prediction

Another example using the skin cancer dataset. Say  $Y$ =Mortality and  $X$ =Latitude.

We want a 95% confidence interval for the individual response of mortality when latitude is 40.


```
predict(model.skin, list(Lat=40), interval="p")
```

|  | fit      | lwr     | upr      |
|--|----------|---------|----------|
|  | 150.0839 | 111.235 | 188.9329 |

How to read the output:

- The predicted mortality rate when  $\text{Lat}=40$  is 150.08
- The 95% confidence interval of the response variable when  $\text{Lat}=40$  is (111.23, 188.93).

# Simple Linear Regression: Estimation and Prediction





# Simple Linear Regression: Extrapolation

Issues with extrapolation.

- When we use the estimated regression line to predict a point whose X-value is outside the range of the X-values of dataset used to estimate the regression equation, it is called extrapolation.
- Example: The dataset has subjects ranging in age from 18 to 35 years old.
- Now say we want to obtain inference on a subject that is 75 years old. This data point is well outside the range of the data used to fit the model.
- Extrapolation does not give reliable predictions because the regression line may not be valid outside the dataset range.
- Our model should be used to predict and obtain inference on subjects within or near the range of 18 to 35 years old.