

Solution to exercise 1

In this exercise, you want to take care of possible overflows in the computation of

$$r := (a * \text{randomNumbers}[\text{indexOfInteger}] + c) \% \text{modulus}$$

under the assumption

$$\text{Long.MAX_VALUE} < a \cdot \text{modulus} + c < 2 \cdot \text{Long.MAX_VALUE}.$$

If the operation $a * \text{randomNumbers}[\text{indexOfInteger}] + c$ produces an overflow, i.e., if the result is bigger than Long.MAX_VALUE , the value produced in Java is negative (note that a multiple overflow is prevented by the assumption $a \cdot \text{modulus} + c < 2 \cdot \text{Long.MAX_VALUE}$).

In this case, two values play an essential role: the true mathematical value of

$$a * \text{randomNumbers}[\text{indexOfInteger}] + c$$

and the number you get in your program. The first one can be written as

$$\text{trueNumber} = \text{Long.MAX_VALUE} + \text{valueOverflow}$$

where valueOverflow is the size of the overflow got in the operation, whereas the number produced by Java is

$$\text{observedNumber} = \text{Long.MIN_VALUE} + \text{valueOverflow} - 1. \quad (1)$$

Note that, if one does not take the overflow into account, Java would simply return

$$\text{observedNumber} \% \text{modulus},$$

which might differ from

$$r := \text{trueNumber} \% \text{modulus}.$$

This is the all point: **if**

$$\text{observedNumber} \% \text{modulus} = \text{trueNumber} \% \text{modulus},$$

then adding the modulus to the value returned by Java would be fine. But in general, this is not the case!

The goal of the exercise is then to find a way to get r only looking at observedNumber .

By the generalized version of the distributive property of the $\%$ operation, we have

$$\begin{aligned} & \text{Long.MAX_VALUE} + \text{valueOverflow} \% \text{modulus} \\ &= ((\text{Long.MAX_VALUE} \% \text{modulus}) + (\text{valueOverflow} \% \text{modulus})) \% \text{modulus} \\ &= (\text{remainderOfMax} + \text{remainderOverflow}) \% \text{modulus}, \end{aligned}$$

where

$$\text{remainderOfMax} = \text{Long.MAX_VALUE} \% \text{modulus}$$

and

$$\text{remainderOverflow} = \text{valueOverflow} \% \text{modulus}.$$

Note that $\text{remainderOfMax} + \text{remainderOverflow}$ is positive and less than Long.MAX_VALUE since by assumption $2 \cdot \text{modulus} < \text{Long.MAX_VALUE}$, so it is not affected by overflows. This is thus the right correction to the overflow that we have to perform before applying $\%$.

Now it only remains to get valueOverflow from the observed number (1), i.e.

$$\text{valueOverflow} = \text{observedNumber} - \text{Long.MIN_VALUE} + 1.$$