

**Exercise 1**

Write a class implementing `AbstractAssetMonteCarloProduct` whose method

```
getValue(double evaluationTime, AssetModelMonteCarloSimulationModel
        underlyingSimulation)
```

returns the central finite difference approximation for a given shift  $\delta > 0$  of the derivative of the Monte-Carlo computation of the price of a European call option with underlying `underlyingSimulation`, with respect to the initial value of `underlyingSimulation` (i.e. the Delta of the option). That is, the  $i$ -th realization of the `RandomVariable` object returned by such a method must be the (discounted) value of

$$\frac{(S_T(i, S_0 + \delta) - K)^+ - (S_T(i, S_0 - \delta) - K)^+}{2\delta},$$

where  $S_T(i, S_0 + \delta)$  and  $S_T(i, S_0 - \delta)$  are the final values for the  $i$ -th simulation of an underlying which is identical to `underlyingSimulation` apart for the initial value, which is now  $S_0 + \delta$  and  $S_0 - \delta$ , respectively, instead of  $S_0$ .

**Hints:** the main point here is to get a clone of `underlyingSimulation` for changed initial value. You can do so by using the method `getCloneWithModifiedData` of `AssetModelMonteCarloSimulationModel`. This method takes a `Map<String, Object>` object specifying what you want to change and which new value you want to give.

**Exercise 2**

Write a class for the approximation of the derivative of the price of a European call option with respect to the initial price  $S_0$  of the asset (i.e., for the Delta of the option), when the underlying is a Black-Scholes model. In particular, the class should have:

- A method computing and returning the Delta of the option by computing the analytical prices of the option when the initial value of the asset is  $S_0 + \delta$  and  $S_0 - \delta$ , for a given  $\delta > 0$ .
- A method which creates an object of type `MonteCarloBlackScholesModel`, for number of simulations and time discretization given as arguments of the method, and then uses the implementation of the class of Exercise 1, for a given shift  $\delta > 0$ , to return a `double` representing the Monte-Carlo Delta of the option for such an underlying.
- A method that, using

```
net.finmath.functions.AnalyticFormulas.blackScholesOptionDelta
```

to have a benchmark, plots the errors given by the two methods above for different values of the shift  $\delta > 0$ .