

Sumcheck-free hypercube inner products from table-form Kronecker encodings

Research note (companion to Kronecker-FRI)

September 2026 — version 7 (Section 5: sparse matrix–vector products)

Abstract

We encode a multilinear polynomial by its *table*, $V_f(X) = \sum_w f(w)X^w$, instead of its coefficients. Two quantities then become single coefficients of univariate products: the evaluation $\tilde{f}(z)$, with a product-form kernel, and the hypercube inner product $\sum_w a(w)b(w)$ of two committed tables, through the reversal $V_b^*(X) = X^{N-1}V_b(1/X)$. Because a smooth domain is closed under inversion, the verifier reads V_b^* from the commitment to b as a virtual word, and the Kronecker-FRI machinery (opening polynomial, virtual word, one folding test) proves $\sum_w a(w)b(w) = S$ with no sumcheck, with soundness error $3M/|\mathbb{F}| + \varepsilon_{\text{fold}} + (1 - \delta)^\kappa$. With geometric weights γ^w and DEEP quotients as further virtual words, the same method gives a sumcheck-free Hadamard check $a \circ b = c$ for three committed tables, with soundness error $2(N - 1)/|\mathbb{F}| + 8M/|\mathbb{F}| + \varepsilon_{\text{fold}} + (1 - \delta)^\kappa$. Finally, any list of evaluations, inner products and Hadamard checks on the same commitments is proved by one folding test, with soundness error $2(N - 1)/|\mathbb{F}| + tM/|\mathbb{F}| + \varepsilon_{\text{fold}} + (1 - \delta)^\kappa$ for $t + 1$ batched words; the term $2(N - 1)/|\mathbb{F}|$ is paid once, whatever the number of Hadamard checks. The results of Sections 1–4 are machine-checked in Lean 4. Section 5 reduces sparse matrix–vector products, and hence R1CS, to the same statements with lookups by logarithmic derivatives, so an R1CS argument needs no sumcheck at all.

Setting. Notation and parameters are those of Kronecker-FRI (KF) [1]: $n \geq 1$, $N = 2^n$, a smooth domain $L \leq \mathbb{F}^\times$ of order $M = 2^{n+R}$ with $R \geq 2$, codes $\mathcal{C}_j$, the folding test $\Pi_{\text{FT}}[\ell, \kappa]$ with error $\varepsilon_{\text{fold}}$ (KF Thm 5.24), fibre distance Δ_0^{fib} , and $0 < \delta \leq \delta^* = (1 - \rho)/2$. A bit vector $w \in \{0, 1\}^n$ is identified with the integer $\sum_k w_k 2^{k-1} \in [0, N - 1]$.

1 Table encoding and two extraction identities

Definition 1 (Table encoding, reversal). For a table $f : \{0, 1\}^n \rightarrow \mathbb{F}$, $V_f(X) = \sum_w f(w)X^w \in \mathbb{F}[X]_{<N}$ and $\text{Enc}^T(f) = \text{ev}_L(V_f)$. For $P \in \mathbb{F}[X]_{<N}$, its reversal is $P^*(X) = X^{N-1}P(1/X)$, so $[X^j]P^* = [X^{N-1-j}]P$ and $(P^*)^* = P$.

$f \mapsto V_f$ is a linear bijection onto $\mathbb{F}[X]_{<N}$, so Enc^T is a linear bijection onto $\mathcal{C}_0$ (as KF Lemma 6.3(1)); it is the Kronecker encoding of the multilinear polynomial whose coefficients are the values of f , and needs no Möbius transform. In particular $\text{Enc}^T(f) = \text{Enc}(f)$ as maps on vectors of length N : a table-form commitment is an ordinary Kronecker-FRI commitment to the vector of values, with the same Merkle tree, and all of KF’s results on commitments (unique decoding, extraction, batching) apply to it unchanged.

Lemma 2 (Evaluation). For $z \in \mathbb{F}^n$ let $E_z^*(X) = \prod_{k=1}^n (z_k + (1 - z_k)X^{2^{k-1}})$. Then $\tilde{f}(z) = [X^{N-1}]V_f E_z^*$, where $\tilde{f}$ is the multilinear extension of f .

Proof. Expanding the product, $E_z(X) := \sum_w \text{eq}(w, z) X^w = \prod_k ((1 - z_k) + z_k X^{2^{k-1}})$, and $X^{N-1} E_z(1/X) = \prod_k X^{2^{k-1}} ((1 - z_k) + z_k X^{-2^{k-1}}) = E_z^*(X)$ since $N - 1 = \sum_k 2^{k-1}$. Hence $[X^{N-1-w}] E_z^* = [X^w] E_z = \text{eq}(w, z)$ and $[X^{N-1}] V_f E_z^* = \sum_w f(w) \text{eq}(w, z) = \tilde{f}(z)$. $\square$

Lemma 3 (Inner product). *For tables a, b : $\sum_w a(w)b(w) = [X^{N-1}] V_a V_b^*$.*

Proof. $[X^{N-1}] V_a V_b^* = \sum_{i=0}^{N-1} [X^i] V_a [X^{N-1-i}] V_b^* = \sum_i [X^i] V_a [X^i] V_b = \sum_w a(w)b(w)$. $\square$

Both are instances of one fact: for $U \in \mathbb{F}[X]_{<N}$ and a public $K \in \mathbb{F}[X]_{<N}$, $[X^{N-1}] UK = \sum_i [X^i] U [X^{N-1-i}] K$. KF's split lemma (Lemma 4.10) and identity lemma (Lemma 4.12) use only $\deg U, \deg K \leq N - 1$, so they hold verbatim for UK replaced by $V_a E_z^*$ or $V_a V_b^*$:

Lemma 4 (Split and identity, general form). *Let $U, K, A, H \in \mathbb{F}[X]_{<N}$ and $v \in \mathbb{F}$. (1) There are $A, H \in \mathbb{F}[X]_{<N}$ with $XUK = A + vX^N + X^{N+1}H$ iff $v = [X^{N-1}] UK$. (2) $\Phi = XUK - A - vX^N - X^{N+1}H$ has degree at most $2N$, and $\Phi = 0$ implies $v = [X^{N-1}] UK$.*

Proof. As in KF Lemmas 4.10 and 4.12: $\deg(UK) \leq 2N - 2$, and the three parts of the right-hand side occupy the disjoint degree ranges $[0, N - 1]$, $\{N\}$, $[N + 1, 2N]$; compare the coefficients of X^N . $\square$

Lemma 5 (Reversed words). *For $w \in \mathbb{F}^L$ put $w^*(\xi) = \xi^{N-1} w(\xi^{-1})$. Then: (1) $w = \text{ev}_L(P)$ with $P \in \mathbb{F}[X]_{<N}$ implies $w^* = \text{ev}_L(P^*)$; in particular $w \mapsto w^*$ maps $\mathcal{C}_0$ onto $\mathcal{C}_0$; (2) $w \mapsto w^*$ is a linear bijection of $\mathbb{F}^L$, and $w^* = c^*$ at ξ iff $w = c$ at ξ^{-1} ; (3) a set $T \subseteq L$ is fibre-closed iff $T^{-1} = \{\xi^{-1} : \xi \in T\}$ is, and $|T^{-1}| = |T|$; (4) $h(\xi) = w^*(\xi)$ is computed from the value of w at $\xi^{-1} \in L$, and a fold check at $\pm\xi$ reads w on the fibre $\{\pm\xi^{-1}\}$.*

Proof. L is a group, so $\xi \mapsto \xi^{-1}$ is a bijection of L mapping $-\xi$ to $-\xi^{-1}$ (as $-1 \in L$); and $\xi^{N-1} \neq 0$. (1) $\xi^{N-1} P(\xi^{-1}) = P^*(\xi)$, and $P \mapsto P^*$ is a bijection of $\mathbb{F}[X]_{<N}$. (2)–(4) follow from the first sentence. $\square$

2 The inner-product protocol

Definition 6 (Π_{IP}). Instance: oracles $w_a, w_b \in \mathbb{F}^L$ and $S \in \mathbb{F}$; honestly $w_a = \text{Enc}^T(a)$, $w_b = \text{Enc}^T(b)$ and $S = \sum_w a(w)b(w)$.

1. The prover sends $w_A = \text{ev}_L(A)$, where $XV_a V_b^* = A + SX^N + X^{N+1}H$ (Lemma 4); the verifier sends $\beta \in \mathbb{F}$.
2. Put the virtual words w_b^* (Lemma 5) and $h(\xi) = (\xi w_a(\xi) w_b^*(\xi) - w_A(\xi) - S\xi^N)/\xi^{N+1}$, and

$$w_0 = w_a + \beta w_b^* + \beta^2 w_A + \beta^3 h.$$

3. Run $\Pi_{\text{FT}}[\ell, \kappa]$ (or Π_{FT}^C , KF Prop. 5.28) on w_0 .

Per query point the verifier reads $w_a(\pm\xi)$, $w_b(\pm\xi^{-1})$ and $w_A(\pm\xi)$, and does $O(1)$ extra field operations beyond KF (no kernel to evaluate).

Theorem 7 (Completeness). *If $w_a = \text{ev}_L(V_a)$, $w_b = \text{ev}_L(V_b)$ and $S = \sum_w a(w)b(w)$, the honest prover is accepted with probability 1.*

Proof. By Lemma 3 and Lemma 4(1), $A, H \in \mathbb{F}[X]_{<N}$ exist; by Lemma 5(1) $w_b^* = \text{ev}_L(V_b^*)$; evaluating the identity at ξ gives $h = \text{ev}_L(H)$. So $w_0 = \text{ev}_L(V_a + \beta V_b^* + \beta^2 A + \beta^3 H)$ with a polynomial of degree $< N$, and KF Lemma 5.9(1) applies. $\square$

Definition 8 (Relation). $\mathcal{R}_{\text{IP}}^\delta$: $((w_a, w_b; S), (a, b))$ with $\Delta_0^{\text{fib}}(w_a, \text{Enc}^T(a)) \leq \delta$, $\Delta_0^{\text{fib}}(w_b, \text{Enc}^T(b)) \leq \delta$ and $\sum_w a(w)b(w) = S$.

Lemma 9 (Batching, degree 3). *Let $\mathcal{G}$ be the set of β for which w_0 agrees with some $c \in \mathcal{C}_0$ on a fibre-closed $T \subseteq L$ with $|T| \geq (1 - \delta)M$. If $|\mathcal{G}| > 3M$, then $(w_a, w_b; S)$ has a witness in $\mathcal{R}_{\text{IP}}^\delta$.*

Proof. The words w_0 form the curve of degree 3 generated by (w_a, w_b^*, w_A, h) , and every $\beta \in \mathcal{G}$ has $\Delta(w_0, \mathcal{C}_0) \leq \delta$. By correlated agreement for curves (KF Thm 2.16, from [2], with $t = 3$) there are $P_0, P_1, P_2, P_3 \in \mathbb{F}[X]_{<N}$ and a set Agr , $|\text{Agr}| \geq (1 - \delta)M$, on which $w_a = \text{ev}(P_0)$, $w_b^* = \text{ev}(P_1)$, $w_A = \text{ev}(P_2)$, $h = \text{ev}(P_3)$ (take Agr maximal).

Value. Let $\Phi = XP_0P_1 - P_2 - SX^N - X^{N+1}P_3$. On Agr , $\Phi(\xi) = \xi w_a(\xi)w_b^*(\xi) - w_A(\xi) - S\xi^N - \xi^{N+1}h(\xi) = 0$ by the definition of h . Since $\deg \Phi \leq 2N < (1 - \delta)M \leq |\text{Agr}|$ (KF Lemma 7.2, which uses $R \geq 2$), $\Phi = 0$, and Lemma 4(2) gives $S = [X^{N-1}]P_0P_1$. Let a, b be the tables with $V_a = P_0$ and $V_b = P_1^*$; then $V_b^* = P_1$ and $S = \sum_w a(w)b(w)$ by Lemma 3.

Bad challenges. For $\xi \notin \text{Agr}$, $d_\xi(Z) = \sum_{i=0}^3 Z^i(u_i - \text{ev}(P_i))(\xi)$, with $(u_0, \dots, u_3) = (w_a, w_b^*, w_A, h)$, is a nonzero polynomial of degree ≤ 3 ; its roots over all such ξ form a set Bad with $|\text{Bad}| \leq 3\delta M < 3M < |\mathcal{G}|$. Take $\beta \in \mathcal{G} \setminus \text{Bad}$ with its c and T , and $\hat{c} = \sum_i \beta^i \text{ev}(P_i) \in \mathcal{C}_0$. Then c and $\hat{c}$ agree with w_0 on $T \cap \text{Agr}$, of size $\geq (1 - 2\delta)M \geq N$, so $c = \hat{c}$; and for $\xi \in T$, $d_\xi(\beta) = w_0(\xi) - \hat{c}(\xi) = 0$ forces $\xi \in \text{Agr}$. So $T \subseteq \text{Agr}$.

Distances. $w_a = \text{ev}(V_a)$ on the fibre-closed T , so $\Delta_0^{\text{fib}}(w_a, \text{Enc}^T(a)) \leq \delta$ (KF Lemma 5.11(2)). $w_b^* = \text{ev}(V_b^*) = \text{ev}(V_b)^*$ on T , so by Lemma 5(2) $w_b = \text{ev}(V_b)$ on T^{-1} , which is fibre-closed of the same size (Lemma 5(3)); so $\Delta_0^{\text{fib}}(w_b, \text{Enc}^T(b)) \leq \delta$. $\square$

Theorem 10 (Soundness). *If $(w_a, w_b; S)$ has no witness in $\mathcal{R}_{\text{IP}}^\delta$, every prover is accepted with probability at most*

$$\frac{3M}{|\mathbb{F}|} + \varepsilon_{\text{fold}} + (1 - \delta)^\kappa < \frac{4M}{|\mathbb{F}|} + (1 - \delta)^\kappa.$$

Proof. Identical to KF Thm 7.8, with Lemma 9 in place of KF Lemma 7.4: for $\beta \notin \mathcal{G}$ (probability $\geq 1 - 3M/|\mathbb{F}|$) and good r (probability $\geq 1 - \varepsilon_{\text{fold}}$, KF Thm 5.24(1)), the codeword chain (KF Prop. 5.23) shows that $\text{dens}_\ell(w_\ell) \geq 1 - \delta$ would put β in $\mathcal{G}$, the witness set $\mathcal{W}_\ell(w_\ell)$ being fibre-closed (KF Lemma 7.7); so the query term is at most $(1 - \delta)^\kappa$ (KF Lemma 3.3). $\square$

Round-by-round knowledge soundness, the extractor (decode w_a and w_b at radius δ^*) and binding follow as in KF Thm 7.11 and Cor. 7.12, with round-1 error $3M/|\mathbb{F}|$.

Remark 11 (One commitment, three kinds of openings). The same table-form commitment $\text{Enc}^T(f)$ supports: evaluation openings at any z (KF's protocol with the kernel E_z^* of Lemma 2, evaluated by the verifier with $O(n)$ operations); hypercube sums ($z = (\frac{1}{2}, \dots, \frac{1}{2})$ up to 2^n , or the kernel $\prod_k (1 + X^{2^{k-1}})$ directly); and inner products with any other committed table (Π_{IP}). Inner products with a *public* table p are the special case $K = V_p^*$, useful when $V_p^*(\xi)$ is cheap to evaluate (product form). All of these batch with KF's batched openings (KF Thm 11.2).

3 A sumcheck-free Hadamard check

SNARKs of Spartan/HyperPlonk type need, beyond linear functionals and inner products, the statement $a(w)b(w) = c(w)$ for all w (a *Hadamard check*), for three committed tables.

Geometric weights. For $\gamma \in \mathbb{F}$,

$$\sum_w \gamma^w a(w)b(w) = [X^{N-1}] V_a(\gamma X) V_b^*(X), \quad \sum_w \gamma^w c(w) = V_c(\gamma),$$

the first by Lemma 3 applied to the tables $w \mapsto \gamma^w a(w)$ and b . If $d = a \circ b - c \neq 0$, then $D_d(X) = \sum_w d(w)X^w$ is a nonzero polynomial of degree $\leq N-1$, and the two sides differ unless $D_d(\gamma) = 0$. The values $V_a(\gamma\xi)$ and $V_c(\gamma)$ lie outside L ; the prover commits to $Q(X) = V_a(\gamma X)$, and both are tied to the commitments by DEEP quotients, which are virtual words like h .

Definition 12 (Quotient words). For $w \in \mathbb{F}^L$ and $y, x \in \mathbb{F}$, put $q[w, y, x](\xi) = (w(\xi) - y)/(\xi - x)$, with the value 0 at $\xi = x$ if $x \in L$.

Lemma 13 (DEEP step on an agreement set). *Let $x, y \in \mathbb{F}$, $P, R \in \mathbb{F}[X]_{<N}$ and $T \subseteq L$ with $|T| > N+1$. If $w = \text{ev}(P)$ and $q[w, y, x] = \text{ev}(R)$ on T , then $P(x) = y$ and $R = (P - y)/(X - x)$. Conversely, if $x \notin L$, $w = \text{ev}_L(P)$ and $P(x) = y$, then $q[w, y, x] = \text{ev}_L((P - y)/(X - x))$, a polynomial of degree $\leq N-2$.*

Proof. $(X - x)R - P + y$ has degree $\leq N$ and vanishes on $T \setminus \{x\}$, which has more than N points, so it is 0; evaluate at x . The converse is the factor theorem. $\square$

Definition 14 (Π_{Had}). Instance: oracles $w_a, w_b, w_c \in \mathbb{F}^L$; honestly $w_f = \text{Enc}^T(f)$ for $f \in \{a, b, c\}$ and $a \circ b = c$.

1. The verifier sends γ , uniform in $\mathbb{F}$. The prover sends $w_Q = \text{ev}_L(Q)$, where $Q(X) = V_a(\gamma X)$.
2. The verifier sends θ , uniform in $\mathbb{F}$. The prover sends $y_1 = V_a(\gamma\theta)$, $y_3 = V_c(\gamma)$ and $w_A = \text{ev}_L(A)$, where $XQV_b^* = A + y_3X^N + X^{N+1}H$ (Lemma 4).
3. The verifier sends β , uniform in $\mathbb{F}$. With $h(\xi) = (\xi w_Q(\xi) w_b^*(\xi) - w_A(\xi) - y_3\xi^N)/\xi^{N+1}$ and

$$(u_0, \dots, u_8) = (w_a, w_c, w_Q, w_b^*, w_A, h, q[w_a, y_1, \gamma\theta], q[w_Q, y_1, \theta], q[w_c, y_3, \gamma]),$$

put $w_0 = \sum_{i=0}^8 \beta^i u_i$.

4. Run $\Pi_{\text{FT}}[\ell, \kappa]$ (or Π_{FT}^C) on w_0 .

The challenges are unrestricted: soundness holds for all values, and a pole $\gamma\theta, \theta, \gamma$ in L affects only completeness. Using y_1 for both $V_a(\gamma\theta)$ and $Q(\theta)$ is the consistency check $V_a(\gamma\theta) = Q(\theta)$. Per query point the verifier reads w_a, w_c, w_Q, w_A on the coset of ξ and w_b on the coset of ξ^{-1} , and does $O(1)$ field operations and three inversions per point (one batched inversion per query) beyond the folding test.

Theorem 15 (Completeness). *If $w_f = \text{Enc}^T(f)$ for $f \in \{a, b, c\}$ and $a \circ b = c$, the honest prover is accepted whenever $\gamma\theta, \theta, \gamma \notin L$, hence with probability at least $1 - 3M/|\mathbb{F}|$.*

Proof. $Q \in \mathbb{F}[X]_{<N}$; $y_1 = V_a(\gamma\theta) = Q(\theta)$; and $y_3 = V_c(\gamma) = \sum_w \gamma^w c(w) = \sum_w \gamma^w a(w)b(w) = [X^{N-1}] QV_b^*$, so A, H exist and $h = \text{ev}_L(H)$ as in Π_{IP} . By Lemma 13 the three quotient words are codewords when the poles lie outside L , and then w_0 is the codeword of a polynomial of degree $< N$. Each of the events $\gamma \in L$, $\theta \in L$, $\gamma\theta \in L$ has probability at most $M/|\mathbb{F}|$ (for $\gamma \neq 0$, at most M values of θ put $\gamma\theta$ in L ; for $\gamma = 0$, none). $\square$

Definition 16 (Relation). $\mathcal{R}_{\text{Had}}^\delta$: $((w_a, w_b, w_c), (a, b, c))$ with $\Delta_0^{\text{fib}}(w_f, \text{Enc}^T(f)) \leq \delta$ for $f \in \{a, b, c\}$ and $a \circ b = c$.

Lemma 17 (Batching, degree 8). *Fix $\gamma, \theta \in \mathbb{F}$ and the prover's w_Q, y_1, y_3, w_A . Let $\mathcal{G}$ be the set of β for which w_0 agrees with some $c' \in \mathcal{C}_0$ on a fibre-closed $T \subseteq L$ with $|T| \geq (1 - \delta)M$. If $|\mathcal{G}| > 8M$, there are tables a, b, c and $\hat{Q} \in \mathbb{F}[X]_{<N}$ with*

1. $\Delta_0^{\text{fib}}(w_f, \text{Enc}^T(f)) \leq \delta$ for $f \in \{a, b, c\}$, and $\Delta_0^{\text{fib}}(w_Q, \text{ev}_L(\hat{Q})) \leq \delta$;
2. $V_a(\gamma\theta) = y_1 = \hat{Q}(\theta)$;
3. $V_c(\gamma) = y_3 = [X^{N-1}] \hat{Q} V_b^*$.

Proof. Correlated agreement on one fibre-closed set (the step shared with Lemma 9; Lean: `curve_agreement`, $t = 8$) gives $P_0, \dots, P_8 \in \mathbb{F}[X]_{<N}$ and a fibre-closed T , $|T| \geq (1 - \delta)M > 2N \geq N + 2$ (KF Lemma 7.2), with $u_i = \text{ev}(P_i)$ on T . As in Lemma 9, $XP_2P_3 - P_4 - y_3X^N - X^{N+1}P_5$ vanishes on T and has degree $\leq 2N$, so $y_3 = [X^{N-1}] P_2P_3$. Lemma 13 with $(P_0, P_6, \gamma\theta)$, (P_2, P_7, θ) and (P_1, P_8, γ) gives $P_0(\gamma\theta) = y_1$, $P_2(\theta) = y_1$ and $P_1(\gamma) = y_3$. Let $V_a = P_0$, $V_c = P_1$, $V_b = P_3^*$ and $\hat{Q} = P_2$; then (2) and (3) hold, and (1) follows from agreement on T (for w_b , on T^{-1} , Lemma 5). $\square$

Theorem 18 (Soundness). *Let $R \geq 2$ and $0 < \delta \leq \delta^*$. If (w_a, w_b, w_c) has no witness in $\mathcal{R}_{\text{Had}}^\delta$, every prover is accepted with probability at most*

$$\varepsilon_{\text{Had}} = \frac{2(N-1)}{|\mathbb{F}|} + \frac{8M}{|\mathbb{F}|} + \varepsilon_{\text{fold}} + (1 - \delta)^\kappa.$$

Proof. For $\delta \leq \delta^*$ each word has at most one codeword within fibre distance δ (KF Lemma 6.3(2)), so the tables and $\hat{Q}$ of Lemma 17 are the decodings of w_a, w_b, w_c, w_Q . It suffices to consider deterministic provers (a randomised prover is an average of deterministic ones), so w_Q is a function of γ and the round-2 messages are functions of (γ, θ) . Call (γ, θ) *good* if, for every choice of the prover's round-2 messages, $|\mathcal{G}| \leq 8M$. For good (γ, θ) the argument of KF Thm 7.8 applied to the rest of the protocol bounds the acceptance probability by $8M/|\mathbb{F}| + \varepsilon_{\text{fold}} + (1 - \delta)^\kappa$. It remains to bound the probability that (γ, θ) is not good.

Case 1: some w_f has no table within fibre distance δ . Lemma 17(1) fails, so every (γ, θ) is good.

Case 2: w_a, w_b, w_c decode to $\hat{a}, \hat{b}, \hat{c}$ with $\hat{a} \circ \hat{b} \neq \hat{c}$. Let $D = D_{\hat{a} \circ \hat{b} - \hat{c}} \neq 0$. The event $D(\gamma) = 0$ has probability at most $(N-1)/|\mathbb{F}|$. Fix γ with $D(\gamma) \neq 0$, and the prover's w_Q (a function of γ). If w_Q has no codeword within fibre distance δ , (γ, θ) is good for every θ by Lemma 17(1). If it decodes to $\hat{Q} = V_{\hat{a}}(\gamma X)$, then Lemma 17(3) would give $\sum_w \gamma^w \hat{c}(w) = V_{\hat{c}}(\gamma) = [X^{N-1}] V_{\hat{a}}(\gamma X) V_{\hat{b}}^* = \sum_w \gamma^w \hat{a}(w) \hat{b}(w)$, i.e. $D(\gamma) = 0$; so every θ is good. Otherwise $\hat{Q} - V_{\hat{a}}(\gamma X)$ is a nonzero polynomial of degree $\leq N-1$, fixed before θ is drawn; the event $\hat{Q}(\theta) = V_{\hat{a}}(\gamma\theta)$ has probability at most $(N-1)/|\mathbb{F}|$, and outside it Lemma 17(2) fails, so θ is good. $\square$

So $\varepsilon_{\text{Had}} < (8M + 2N)/|\mathbb{F}| + \varepsilon_{\text{fold}} + (1 - \delta)^\kappa$, against $3M/|\mathbb{F}|$ for Π_{IP} and $2M/|\mathbb{F}|$ for an evaluation. The curve has degree 8 because w_a and w_c must be in the batch themselves: a quotient word alone determines w_a on T only as a polynomial of degree $\leq N$, which does not identify the decoding.

Knowledge follows: if the acceptance probability exceeds ε_{Had} , the instance has a witness, and the extractor decodes w_a, w_b, w_c at radius δ (KF Cor. 7.12).

Remark 19 (Consequence). Linear functionals (evaluations, with the kernel E_z^*), inner products (Π_{IP}) and Hadamard checks (Π_{Had}) are all proved on table-form Kronecker-FRI commitments by one folding test each, with no sumcheck and a verifier that is an FRI verifier plus $O(1)$ operations per query point (plus $O(n)$ for E_z^*). For R1CS-type SNARKs what remains is the sparse matrix-vector products Az, Bz, Cz , which Spartan handles with its own sparse-polynomial commitment; whether those admit a coefficient-extraction form is the next question.

4 One folding test for a batch of statements

A SNARK needs many statements at once on the same commitments: Hadamard checks, inner products and evaluations. Since every statement of Sections 1–3 is checked by words in one curve, all of them can share one curve and one folding test.

Definition 20 (Statements). Let $w_1, \dots, w_r \in \mathbb{F}^L$ be commitments. A *statement* is one of the following, with indices in $\{1, \dots, r\}$:

- **Eval**(i, z, v) with $z \in \mathbb{F}^n, v \in \mathbb{F}$: the table f_i of w_i has $\tilde{f}_i(z) = v$;
- **IP**(i, i', S) with $S \in \mathbb{F}$: $\sum_w f_i(w) f_{i'}(w) = S$;
- **Had**(i, i', k): $f_i \circ f_{i'} = f_k$.

For a list $\Sigma = (\sigma_1, \dots, \sigma_J)$ of $J \geq 1$ statements, let D be the set of commitment indices read directly (i in **Eval** and **IP**; i, k in **Had**), B the set of indices read reversed (i' in **IP** and **Had**), and $\mathcal{H} \subseteq \{1, \dots, J\}$ the positions of the Hadamard checks. A table may play several roles. The relation $\mathcal{R}_{\text{Batch}}^\delta$ holds for $((w_i)_i, \Sigma)$ and tables $(f_i)_{i \in D \cup B}$ if $\Delta_0^{\text{fib}}(w_i, \text{Enc}^T(f_i)) \leq \delta$ for all $i \in D \cup B$ and every statement of Σ holds for these tables.

For each statement σ_j the protocol uses a triple (U_j, K_j, v_j) , with $U_j, K_j \in \mathbb{F}[X]_{<N}$, and the identity $XU_jK_j = A_j + v_jX^N + X^{N+1}H_j$ of Lemma 4: (V_{f_i}, E_z^*, v) for **Eval**(i, z, v) (Lemma 2); $(V_{f_i}, V_{f_{i'}}^*, S)$ for **IP**(i, i', S) (Lemma 3); $(Q_j, V_{f_{i'}}^*, y_3^{(j)})$ with $Q_j(X) = V_{f_i}(\gamma X)$ for $\sigma_j = \text{Had}(i, i', k)$ (Section 3). The verifier reads $U_j(\xi)$ and $K_j(\xi)$ as $w_i(\xi)$ and $E_z^*(\xi)$, as $w_i(\xi)$ and $w_{i'}^*(\xi)$, or as $w_Q^{(j)}(\xi)$ and $w_{i'}^*(\xi)$.

Definition 21 (Π_{Batch}). Instance: $(w_i)_i$ and Σ .

1. The verifier sends γ , uniform in $\mathbb{F}$. For each $j \in \mathcal{H}$ the prover sends $w_Q^{(j)} = \text{ev}_L(Q_j)$.
2. The verifier sends θ , uniform in $\mathbb{F}$. For each $j \in \mathcal{H}$, with $\sigma_j = \text{Had}(i, i', k)$, the prover sends $y_1^{(j)} = V_{f_i}(\gamma\theta)$ and $y_3^{(j)} = V_{f_k}(\gamma)$; for each $j \leq J$ it sends $w_A^{(j)} = \text{ev}_L(A_j)$.
3. The verifier sends β , uniform in $\mathbb{F}$. The words are, in a fixed order: w_i for $i \in D$; w_i^* for $i \in B$; $w_Q^{(j)}$ for $j \in \mathcal{H}$; $w_A^{(j)}$ and $h_j(\xi) = (\xi U_j(\xi)K_j(\xi) - w_A^{(j)}(\xi) - v_j\xi^N)/\xi^{N+1}$ for $j \leq J$; and $q[w_i, y_1^{(j)}, \gamma\theta]$, $q[w_Q^{(j)}, y_1^{(j)}, \theta]$, $q[w_k, y_3^{(j)}, \gamma]$ for $j \in \mathcal{H}$, $\sigma_j = \text{Had}(i, i', k)$. They are $t+1 = |D| + |B| + 2J + 4|\mathcal{H}|$ words $u_0, \dots, u_t$; put $w_0 = \sum_{s=0}^t \beta^s u_s$.
4. Run $\Pi_{\text{FT}}[\ell, \kappa]$ (or Π_{FT}^C) on w_0 .

γ and θ are shared by all Hadamard checks. All $w_Q^{(j)}$ can be committed in one Merkle tree whose leaves hold one value of each, and likewise all $w_A^{(j)}$; a query then opens, beyond the folding test, one path per commitment in D , one per commitment in B (on the inverse coset), and at most two paths for the prover's oracles (one if $\mathcal{H} = \emptyset$), whatever J .

Theorem 22 (Completeness). *If $w_i = \text{Enc}^T(f_i)$ and every statement holds, the honest prover is accepted whenever $\mathcal{H} = \emptyset$ or $\gamma\theta, \theta, \gamma \notin L$; so with probability at least $1 - 3M/|\mathbb{F}|$.*

Proof. Each $w_A^{(j)}$, h_j and quotient word is a codeword, by the completeness proofs of Sections 2 and 3 and Lemma 2 for **Eval** (where E_z^* is a public polynomial of degree $< N$). $\square$

Lemma 23 (Batching, degree t). *Fix γ, θ and the prover's messages, and let $\mathcal{G}$ be the set of β for which w_0 agrees with some $c' \in \mathcal{C}_0$ on a fibre-closed $T \subseteq L$ with $|T| \geq (1 - \delta)M$. Let $R \geq 2$ and $0 < \delta \leq \delta^*$. If $|\mathcal{G}| > tM$, there are tables $(f_i)_{i \in D \cup B}$ and polynomials $\hat{Q}_j \in \mathbb{F}[X]_{<N}$ ($j \in \mathcal{H}$) such that $\Delta_0^{\text{fib}}(w_i, \text{Enc}^T(f_i)) \leq \delta$, $\Delta_0^{\text{fib}}(w_Q^{(j)}, \text{ev}_L(\hat{Q}_j)) \leq \delta$, and*

1. *every Eval and IP statement of Σ holds for the f_i ;*
2. *for every $j \in \mathcal{H}$ with $\sigma_j = \text{Had}(i, i', k)$: $V_{f_i}(\gamma\theta) = y_1^{(j)} = \hat{Q}_j(\theta)$ and $V_{f_k}(\gamma) = y_3^{(j)} = [X^{N-1}] \hat{Q}_j V_{f_{i'}}^*$.*

Proof. Correlated agreement on one fibre-closed set (**curve_agreement** with this t) gives polynomials of degree $< N$ for all the words and a fibre-closed T with $|T| \geq (1 - \delta)M > 2N$ on which all

words agree with them. Tables: V_{f_i} is the polynomial of w_i for $i \in D$; for $i \in B$, V_{f_i} is the reversal of the polynomial of w_i^* , and w_i agrees with $\text{Enc}^T(f_i)$ on T^{-1} (Lemma 5). If $i \in D \cap B$, both give tables within fibre distance δ of w_i , which coincide by unique decoding. For each j , the identity of h_j on T and Lemma 4(2) give $v_j = [X^{N-1}]U_jK_j$ with the polynomials of the words; for **Eval** this is $v = f_i(z)$ by Lemma 2, for **IP** it is $S = \sum_w f_i(w)f_{i'}(w)$ by Lemma 3, and for **Had** it is the second equality of (2) with $\hat{Q}_j$ the polynomial of $w_Q^{(j)}$. The three quotient words of each $j \in \mathcal{H}$ give the other equalities of (2) by Lemma 13, as in Lemma 17. $\square$

Theorem 24 (Soundness). *Let $R \geq 2$ and $0 < \delta \leq \delta^*$. If $((w_i)_i, \Sigma)$ has no witness in $\mathcal{R}_{\text{Batch}}^\delta$, every prover is accepted with probability at most*

$$\varepsilon_{\text{Batch}} = \frac{2(N-1)}{|\mathbb{F}|} + \frac{tM}{|\mathbb{F}|} + \varepsilon_{\text{fold}} + (1-\delta)^\kappa.$$

Proof. As for Theorem 18: call (γ, θ) good if $|\mathcal{G}| \leq tM$ for all round-2 messages; for good (γ, θ) the acceptance probability is at most $tM/|\mathbb{F}| + \varepsilon_{\text{fold}} + (1-\delta)^\kappa$. If some w_i , $i \in D \cup B$, has no table within fibre distance δ , every pair is good by Lemma 23. Otherwise let $\hat{f}_i$ be the decodings; some statement σ_{j^*} is false for them, and j^* depends only on the instance. If σ_{j^*} is an **Eval** or **IP** statement, Lemma 23(1) fails, so every pair is good. If $\sigma_{j^*} = \text{Had}(i, i', k)$, the proof of Theorem 18 applies verbatim to this one check, with Lemma 23(2) for j^* in place of Lemma 17: the pair is good unless $D_{\hat{f}_i \circ \hat{f}_{i'} - \hat{f}_k}(\gamma) = 0$, or $\hat{Q}_{j^*} \neq V_{\hat{f}_i}(\gamma X)$ and $\hat{Q}_{j^*}(\theta) = V_{\hat{f}_i}(\gamma \theta)$, events of probability at most $(N-1)/|\mathbb{F}|$ each. $\square$

When $\mathcal{H} = \emptyset$ the same proof shows that every pair is good, so the first term can be omitted (this refinement is not in the Lean formalisation). Only one false statement is needed in the proof, so the term $2(N-1)/|\mathbb{F}|$ from γ, θ is paid once, not once per Hadamard check; the term $tM/|\mathbb{F}|$ grows linearly in the number of words, by $4M/|\mathbb{F}|$ per Hadamard check and $2M/|\mathbb{F}|$ per other statement. For $|\mathbb{F}| = p^2 \approx 2^{128}$, $M = 2^{22}$ and $t = 100$, this term is about 2^{-99} . Knowledge follows as before: if the acceptance probability exceeds $\varepsilon_{\text{Batch}}$, the instance has a witness, obtained by decoding each w_i .

n	one batched proof ($t+1=16$)			four separate proofs		
	prove (ms)	verify (ms)	proof (KiB)	prove (ms)	verify (ms)	proof (KiB)
12	38.2	7.21	397.9	54.3	13.85	795.4
14	165.2	8.20	479.5	224.9	16.17	1017.2
16	716.2	8.90	587.5	959.8	18.70	1315.3
18	3154.6	10.22	701.2	4116.7	22.59	1658.7
20	13187.4	11.32	804.5	18012.6	25.43	1957.9

Table 1: The statements **Had**(a, b, c), **IP**(a, d, S), **Eval**(a, z_1, v_1), **Eval**(c, z_2, v_2) on four table-form commitments, proved by one Π_{Batch} proof or by four (each a Π_{Batch} proof of one statement). Configuration KF-8, 148 queries, 32-byte salts, $\mathbb{F}_{p^2}$; one core; medians. Source: `bench/batch.csv`, `cargo run --release --example batch_bench`. The prototype prover is not optimised.

One proof for the four statements is 2.0 to 2.4 times smaller and 1.9 to 2.2 times faster to verify than four proofs (2.4 and 2.2 at $n = 20$), and its prover takes 23 to 30% less time.

5 Sparse matrix–vector products without sumcheck

An R1CS instance asks for z with $(Az) \circ (Bz) = Cz$ for public sparse matrices A, B, C . Committing to z and to $a = Az$, $b = Bz$, $c = Cz$, the statement $a \circ b = c$ is a Hadamard check. This section

proves $a = \mathbf{M}z$ for a public sparse matrix $\mathbf{M}$ (a *lincheck*) with the statements of Section 4 on affine forms of committed and public tables; lookups are handled with logarithmic derivatives (LogUp, Haböck [4]). Throughout, N is at least the length of z and the number of nonzero entries of each matrix, and all tables are padded to length $N = 2^n$; $R \geq 2$, $0 < \delta \leq \delta^*$ and $\text{char } \mathbb{F} > N$.

5.1 Affine forms and public tables

A *public table* is a table P whose polynomial V_P (and hence V_P^*) the verifier evaluates at any point with few operations. We use

$$\mathbf{1}(w) = 1, \quad \mathbf{0}(w) = 0, \quad \text{id}(w) = w, \quad G_\eta(w) = \eta^w \ (0^0 = 1),$$

with w read as an integer in $[0, N-1]$ and then as an element of $\mathbb{F}$, for which $V_1(X) = \prod_{k < n} (1 + X^{2^k}) = V_1^*(X)$, $V_{G_\eta}(X) = V_1(\eta X)$, $V_{G_\eta^*}(X) = \prod_{k < n} (X^{2^k} + \eta^{2^k})$, and $V_{\text{id}} = T_n$ with $S_0 = 1$, $T_0 = 0$, $S_{k+1} = (1 + X^{2^k})S_k$, $T_{k+1} = (1 + X^{2^k})T_k + 2^k X^{2^k} S_k$; each costs $O(n)$ operations. An *affine form* is $\lambda = \sum_i c_i w_i + \sum_j d_j P_j$ with commitments w_i , public tables P_j and coefficients $c_i, d_j \in \mathbb{F}$. Its word at ξ is $\sum_i c_i w_i(\xi) + \sum_j d_j V_{P_j}(\xi)$, and for tables f_i of the w_i its table is $\sum_i c_i f_i + \sum_j d_j P_j$.

Proposition 25 (Π_{Batch} on affine forms). *Let the arguments of the statements of Section 4 be affine forms, the coefficients and the parameters of the public tables being part of the instance.*

- **Relation:** *tables are required for the committed words that occur with a nonzero coefficient in some argument, within fibre distance δ ; every statement holds for the tables of its affine forms. Public tables carry no condition.*
- **Curve:** *the committed words occurring directly, the reversals of those occurring reversed, every $w_Q^{(j)} = \text{ev}_L(V_\lambda(\gamma X))$ (public part included), every $w_A^{(j)}$ and h_j , and the quotient words $q[\lambda, y, x]$ of affine forms λ with a committed part. When the form of a quotient word is purely public, the verifier checks $y = V_\lambda(x)$ directly instead.*

Then completeness, Lemma 23 and Theorem 24 hold with $t+1$ the number of words in the curve. The verifier's work per query point grows by $O(n)$ per public table.

Proof. On the agreement set T of Lemma 23, the word of an affine form agrees with the polynomial $\sum_i c_i P_{w_i} + \sum_j d_j V_{P_j}$ of degree $< N$, since public tables agree with their polynomials everywhere. The proofs of Lemma 23 and Theorem 24 use only such agreements, Lemmas 4 and 13, and the decodings of the committed words, which determine the tables of the affine forms; the case analysis of Theorem 18 uses only the polynomial V_λ of the decoded form. A direct check $y = V_\lambda(x)$ gives the conclusion of Lemma 13 for a public form. $\square$

5.2 The lincheck

Definition 26 (Index, relation). A sparse $\mathbf{M} \in \mathbb{F}^{N \times N}$ with $\nu \leq N$ nonzero entries is listed as $(r_k, c_k, v_k)_{k < N}$, padded with $v_k = 0$, $r_k = c_k = 0$. Its *index* consists of the commitments Enc^T of the tables $\text{row}(k) = r_k$, $\text{col}(k) = c_k$, $\text{val}(k) = v_k$ and of the multiplicities $m_R(w) = \#\{k : r_k = w\}$, $m_C(w) = \#\{k : c_k = w\}$, computed once by an honest indexer (soundness uses only that $\text{row}, \text{col}, \text{val}$ are honest). The lincheck relation holds for commitments (w_z, w_a) and tables (z, a) if $\Delta_0^{\text{fib}}(w_z, \text{Enc}^T(z)) \leq \delta$, $\Delta_0^{\text{fib}}(w_a, \text{Enc}^T(a)) \leq \delta$ and $a = \mathbf{M}z$.

The reduction rests on $\sum_w \eta^w (\mathbf{M}z)(w) = \sum_k v_k \eta^{r_k} z(c_k)$: the prover commits to $e_k = \eta^{r_k}$, $\zeta_k = z(c_k)$ and $p_k = e_k \zeta_k$, and proves the lookups $(r_k, e_k) \in \{(w, \eta^w)\}$ and $(c_k, \zeta_k) \in \{(w, z(w))\}$ by LogUp.

Definition 27 (Π_{Lin}). Instance: the index of $\mathbf{M}$ and commitments w_z, w_a .

1. The verifier sends η , uniform in $\mathbb{F}$. The prover commits to tables e, ζ, p (honestly $e_k = \eta^{r_k}$, $\zeta_k = z(c_k)$, $p_k = e_k \zeta_k$).
2. The verifier sends x_R, y_R, x_C, y_C , uniform in $\mathbb{F}$. The prover commits to tables $\phi^R, \psi^R, \phi^C, \psi^C$ and sends $s, s_R, s_C \in \mathbb{F}$; honestly

$$\begin{aligned}\phi_k^R &= \frac{1}{x_R - r_k - y_R e_k}, & \psi_w^R &= \frac{m_R(w)}{x_R - w - y_R \eta^w}, \\ \phi_k^C &= \frac{1}{x_C - c_k - y_C \zeta_k}, & \psi_w^C &= \frac{m_C(w)}{x_C - w - y_C z(w)},\end{aligned}$$

with $\psi_w = 0$ when $m(w) = 0$; the honest prover aborts if a denominator with $m(w) \neq 0$ vanishes.

3. The statements, proved by one Π_{Batch} on affine forms (Proposition 25) whose own challenges γ, θ, β are drawn after step 2:

$$\begin{aligned}& \text{IP}(a, G_\eta, s), \quad \text{IP}(\text{val}, p, s), \quad \text{Had}(e, \zeta, p), \\ & \text{Had}(\phi^R, x_R \mathbf{1} - \text{row} - y_R e, \mathbf{1}), \quad \text{Had}(\psi^R, x_R \mathbf{1} - \text{id} - y_R G_\eta, m_R), \\ & \text{IP}(\phi^R, \mathbf{1}, s_R), \quad \text{IP}(\psi^R, \mathbf{1}, s_R), \\ & \text{Had}(\phi^C, x_C \mathbf{1} - \text{col} - y_C \zeta, \mathbf{1}), \quad \text{Had}(\psi^C, x_C \mathbf{1} - \text{id} - y_C z, m_C), \\ & \text{IP}(\phi^C, \mathbf{1}, s_C), \quad \text{IP}(\psi^C, \mathbf{1}, s_C).\end{aligned}$$

The honest prover aborts with probability at most $2N/|\mathbb{F}|$ (a denominator $x - w - y \cdot$ with $m(w) \neq 0$ vanishes; the denominators of ϕ are among them), and is otherwise accepted with the probability of Proposition 25, at least $1 - 3M/|\mathbb{F}|$.

Lemma 28 (LogUp with pairs). *Let $\text{char } \mathbb{F} > N$, pairs $(\alpha_k, \alpha'_k)_{k < N}$ and $(\omega_w, \omega'_w)_{w < N}$ in $\mathbb{F}^2$, and $m_w \in \mathbb{F}$. If some (α_k, α'_k) is not among the (ω_w, ω'_w) , then for x, y uniform in $\mathbb{F}$,*

$$\Pr \left[\sum_k \frac{1}{x - \alpha_k - y \alpha'_k} = \sum_w \frac{m_w}{x - \omega_w - y \omega'_w} \text{ with all denominators nonzero} \right] \leq \frac{2N-1}{|\mathbb{F}|}.$$

Proof. In $\mathbb{F}(X, Y)$ let $\Phi = \sum_k (X - \alpha_k - Y \alpha'_k)^{-1} - \sum_w m_w (X - \omega_w - Y \omega'_w)^{-1}$, and let Den be the product of the $2N$ linear forms. Two such linear forms are equal iff their pairs are. Let $\ell = X - \alpha_k - Y \alpha'_k$ with (α_k, α'_k) not a table pair, and $c = \#\{k' : (\alpha_{k'}, \alpha'_{k'}) = (\alpha_k, \alpha'_k)\} \in [1, N]$. Multiply Φ by the product of the linear forms different from ℓ and substitute $X = \alpha_k + Y \alpha'_k$: every term vanishes except c times a product of nonzero polynomials in Y , so $\Phi \neq 0$ since $c \neq 0$ in $\mathbb{F}$. Hence ΦDen is a nonzero polynomial of total degree $\leq 2N - 1$; where $\text{Den} \neq 0$, $\Phi = 0$ iff $\Phi \text{Den} = 0$, and the Schwartz-Zippel lemma concludes (the pairs and m are fixed before x, y). $\square$

Theorem 29 (Lincheck reduction). *Suppose the index is honest, the decodings z, a of w_z, w_a exist and $a \neq \mathbf{M}z$, and fix a deterministic prover. Let E be the event, over η, x_R, y_R, x_C, y_C , that every word committed by the prover in steps 1–2 has a decoding within fibre distance δ and that these decodings, with z, a and the index, satisfy all the statements of step 3. Then*

$$\Pr[E] \leq \varepsilon_{\text{Lin}} = \frac{N-1}{|\mathbb{F}|} + 2 \left(\frac{2N-1}{|\mathbb{F}|} + \frac{N}{|\mathbb{F}|} \right) = \frac{7N-3}{|\mathbb{F}|}.$$

Proof. On E : from $\text{Had}(\phi^R, x_R \mathbf{1} - \text{row} - y_R e, \mathbf{1})$, every $x_R - r_k - y_R e_k$ is nonzero and ϕ_k^R is its inverse. From $\text{Had}(\psi^R, \dots, m_R)$, $\psi_w^R(x_R - w - y_R \eta^w) = m_R(w)$; unless $x_R = w + y_R \eta^w$ for some

w (probability $\leq N/|\mathbb{F}|$ over x_R , for every y_R), $\psi_w^R = m_R(w)/(x_R - w - y_R\eta^w)$. The two IP with **1** give $\sum_k \phi_k^R = \sum_w \psi_w^R$. The decoding e is fixed before x_R, y_R , so by Lemma 28, except with probability $(2N-1)/|\mathbb{F}|$, every (r_k, e_k) is a pair (w, η^w) , that is $e_k = \eta^{r_k}$. In the same way, except with probability $(2N-1)/|\mathbb{F}| + N/|\mathbb{F}|$, $\zeta_k = z(c_k)$ for all k (the decodings ζ and z are fixed before x_C, y_C). Then $\text{Had}(e, \zeta, p)$ gives $p_k = \eta^{r_k} z(c_k)$, and the first two statements give $\sum_w \eta^w a(w) = s = \sum_k v_k \eta^{r_k} z(c_k) = \sum_w \eta^w (\mathbf{M}z)(w)$. So η is a root of the nonzero polynomial $\sum_w (a - \mathbf{M}z)(w) X^w$ of degree $\leq N-1$, fixed before η : probability $\leq (N-1)/|\mathbb{F}|$. $\square$

Theorem 30 (Soundness of the lincheck). *If the index is honest and (w_z, w_a) has no witness for the lincheck relation, every prover of Π_{Lin} followed by Π_{Batch} is accepted with probability at most $\varepsilon_{\text{Lin}} + \varepsilon_{\text{Batch}}$, with $\varepsilon_{\text{Batch}}$ of Theorem 24 for the $t+1$ words of step 3.*

Proof. It suffices to consider deterministic provers. Let τ be the transcript of steps 1–2; the batch instance $I(\tau)$ and the event E of Theorem 29 are functions of τ , and given τ the rest of the prover is a prover of Π_{Batch} facing fresh challenges. If w_z or w_a has no decoding, $I(\tau)$ has no witness for every τ . Otherwise $a \neq \mathbf{M}z$, and for $\tau \notin E$, $I(\tau)$ has no witness: by unique decoding a witness would consist of the decodings, which either do not exist or violate a statement. In both cases Theorem 24 gives $\Pr[\text{accept} \mid \tau] \leq \varepsilon_{\text{Batch}}$ for $\tau \notin E$, so the acceptance probability is at most $\Pr[E] + \varepsilon_{\text{Batch}}$. $\square$

5.3 R1CS

Definition 31 (R1CS relation). Public: sparse $A, B, C \in \mathbb{F}^{N \times N}$ (with honest indices), input positions $I \subseteq [0, N-1]$ and an input $x \in \mathbb{F}^I$. Let $\hat{x}$ be the table equal to x on I and 0 elsewhere, and χ the 0/1 table of I (both public tables; for $I = [0, |I|-1]$, $V_\chi(X) = (X^{|I|} - 1)/(X - 1)$, and in general $V_{\hat{x}}, V_\chi$ cost $O(|I|)$). The relation holds for commitments (w_w, w_a, w_b, w_c) and tables (w, a, b, c) if all four are within fibre distance δ , w vanishes on I , and with $z = \hat{x} + w$: $a = Az$, $b = Bz$, $c = Cz$ and $a \circ b = c$.

Theorem 32 (R1CS without sumcheck). *Run the three linchecks Π_{Lin} for (A, a) , (B, b) , (C, c) on the affine form $z = \hat{x} + w_w$ (they may share their challenges), and prove their statements, $\text{Had}(a, b, c)$ and $\text{Had}(w, \chi, \mathbf{0})$ in one Π_{Batch} on affine forms. If the instance has no witness, every prover is accepted with probability at most*

$$\varepsilon_{\text{Lin}} + \varepsilon_{\text{Batch}} \leq \frac{9N - 5 + tM}{|\mathbb{F}|} + \varepsilon_{\text{fold}} + (1 - \delta)^\kappa.$$

Proof. As in Theorem 30, with one false statement: if some decoding is missing, the batch instance never has a witness; if $a \neq Az$ (or $b \neq Bz$, $c \neq Cz$) for the decodings, only the event E of that lincheck matters, of probability $\leq \varepsilon_{\text{Lin}}$; otherwise $a \circ b \neq c$ or w does not vanish on I , and $\text{Had}(a, b, c)$ or $\text{Had}(w, \chi, \mathbf{0})$ is false for every value of the challenges. $\square$

Every step is a coefficient extraction, a lookup by logarithmic derivatives, or a Hadamard check: an R1CS argument on Kronecker-FRI commitments needs no sumcheck.

Status. The algebra of Π_{Lin} is checked in the clear by the Rust module `lincheck`: the closed forms of V_1 , V_{id} , V_{G_η} ; honest runs with padded matrices; a wrong a caught by the weighted sums; a forged ζ_k or e_k that repairs the weighted sums caught by the column or row lookup. The compiled protocol (Proposition 25 in `batch`) and the Lean formalisation of this section are the next steps.

6 Relation to known techniques and status

Writing an inner product as the middle coefficient of a product with a reversed polynomial is classical, and appears in pairing-based multilinear commitments (Mercury [3] uses the table form with KZG). DEEP quotients for out-of-domain values are standard in FRI-based STARKs. The new points here are: (1) the reversed virtual word w_b^* , which reads the second vector from the *same* Reed–Solomon commitment thanks to $L^{-1} = L$; (2) the reuse of Kronecker-FRI’s opening polynomial, virtual word and single folding test, with a fully proved soundness bound (Lemma 9); (3) the table-form evaluation kernel E_z^* , which gives evaluation openings on the same commitments; (4) a Hadamard check in which the out-of-domain values are DEEP quotient words inside the same batch as the opening polynomial, so that one degree-8 curve and one folding test carry the whole argument (Lemma 17).

Prover cost and the sumcheck baseline. The prover of Π_{IP} computes one product $V_a V_b^*$ of two polynomials of degree $< N$ over the base field (three NTTs of size $2N$), one NTT of size M and one Merkle tree for w_A , and then runs the folding test; every round-1 object lies in the base field, since V_a, V_b do. The natural baseline is the sumcheck route: n rounds of a degree-2 sumcheck on $\sum_w a(w)b(w)$, followed by openings of a and b at the random point. For that baseline the fastest prover we know of in our own work is the tree circuit of [5], which produces the evaluations, hypercube sums and sumcheck round polynomials of a multilinear polynomial from one binary-tree recursion; the comparison of the two routes at equal soundness is left to the implementation paper.

Status.

- *Lean 4*. Sections 1–2 are formalised in the Kronecker-FRI Lean project, files `TableForm.lean` and `InnerProduct.lean`, without `sorry` and with the single axiom of that project (correlated agreement for curves, KF Thm 2.16, here with $t = 3$). Lemma 2: `eval_table`; Lemma 3: `inner_product`; Lemma 4: `splitG_iff`, `natDegree_PhiG_le`, `identityG`; Lemma 5: `wrev_ev`, `wrev_eq_iff`, `inv_image_closed`, `ncard_inv_image`; completeness: `honestIP_prob_one`; Lemma 9: `batching_IP`; soundness: `soundness_IP`, and `soundnessC_IP` for any folding arity 2^k ; knowledge and binding: `knowledge_IP`, `binding_IP`. The common step of KF Lemma 7.4 and Lemma 9, correlated agreement on one fibre-closed set for a curve of any degree, is the separate theorem `curve_agreement`.
- *Rust*. The module `ip` of the Kronecker-FRI crate implements Π_{IP} with Merkle caps, arity 2^k and Fiat–Shamir, on the commitments of `pcs`. Its tests check Lemmas 3, 4 and 5(1), completeness for $n \leq 10$, two rates and arities 2 to 16, and the rejection of three cheating provers (a wrong S ; a wrong S absorbed into a polynomial A of degree N , as in KF Remark 4.13; opening polynomials computed from a table $b' \neq b$).
- *Lean 4, Section 3*. File `Hadamard.lean`, same axiom, no `sorry`. Lemma 13: `deep_step`, `deep_honest`; completeness: `honestHad_accepts`, and `honestHad_prob` for the bound $1 - 3M/|\mathbb{F}|$; Lemma 17: `batching_Had`; Theorem 18: `prob_bad_le` (the bad pairs (γ, θ)), `soundness_Had`, and `soundnessC_Had` for any folding arity 2^k ; knowledge: `knowledge_Had`. The step “for good (γ, θ) , the argument of KF Thm 7.8” is the separate theorem `ft_family_bound`: for any family of words $w_0(\beta)$, the acceptance probability is at most $|\mathcal{G}|/|\mathbb{F}| + \varepsilon_{\text{fold}} + (1 - \delta)^\kappa$.
- *Lean 4, Section 4*. File `BatchStmts.lean`, same axiom, no `sorry`. Statements: `BStmt` (`eval`, `ip`, `had`), `Dset`, `Bset`, `RelBatchS`; the words are indexed by the finite type `BWord` ($D \oplus B \oplus \mathcal{H} \oplus [J] \oplus [J] \oplus \mathcal{H} \times [3]$) with a fixed enumeration, and `curve_agreement_idx` is correlated agreement for such an index type. Lemma 23: `batching_B`; Theorem 24: `prob_bad_had`

(one false Hadamard check), `prob_bad_B`, `soundness_B`, `soundnessC_B` (any folding arity); knowledge: `knowledge_B`; completeness: `bwords_honest`, `honestB_accepts`, `honestB_prob`.

- *Rust, Section 4.* The module `batch` implements Π_{Batch} with one multi-value Merkle tree for all w_Q and one for all w_A ; its tests check Lemma 2 for E_z^* , completeness for mixed batches (two Hadamard checks sharing a table, an inner product, two evaluations) for $n \leq 10$ and arities 2 to 16, and the rejection of a batch with one false evaluation, one false inner product, or one false Hadamard check (also against a prover that sets y_3 to the weighted inner product, which only the quotient word $q[w_c, y_3, \gamma]$ catches).
- *Rust, Section 3.* The module `had` implements Π_{Had} on the shared folding-test code of module `ft`, drawing γ, θ by rejection sampling outside the base field $\mathbb{F}_p \supseteq L$, so that no pole lies in L and completeness is perfect (the two first terms of ε_{Had} become $(N - 1)/(|\mathbb{F}| - p)$ each); its tests check completeness for $n \leq 10$, two rates and arities 2 to 16, and the rejection of a wrong y_1 , of a wrong Q (with $y_1 = Q(\theta)$), and of a false instance (c differing from $a \circ b$ in one entry) with three choices of y_3 . Removing $q[w_c, y_3, \gamma]$ from the batch makes the false instance accepted, and a wrong pole in the verifier breaks completeness, so the tests exercise each part.

n	prove (ms)			verify (ms)			proof (KiB)		
	KF	IP	Had	KF	IP	Had	KF	IP	Had
12	7.5	6.2	17.6	2.09	2.13	4.31	138.1	171.4	289.1
14	32.2	25.9	74.0	2.56	2.60	4.97	182.3	233.4	371.1
16	141.4	109.2	314.7	3.24	3.25	5.63	249.8	308.1	458.8
18	548.2	437.8	1353.6	3.87	4.01	6.40	325.0	395.3	565.7
20	2736.6	1947.4	6004.5	4.59	5.13	7.39	391.7	471.1	662.0

Table 2: KF: one evaluation opening (Π_{KF}); IP: one inner product (Π_{IP}); Had: one Hadamard check (Π_{Had}). Configuration KF-8 of the Kronecker-FRI paper (rate 1/4, arity 8, caps of 128 nodes, $\ell = n - 8$), 148 queries, 32-byte salts, $\mathbb{F}_{p^2}$ over Goldilocks; one core of an Intel Xeon at 2.10 GHz; medians; commitments not included. Source: `bench/ip.csv`, `cargo run --release --example ip_bench`.

An inner product of two committed tables costs less prover time than one evaluation opening (the round-1 work is in the base field), about the same verifier time, and about 20% more proof, the extra openings of w_b on the inverse cosets. A Hadamard check costs about 2.2 evaluation openings of prover time (its polynomials Q, A, H lie in the extension field) and 1.7 times the proof, since five oracles are opened per query instead of two; when a, b, c are committed together, one tree with three values per leaf removes two of the five authentication paths per query.

Next. The compiled lincheck (Proposition 25 in `batch`), then the Lean formalisation of Section 5.

References

- [1] A. Mkhida. Kronecker-FRI: A Hash-Based Multilinear Polynomial Commitment from a Coefficient-Extraction Identity. Preprint, 2026. <https://github.com/qoosmo/kronecker-fri>.
- [2] E. Ben-Sasson, D. Carmon, Y. Ishai, S. Kopparty, S. Saraf. Proximity gaps for Reed–Solomon codes. J. ACM 70(5), 2023.
- [3] L. Eagen, A. Gabizon. MERCURY: a multilinear polynomial commitment scheme with constant proof size and no prover FFTs. IACR ePrint 2025/385.

- [4] U. Haböck. Multivariate lookups based on logarithmic derivatives. IACR ePrint 2022/1530.
- [5] A. Mkhida, A. Igider. Multilinear polynomials via tree-based circuit and the sumcheck protocol. IACR ePrint 2026/1469.