UNDER REVIEW

RESCIENCE C

Replication / Statistics

[Re] An independent replication of bootstrap confidence-interval calibration for small samples

Kyunghan Bae[1]

[1]Independent researcher, South Korea

Edited by
(Editor)

Received
–

Published
–

DOI
–

Abstract

We independently implement seven confidence-interval procedures from Hesterberg's study of resampling inference. All 20 printed numerical cells of Table 5 in the extended manuscript agree with direct calculations at the published precision. A fixed-seed experiment evaluates normal and exponential sample means at n=10,20,40, using 5,000 original samples and 1,999 bootstrap resamples per condition. Conditioning on centered or normalized residuals reduces outer simulation noise. The experiment compares one-sided miss probabilities and their Monte Carlo uncertainty, and a paired experiment checks sensitivity to 9,999 resamples. For exponential data at n=10, bootstrap-t coverage is 0.9407, compared with 0.9001 for Student t. The largest-magnitude finite-B tail-probability change is -0.00081. The replication concerns a prespecified small-sample subset of the original study.

1 Original study and target

Hesterberg (2015), Figure 10, compares one-sided miss probabilities for normal and exponential populations. We independently implement its selected small-sample conditions. The publicly accessible 2014 extended manuscript provides detailed methods and exact numerical targets [1, 2]. Table 5 and Figures 20–22 refer to that extended manuscript; Figure 10 refers to the journal article. The exact anchor supplements the simulation target. The interval algorithms are newly written from the equations.

2 Methods

The sample mean is the parameter estimate. The seven intervals are Student t, skewness-corrected t, t with bootstrap standard error, percentile, reverse percentile, expanded percentile, and bootstrap t. Their endpoints are implemented in `replicate.py`, directly from the manuscript definitions. All variances used for Student standard errors have divisor n-1. Bootstrap standard error uses the sample standard deviation of the B resampled means. Empirical quantiles use linear interpolation (Hyndman–Fan type 7). Write the sample mean as $\bar{x}$, its standard error as $h = s/\sqrt{n}$, and $c = t_{n-1,0.975}$. For bootstrap means $\bar{x}^*_b$, let Q_p denote the empirical p quantile, h_B their sample standard deviation, and $T^*_b = (\bar{x}^*_b - \bar{x})/(s^*_b/\sqrt{n})$. The intervals are:

Copyright © 2026 K. Bae, released under a Creative Commons Attribution 4.0 International license.
Correspondence should be addressed to Kyunghan Bae (qorud001@gmail.com)
The author declares no competing interests.
Code is available at https://github.com/qorud02/bootstrap-ci-replication – DOI 10.5281/zenodo.23298767..
Data is available at https://github.com/qorud02/bootstrap-ci-replication – DOI 10.5281/zenodo.23298767.

Procedure	Lower and upper endpoints
Student t	$\bar{x} \pm ch$
t with bootstrap SE	$\bar{x} \pm ch_B$
Percentile	$[Q_{0.025}(\bar{x}^*), Q_{0.975}(\bar{x}^*)]$
Reverse percentile	$[2\bar{x} - Q_{0.975}(\bar{x}^*), 2\bar{x} - Q_{0.025}(\bar{x}^*)]$
Expanded percentile	$[Q_{a_n}(\bar{x}^*), Q_{1-a_n}(\bar{x}^*)]$
Bootstrap t	$[\bar{x} - Q_{0.975}(T^*)h, \bar{x} - Q_{0.025}(T^*)h]$

Here $a_n = \Phi(-c\sqrt{n/(n-1)})$. For the skewness-corrected t interval define $g(q) = q + \kappa(1 + 2q^2)$ and $\kappa = m_3/(6s^3\sqrt{n})$, where $m_3 = n^{-1}\sum_i (x_i - \bar{x})^3$. Its endpoints are $\bar{x} + hg(-c)$ and $\bar{x} + hg(c)$ after the flattening rule below. If $\kappa > 0$, arguments below $-1/(4\kappa)$ are set to that value; if $\kappa < 0$, arguments above it are set to it. The zero-skewness limit equals Student t.

The skewness correction uses m_3/s^3. Its quadratic is flattened at the extremum when a nominal endpoint lies beyond it, specifying the source's instruction explicitly. Constant resamples have undefined studentized statistics; the implementation counts and excludes only those statistics from bootstrap-t quantiles. It fails if fewer than 99% remain finite.

Two source populations are studied, N(0,1) and Exponential(mean=1). The six main conditions each use 5,000 independently generated original samples and B=1,999 resamples. The finite-B comparison uses 1,500 fresh exponential samples at n=10 and n=40, with B=1,999 and B=9,999 sharing original samples and the initial resample prefix. PCG64DXSM seeds are fixed before calculation.

Let a left miss mean $L > \mu$ and a right miss mean $U < \mu$. For normal data, $\bar{X}$ is independent of $V = X - \bar{X}$, and the intervals are translation equivariant. Conditional on V and the resample indices, $L = \bar{X} + L_V$ and $U = \bar{X} + U_V$. Since $\bar{X} \sim N(0, 1/n)$, the conditional probabilities are

$$p_L(V) = \Phi(\sqrt{n}L_V), \qquad p_R(V) = \Phi(-\sqrt{n}U_V).$$

For exponential data, $V = X/\bar{X}$ is independent of $\bar{X} \sim \text{Gamma}(\text{shape} = n, \text{scale} = 1/n)$ and the intervals are positively scale equivariant. Thus $L = \bar{X}L_V$ and $U = \bar{X}U_V$. If G_n is the CDF of that gamma law, the conditional probabilities are

$$p_L(V) = 1 - G_n(1/L_V), \qquad p_R(V) = G_n(1/U_V),$$

when the corresponding residual endpoint is positive. For $L_V \leq 0$ the left probability is zero; for $U_V \leq 0$ the right probability is one. We average these conditional weights over independent residual vectors and independently drawn resample indices. The law of total expectation gives the same unconditional miss probability as binary simulation, while conditioning removes the sample-mean component of Monte Carlo variance.

Reported Monte Carlo standard errors are the sample standard deviation of the conditional weights divided by sqrt(R); pointwise 95% intervals use a Student critical value with R-1 degrees of freedom. Paired B differences use the standard error of within-sample differences and the normal critical value 1.96. These intervals quantify outer simulation uncertainty at the specified B. The separate paired experiment assesses finite-B sensitivity. Prespecified sample sizes cover three small-sample conditions.

For descriptive comparisons we use $E = |p_L - 0.025| + |p_R - 0.025|$. This operationalizes the source's qualitative statement about overall coverage accuracy; the source does not specify this particular loss function. Individual tail estimates and their uncertainty remain the primary simulation results.

3 Results

All 20/20 numerical targets in Table 5 agree at their printed precision. The full values and discrepancies are available in `table5_comparison.csv`.
For the deterministic anchor, let $d_n = \sqrt{(n-1)/n}$ and $z = \Phi^{-1}(0.975)$. The four calculated quantities are d_n, z/c, $1 - F_{t_{n-1}}(zd_n)$, and $\Phi(-c/d_n)$, respectively. The following table reports our calculations with more digits than the source's three or four decimal places.

n	Narrowness	z/t	One-sided size	Adjusted alpha/2
5	0.894427	0.705926	0.077233	0.000954
10	0.948683	0.866414	0.047954	0.008551
20	0.974679	0.936427	0.035651	0.015881
40	0.987421	0.968988	0.030114	0.020258
80	0.993730	0.984684	0.027503	0.022588

The exponential experiment gives the following newly computed results. A nominal 95% interval has target left and right miss probabilities of 0.025 each.

n	Method	Left miss	Right miss	Coverage	MCSE(L)	MCSE(R)
10	Student t	0.0040	0.0959	0.9001	0.00017	0.00060
10	Skew-corrected t	0.0122	0.0714	0.9163	0.00027	0.00068
10	t with bootstrap SE	0.0056	0.1049	0.8895	0.00021	0.00063
10	Percentile	0.0181	0.1206	0.8613	0.00036	0.00074
10	Reverse percentile	0.0112	0.1497	0.8391	0.00032	0.00059
10	Expanded percentile	0.0068	0.0863	0.9069	0.00021	0.00066
10	Bootstrap t	0.0096	0.0497	0.9407	0.00024	0.00074
20	Student t	0.0058	0.0749	0.9193	0.00014	0.00044
20	Skew-corrected t	0.0213	0.0506	0.9281	0.00023	0.00046
20	t with bootstrap SE	0.0070	0.0792	0.9138	0.00016	0.00045
20	Percentile	0.0158	0.0807	0.9034	0.00024	0.00051
20	Reverse percentile	0.0084	0.1058	0.8858	0.00019	0.00044
20	Expanded percentile	0.0097	0.0653	0.9250	0.00017	0.00046
20	Bootstrap t	0.0160	0.0381	0.9459	0.00021	0.00049
40	Student t	0.0086	0.0595	0.9319	0.00013	0.00031
40	Skew-corrected t	0.0255	0.0389	0.9355	0.00019	0.00031
40	t with bootstrap SE	0.0095	0.0616	0.9289	0.00014	0.00032
40	Percentile	0.0165	0.0589	0.9246	0.00018	0.00035
40	Reverse percentile	0.0090	0.0779	0.9131	0.00015	0.00032
40	Expanded percentile	0.0131	0.0520	0.9349	0.00015	0.00033
40	Bootstrap t	0.0207	0.0321	0.9472	0.00019	0.00033

At n=10, Bootstrap t has the smallest point-estimated sum of absolute one-sided calibration errors (0.0401). At n=20, Bootstrap t has the smallest point-estimated sum of absolute one-sided calibration errors (0.0221). At n=40, Bootstrap t has the smallest point-estimated sum of absolute one-sided calibration errors (0.0114).
For exponential data at n=10, the bootstrap-t right miss probability is 0.0497 (95% Monte Carlo interval [0.0483, 0.0512]), compared with 0.0959 for Student t. Their total coverages are 0.9407 and 0.9001, respectively. Bootstrap t ranks first here while retaining appreciable right-tail error relative to the 0.025 target.
For normal data, all three Student-t estimates have Monte Carlo intervals containing their exact one-sided probability of 0.025. Bootstrap-t miss estimates at n=20 and n=40 are slightly above the nominal target; their pointwise intervals and the percentile undercoverage are shown below and in the comparison ledger.
The largest-magnitude finite-B point change is -0.00081 (expanded_B9999_minus_B1999, n=10, right tail; paired 95% Monte Carlo interval [-0.00109, -0.00053]). All paired differences are reported in `paired_comparisons.csv`.
The run recorded 0 degenerate studentized resamples and 57 skewness-flattening events across the main and sensitivity calculations. The latter count includes the same sensitivity original samples once per B. The primary experiment took 731.6 seconds.




Figure 1. One-sided miss probabilities with pointwise 95% Monte Carlo intervals. All points are newly computed; the dashed line is 0.025. The exponential panels use different vertical ranges.




Figure 2. Paired changes in right-tail miss probability from 1,999 to 9,999 resamples, with 95% Monte Carlo intervals. Original samples and initial resamples are shared; the dashed line is zero.

4 Interpretation and scope

The deterministic anchor is reproduced at source precision. The simulation comparison provides independent small-sample evidence about the original interval-calibration conclusions. The original curves are supplied as figures without tabulated ordinates. `table5_comparison.csv` contains the exact numerical anchor. `original_results_comparison.csv` distinguishes theoretical Student-t probabilities, nominal calibration targets and qualitative ranking checks. Point-estimate rankings are descriptive, and the Monte Carlo intervals are pointwise.
We explicitly reduce R and B relative to the source and quantify both outer uncertainty and finite-B sensitivity. The target is the selected small-sample regime. The large-n cutoffs in extended Figure 22, second-order convergence rates and dependent, discrete or heavy-tailed populations require separate experiments. The conditioning argument uses the stated normal and exponential distributional assumptions.

5 Reproducibility

Python 3.12 and the pinned requirements suffice. `python run_all.py` generates all numerical results; `python analyze.py` creates this article and figures. The numeric source trace, seeds, versions, per-condition timings and failure counts are supplied. All main and finite-B conditional weights are archived in `conditional_miss_samples.npz`, allowing independent reanalysis. The source paper and figures are linked in `SOURCE_TRACE.md`. All observations are generated from the stated probability distributions.

Funding

This research received no external funding.

References

1. T. Hesterberg. **What Teachers Should Know about the Bootstrap: Resampling in the Undergraduate Statistics Curriculum**. 2014. arXiv:1411.5279v1. URL: https://arxiv.org/abs/1411.5279v1.
2. T. C. Hesterberg. "What Teachers Should Know About the Bootstrap: Resampling in the Undergraduate Statistics Curriculum." In: **The American Statistician** 69.4 (2015), pp. 371–386. DOI: 10.1080/00031305.2015.1089789.