Mean

What it is. The value obtained by adding all values of a set number of observations divided by the number of observations. The mean is also known as the average.

Important. The mean is often used as a measure for central tendency`in symmetric distributions. Since outliers can have a large effect on the mean, for skew distributions, we prefer the median as a measure of location.

Range

What it is. The range is the difference between the highest and lowest value.

Important. When the range is big, it means that there is large variety in the dataset. If the range is small, it means that there is little variety in the dataset.

Event

What it is. A thing that happens or takes place.

Important. nan

Population

What it is. The population is a well defined set of subjects or objects A statistical population can be a group of existing objects. This group can potentially be of infinite size.

Important. nan

Sample

What it is. A proportion of subjects collected from a population on which you are going to perform research.

Important. nan

Sensitivity

What it is. Sensitivity (also called the true positive rate) measures the proportion of actual positives that are correctly identified as such (e.g., the percentage of diseased people who are correctly identified as having the condition). (30)

Important. When the sensitivity is high, it means that a low amount of people who are actually ill are tested negative on that test. When the sensitivity is low it means that a lot of people who are ill are missed.

Unit

What it is. 1. Unit of observation: the who or what within a particular research on which data is collected 2. Unit of measurement: the type of measurement in which a certain magnitude is expressed

Important. nan

Variable

What it is. A characteristic or numerical value which can take on different values.

Important. nan