

CBEA: Competitive balances for taxonomic enrichment analysis

Quang P. Nguyen^{1,2}, Anne G. Hoen^{1,2†}, H. Robert Frost^{1*††},

1 Department of Biomedical Data Science, Geisel School of Medicine at Dartmouth College, Hanover, NH, USA

2 Department of Epidemiology, Geisel School of Medicine at Dartmouth College, Hanover, NH, USA

†These authors jointly supervised this work.

†Corresponding author

* hildreth.r.frost@dartmouth.edu

Abstract

Research in human associated microbiomes often involves the analysis of taxonomic count tables generated via high-throughput sequencing. It is difficult to apply statistical tools as the data is high-dimensional, sparse, and compositional. An approachable way to alleviate high-dimensionality and sparsity is to aggregate variables into pre-defined sets. Set-based analysis is ubiquitous in the genomics literature, and has demonstrable impact in improving interpretability and power of downstream analysis. Unfortunately, there is a lack of sophisticated set-based analysis methods specific to microbiome taxonomic data, where current practice often employs abundance summation as a technique for aggregation. This approach prevents comparison across sets of different sizes, does not preserve inter-sample distances, and amplifies protocol bias. Here, we attempt to fill this gap with a new single sample taxon enrichment method with using a novel log-ratio formulation based on the competitive null hypothesis commonly used in the enrichment analysis literature. Our approach, titled competitive balances for taxonomic enrichment analysis (CBEA), generates sample-specific enrichment scores as the scaled log ratio of the subcomposition defined by taxa within a set and the subcomposition defined by its complement. We provide sample-level significance testing by estimating an empirical null distribution of our test statistic with valid p-values. Herein we demonstrate using both real data applications and simulations that CBEA controls for type I error even under high sparsity and high inter-taxa correlation scenarios. Additionally, it provides informative scores that can be inputs to downstream analyses such as prediction tasks.

Author summary

The study of human associated microbiomes relies on genomic surveys via high-throughput sequencing. However, microbiome taxonomic data is sparse and high dimensional which prevents the application of standard statistical techniques. One approach to address this problem is to perform analyses at the level of taxon sets. Set-based analysis has a long history in the genomics literature, with demonstrable

impact in improving both power and interpretability. Unfortunately, there is not a lot of research in developing new set-based tools for microbiome taxonomic data specifically, given its unique features compared to other 'omics data types. We developed a new tool to generate taxon set enrichment scores at the sample level through a novel log-ratio formulation based on the competitive null hypothesis. Our scores can be used for statistical inference at both the sample and population levels, as well as inputs to other downstream analyses such as prediction models. We demonstrate the performance of our method against competing approaches across both real data analyses and simulation studies.

Introduction

The microbiome is the collection of microorganisms (bacteria, protozoa, archaea, fungi, and viruses) which co-exist with their host. Previous research has shown that changes in the composition of the human gut microbiome are associated with important health outcomes such as inflammatory bowel disease [?], type II diabetes [?], and obesity [?]. To understand the central role of the microbiome in human health, researchers have relied on high-throughput sequencing methods, either by targeting a specific representative gene (i.e. amplicon sequencing) or by profiling all the genomic content of the sample (i.e. whole-genome shotgun sequencing) [?]. Raw sequencing data is then processed through a variety of bioinformatic pipelines [?, ?], yielding various data products, one of which are taxonomic tables which can be used to study associations between members of the microbiome and an exposure or outcome of interest.

However, there exists unique challenges in the analysis of these taxonomic count tables [?, ?]. The data is sparse, high-dimensional, and likely compositional [?, ?, ?]. Even though these problems are challenging, a very approachable solution is to use set-based analysis methods, also termed gene set testing in the genomics literature [?, ?]. Aggregated variables can be less sparse, and testing on a smaller number of features can reduce the multiple-testing burden. As such, gene set testing methods have been shown to increase power and reproducibility of genomic analyses. Furthermore, through the usage of functionally informative sets defined apriori based on historical experiments (for example MSigDB [?], and Gene Ontology [?]), gene set analysis also allows for more biologically informative interpretations.

There exists a diverse set of methods already developed in this field. Traditional methods utilize the hypergeometric distribution to test for the overrepresentation of a gene set using a candidate list of genes screened based on a marginal model [?]. Unfortunately, these approaches are sensitive to the differential expression test as well as the chosen threshold when trying to select genes for the candidate list. Aggregate score methods, which are generally more preferred [?], instead assigns a score for each gene set based on gene-specific statistics such as z-scores or fold change. Of these approaches, methods such as GSEA [?] performs a test for each gene set at the population level, summarizing information across all samples. Conversely, methods such as GSVA [?] and VAM [?], generate enrichment scores at the sample level and are more akin to a transformation. In addition to being able to screen for enriched sets per sample, this strategy also allows for the flexible incorporation of different downstream analyses, such as fitting prediction models, or performing dimension reduction.

In microbiome research, even though no explicit enrichment analysis is performed, researchers often aggregate taxa to higher Linnean classification levels such as genus, family, or phylum. However, despite this interest, there is limited research done to

extend existing set-based methods to microbiome relative abundance data. Some software suites, such as *MicrobiomeAnalyst*, do offer tools to perform enrichment testing with curated taxon sets [?]. However, the approach used in *MicrobiomeAnalyst* is a form of overrepresentation analysis at the population level and therefore similarly sensitive to the differential abundance approach used and p-value threshold. One of the primary challenges for adapting gene set analysis to the microbiome context is the compositional nature of the data. Sequencing technologies constrain the total number of reads, and samples are expected to have the same number of reads instead of DNA content [?, ?]. However, different samples still yield arbitrarily different total read counts [?, ?], suggesting the use of normalization methods to allow for proper comparison of feature abundances across samples. However, microbiome data sets do not follow certain assumptions that enable the cross-application of methods from similar fields (such as RNA-seq) [?, ?]. For example, DESeq2's *estimateSizeFactors* [?] assumes that the majority of genes acts as housekeeping genes with constant expression levels across samples. As such, practitioners often rely on total sum normalization to transform count data into relative proportions that sum to one [?]. Some studies have provided empirical performance evaluations supporting this normalization schema [?]. Since this approach imposes a sum constraint on the data, post normalization microbiome data sets are compositional [?], which means that the abundance of any taxa can only be interpreted relative to another. Under this scenario, log-ratio based approaches from the compositional data analysis (CoDA) literature [?] are motivated to address this issue.

Unfortunately, the standard practice for aggregating variables using element-wise summations (referred to as amalgamations in the CoDA literature), does not adequately address the compositional issue [?]. First, inter-sample Aitchison distances computed on original parts are not preserved after amalgamation [?]. This means that cluster analyses might show different results depending on the level of amalgamation and differs from the those computed from original variables. Second, amalgamations do not allow for comparison between sets of different sizes within the same experimental condition since larger sets will have larger means and variances. Third, considering that each taxa has specific measurement biases [?], an amalgamation based approach would make the bias of the amalgamated variable dependent on the relative abundance of the its constituents. In other words, if taxon 1 has abundance A_1 and bias B_1 , while taxon 2 has abundance A_2 and bias B_2 , then the bias of the aggregate variable (for example, a genera) is $(A_1B_1 + A_2B_2)/(A_1 + A_2)$ (see Appendix 1. from McLaren et al. [?]). This means that bias invariant approaches (such as analyses of ratios) would no longer be invariant when applied to amalgamated variables as bias now varies across samples. The alternative would be to multiply the proportions rather than to sum them [?].

Here, we present a taxon-set testing method for microbiome relative abundance data that addresses the aforementioned issues. Our approach generates enrichment scores at the sample level similar to GSVA [?] and VAM [?]. We leverage the concept of the Q_1 competitive hypothesis presented in Tian et al. [?] to formulate the enrichment of a set as the compositional balance [?] of taxa within the set and remainder taxa using multiplication as the method of aggregating proportions [?]. This well-defined null hypothesis allows us to perform significance testing with interpretable results through estimating the empirical distribution of our statistic under the null that can also account for variance inflation due to inter-taxa correlation [?].

In the following sections, we present our approach titled competitive balances for taxonomic enrichment analysis (CBEA). First, we presented the step-by-step formulation of CBEA and discuss its statistical properties. Second, we detailed our evaluation strategy using both real data and parametric simulations, and the methods we're

comparing against. Third, we presented results on enrichment testing using CBEA for single samples as well as at the population level. Fourth, we showed the performance of CBEA in downstream disease prediction. Finally, we discussed our results and the limitations of our method. An R package implementation of CBEA can be found on GitHub (qpmnguyen/CBEA).

Materials and Methods

Competitive balances for taxonomic enrichment analysis (CBEA)

The CBEA method generates sample-specific enrichment scores for microbial sets using products of proportions [?]. Details on the computational implementation of CBEA can be found in the Supplementary Materials. The CBEA method takes two inputs:

- **X**: n by p matrix of positive proportions for p taxa and n samples measured through either targeted sequencing (such as of the 16S rRNA gene) or whole genome shotgun sequencing. Usually **X** is generated from standard taxonomic profiling pipelines such as DADA2 [?] for 16S rRNA sequencing, or MetaPhlAn2 [?] for whole genome shotgun sequencing. CBEA does not accept X matrices with zeroes since it invalidates the log-ratio transformation. Users can generate a dense matrix X using a method of choice, however the default mode for CBEA would be add a pseudocount of 10^{-5} if zeroes are detected in the matrix.
- **A**: p by m indicator matrix annotating the membership of each taxon p to m sets of interest. These sets can be Linnean taxonomic classifications annotated using databases such as SILVA [?], or those based on more functionally driven categories such as tropism or ecosystem roles ($A_{i,j} = 1$ indicates that microbe i belongs to set j).

The CBEA method generates one output:

- **E**: n by m matrix indicating the enrichment score of m pre-defined sets identified in **A** across n samples.

The procedure is as follows:

1. **Compute the CBEA statistic:** Let **M** be a n by m matrix of CBEA scores. Let $M_{i,k}$ be CBEA scores for set k of sample i :

$$M_{i,k} = \sqrt{\frac{\sum_k A_{ik}(p - \sum_k A_{ik})}{p}} \ln \left(\frac{g(\mathbf{X}_{i,j} | \mathbf{A}_{j,k} = 1)}{g(\mathbf{X}_{i,j} | \mathbf{A}_{j,k} \neq 1)} \right) \quad (1)$$

where $g(\cdot)$ is the geometric mean. This represents the ratio of the geometric mean of the relative abundance of taxa assigned to set k and remainder taxa.

2. **Estimate the empirical null distribution** Enrichment scores represent the test statistic for the Q_1 null hypothesis H_o that relative abundances in **X** of members of set k are not enriched compared to those not in set k . Since the distribution of CBEA under the null vary depending on data characteristics (Fig ??), an empirical null distribution will be estimated from data.

- **Compute the CBEA statistic on permuted and un-permuted X.** Let \mathbf{X}_{perm} be the column permuted relative abundance matrix, and \mathbf{M}_{perm}

be the corresponding CBEA scores generated from \mathbf{X}_{perm} . Similarly, we have \mathbf{M}_{unperm} be CBEA scores generated from \mathbf{X} .

- **Estimate correlation-adjusted empirical distribution for each set.**
For each set, a fit a parametric distribution to both \mathbf{M}_{perm} and \mathbf{M}_{unperm} . The location measure estimated from \mathbf{M}_{perm} and the spread measure estimated from \mathbf{M}_{unperm} will be combined as the correlation-adjusted empirical null distribution \mathbf{P}_{emp} for each set. Two available options are the normal distribution and the mixture normal distribution. For the normal distribution, parameters were estimated using the method of maximum likelihood implemented in the *fitdistr* package [?]. For the mixture normal distribution, parameters were estimated using an expectation-maximization algorithm implemented in the *mixtools* package [?].

3. **Calculate finalized CBEA scores with respect to the empirical null.**
Enrichment scores $\mathbf{E}_{i,k}$ are calculated as the cumulative distribution function (CDF) values or z-scores with respect to \mathbf{P}_{emp} distribution. Raw p-values can be calculated by subtracting \mathbf{E} from 1.

Properties of CBEA

CBEA and balances of groups of parts

The CBEA statistic is based on the multiplication-based aggregation approach used to calculate balances between groups of parts [?]. These balances are computed using the isometric log ratio (ILR) transformation [?] formula. For a given balance i splitting variables across sets R and S , we have the balance coordinate x_i^* as:

$$x_i^* = \sqrt{\frac{rs}{r+s}} \log \left(\frac{g(X_{j|j \in R})}{g(X_{j|j \in S})} \right) \quad (2)$$

where r and s are the cardinalities of sets R and S respectively, $g(z)$ is the geometric mean, and X_j are values of the original predictors with indexes defined by membership in R and S .

CBEA belongs to a set of methods that seeks to leverage compositional balances for the analysis of microbiome data [?, ?, ?, ?]. Unlike methods such as PhILR [?], CBEA does not provide a sequential binary partition that forms the basis for ILR procedure [?] and is therefore not a subclass of ILR. A similar method to CBEA would be phylofactor [?]. However, instead of performing an optimization procedure to identify interesting balances, CBEA constructs balances apriori using pre-defined sets, and formulates the enrichment of a set as the scaled log-ratio between the center of the subcomposition represented by microbes within the set and the center of the subcomposition represented by remainder taxa. This formulation aligns with the Q_1 null hypothesis from the gene set testing literature [?].

Estimating the null distribution

We can assume that the CBEA statistic, similar to other log-ratio based transforms, follows a normal distribution [?, ?]. However, when applying CBEA for hypothesis testing at the sample level, it is expected that the researcher would be testing a large number of hypotheses. Under the assumption that the number of truly significant hypotheses is low, Efron [?] showed that estimating the null distribution of the test

statistic directly (termed the empirical null distribution) is much more preferable than using the theoretical null due to unobserved confounding effects inherently part of observational studies. As such, to perform significance testing using CBEA, we also estimated the null distribution from observed raw CBEA variables.

This assumption is also supported by preliminary simulation studies (detailed below). In panel A of Fig ??, we simulated microbiome taxonomic count data under the global null across different data features and compute raw CBEA scores and compute kurtosis and skewness. It can be seen that the characteristics of the null change depending on sparsity and inter-taxa correlation. Sparsity seems to drive the distribution to be more positively skewed while inter-taxa correlation encourages platykurtic (negative kurtosis). The effect is most dramatic under both high inter-taxa correlation and sparsity. This heterogeneity further supports the decision to estimate an empirical null distribution, as suggested by Efron [?].

Additionally, the degree of kurtosis and skewness also suggests that the normal distribution itself might not be a good approximation of the null. To address this issue, we also evaluated a two-component normal mixture distribution. Panel B of Fig ?? demonstrates the goodness of fit of the mixture normal and the normal distribution using Kolmogorov-Smirnov (KS) test statistic computed on fitted normal and mixture normal distribution when fitted on CBEA scores in simulation scenarios under the global null. We can see that the mixture normal distribution is a better fit (lower KS scores) than the normal distribution across both sparsity and correlation settings.

We performed our empirical null estimation by fitting our distribution of choice and computing relevant parameters on raw CBEA scores on taxa-permuted data (equivalent to gene permutation in the gene expression literature). As such, the null distribution is characterized by scores computed on sets of equal size with randomly drawn taxa.

Variance inflation due to inter-taxa correlation

When taxa within a set are highly correlated, the variance of the sample mean of taxon-wise statistics is inflated. Without loss of generalizability, for a set of taxa with taxa-specific statistics x_1, \dots, x_p we have the variance of the mean \bar{x} to be:

$$Var(\bar{x}) = \frac{1}{m^2} \left(\sum_{i=1}^m (\sigma_i^2) + \sum_{i < j} \rho_{ij} \sigma_i \sigma_j \right) \quad (3)$$

where σ_i is the standard deviation of taxon i while ρ_{ij} is the correlation between i and j . The second term of (??) is the correlation dependent variance component, which goes to 0 if there is no correlation. The CBEA statistic follows a similar pattern. Since the geometric mean of a set of variables is equivalent to the exponential of the arithmetic mean of their logarithms, we can re-write CBEA score for a set k with size K as follows:

$$M_{i,k} = \sqrt{\frac{K(p-K)}{K+(p-K)}} (\overline{\log X_{i,j|j \in K}} - \overline{\log X_{i,j|j \notin K}}) \quad (4)$$

where p is the overall number of taxa, j is the index of a taxa and K is the set of indices of taxa in set k . The CBEA statistic then looks similar to a t-statistic for difference in means of log-transformed proportions. As such, the pooled variance of CBEA is dependent on the variance inflation of both mean components $\overline{\log X_{i,j|j \in K}}$ and $\overline{\log X_{i,j|j \notin K}}$. The result of this variance inflation is inflated type I error since highly correlated sets are also detected as significantly enriched.




Fig 1. Properties of the null distribution of CBEA under the global null simulations. Panel (B) presents kurtosis and skewness of CBEA scores while panel (A) presents the goodness of fit (as Kolmogorov-Smirnov D statistic) for mixture normal and normal distributions. Panel (C) is a density plot of the shape of the null distribution. Results indicated the necessity of estimating an empirical null and demonstrating that the mixture distribution was the better fit compared to the basic normal.

However, as Wu et al. [?] showed, performing column permutation to estimate the null distribution of a competitive test statistic doesn't allow for adequate capture of this variance inflation factor since the permutation procedure disrupts the natural correlation structure of the original variables. It is important to address this problem since there is strong inter-taxa correlation within the microbiome [?]. Our strategy for addressing issue is location (or mean) estimate from the column permuted raw score matrix with the spread (or variance) estimate from the original un-permuted scores. This still allows us to leverage the null generated via column permutation while using the proper variance estimate taken from scores where the correlation structure has not been disrupted. As such, this procedure assumes that the variance of the test statistic under the alternate hypothesis is the same as that of the null. Details of the computational implementation to this estimation process can be found in the

Supplementary Materials. 223

However, set-based analysis is an exploratory approach that can help generate 224
functionally informative hypotheses, and as such users might not want strict type I error 225
control in favor of higher power. This is especially true for competitive hypotheses, 226
where its stricter formulation compared to the self-contained approach implies that the 227
test naturally has lower power [?, ?]. Furthermore, sets that are highly correlated 228
compared to background can be biologically relevant. Therefore, CBEA provides an 229
option for users to specify whether correlation adjustment is desired. 230

Evaluation 231

We based our evaluation strategy on gene set testing benchmarking standards set by 232
Geistlinger et al. [?] and utilized the same approaches whenever possible. All data sets 233
are obtained from either the *curatedMetagenomicData* [?] and *HMP16SData* [?] R 234
packages (2020-10-02 snapshot), or downloaded from the Qiita platform [?]. All code 235
and data sets used for evaluation of this method is publicly available and can be found 236
on GitHub (qpmnguyen/CBEA_analysis). Additional packages used to support this 237
analysis includes: *tidyverse* [?], *pROC* [?], *phyloseq* [?], *mia* [?], *targets* [?]. 238

Statistical significance 239

We evaluate the inference procedure of CBEA compared to alternate methods using two 240
approaches: randomly sampled taxa sets and sample label permutation. These analyses 241
were performed on the 16S rRNA gene sequencing of the oral microbiome from the 242
Human Microbiome Project [?, ?]. This data set contains 369 samples split into two 243
subsites: supragingival and subgingival. We processed this data set by removing all 244
samples with total read counts less than 1000 and OTUs whose presence (at least 1 245
count) is in 10% of samples or less. 246

Sample-level inference 247

Due to CBEA's self-contained null hypothesis, we can perform inference at the sample 248
level for the enrichment of a set. We evaluated this application by generating one 249
random taxa of different sizes $S \in \{20, 50, 100, 150, 200\}$ across 500 iterations. Random 250
sets can act as our estimate for type I error since this matches the CBEA null 251
hypothesis stated in ??, where we expect within each sample sets of randomly drawn 252
taxa should not be significantly enriched compared to the remainder background taxa. 253
For this evaluation, we estimated type I error as the fraction of samples that detects our 254
random set as significant at a p-value threshold of 0.05 with confidence bands computed 255
from the standard error across all iterations. Additionally, this analysis also 256
demonstrate whether CBEA is sensitive to different set sizes. 257

Population-level inference 258

We can perform enrichment testing at the population level by generating corresponding 259
sample level CBEA scores and performing a two-sample test such as Welch's t-test. In 260
order to evaluate CBEA under this context, we generated CBEA scores of sets 261
representing genus-level annotation in above gingival data set [?, ?] and applied a t-test 262

to test for enrichment (similar to GSVA [?]) across a randomly generated variable indicating case/control status (repeated 500 times). Type I error is estimated as the fraction of sets per iteration found to be significantly enriched with confidence bands computed from the standard error across all iterations. In addition, we also performed a random set analysis assessment, where we generated 100 sets of different set sizes $S \in \{20, 50, 100, 150, 200\}$ and evaluated the fraction of genera that were found to be differentially abundant across the original labels (supragingival versus subgingival subsite). 95% confidence intervals were computed using the Agresti-Couli approach [?].

Phenotype relevance

We want to evaluate whether sets found to be significantly enriched by CBEA are relevant to the research question. To perform this assessment, we relied on the gingival data set mentioned prior [?,?]. This data set was chosen due to its clear biological interpretation that can serve as the ground truth. Specifically, we expect aerobic microbes to be enriched in the supragingival subsite where the biofilm is exposed to the open air, while conversely anaerobic microbes thrive in the subgingival site [?]. Genus-level annotations for microbial metabolism from Beghini et al. [?] and was obtained from the GitHub repository associated with Calagaro et al. [?]. For sample-level inference, we assessed power as the fraction of supragingival samples where aerobic microbes are significantly enriched. For population-level inference, power is the fraction of sets representing genus level taxonomic assignments that were significant across subsite labels.

In addition to statistical power, we also assessed phenotype relevance through evaluating whether highly ranked sets based on CBEA scores are more likely to be enriched according to ground truth. This is represented by the area under the receiving operator curve (AUROC/AUC) scores computed on CBEA scores against true labels (similar approach was used to evaluate VAM [?]). DeLong 95% confidence intervals for AUROC [?] were obtained for each estimate.

Disease Prediction

CBEA scores can also be used downstream analyses such as disease prediction tasks. We utilized two data sets for this evaluation:

1. Whole genome sequencing of stool samples from inflammatory bowel disease (IBD) patients in the MetaHIT consortium [?]. This data set contains 396 samples from a cohort of European adults, where 195 adults were classified as having IBD (which includes patients diagnosed with either ulcerative colitis or Crohn’s disease). We processed this data by removing all samples with less than 1,000 total read counts as well as any OTU who was present (with non-zero proportions) in 10% of samples or less. Prior to model fitting, we back-transformed relative abundances into count data (to align the format with our 16S rRNA gene sequencing data set) using provided total number of reads aligned to MetaPhlan marker genes (per sample).
2. 16S rRNA gene sequencing of stool samples from IBD patients in the pediatric RISK cohort [?]. This data set contains 16S rRNA gene sequencing samples from a cohort of pediatric patients (ages < 17) from the RISK cohort enrolled in the United States and Canada. Of the 671 samples obtained, 500 samples belong to

patients with IBD. We processed this data set by removing all samples with less than 1,000 total read counts as well as any OTU who was present (at least 1 count) in 10% of samples or less.

We evaluate disease prediction performance by fitting a random forest model [?] using inputs as CBEA scores to classify samples of patients with IBD and healthy controls. Random forest was chosen as a baseline learner due to its flexibility as an out-of-the-box model that is easy to fit. In this instance we evaluated predictive performance of a default random forest model (without hyperparameter tuning) AUROC after 10-fold cross validation. Additionally, we utilized SMOTE to correct for class imbalances [?]. Implementation was done using the *tidymodels* suite of packages [?].

Comparison Methods

We benchmarked the statistical properties of CBEA against existing baseline approaches. For sample-level inference analyses, utilized the Wilcoxon rank-sum test, which non-parametrically tests the difference in mean counts between taxa from a pre-defined set and its remainder similar to CBEA. For assessments at the population level, we compared CBEA against performing a standard test for differential abundance with set-level features generated via element-wise summations instead. We chose DESeq2 [?] and corncob [?] because they represent both methods extrapolated from RNA-seq [?] and those developed specifically for microbiome data.

Since disease prediction models and rankings-based phenotype relevance analyses seek to evaluate the informativeness of CBEA scores instead of relying on computing p-values, we compared performance against other single sample based approaches from the gene set testing literature, specifically ssGSEA [?] and GSVA [?]. Additionally, for evaluating prediction, we also compared performance against a standard analysis plan where inputs are count-aggregated sets with the centered log-ratio (CLR) transformation.

Results

In this section, we present results for evaluating statistical significance, phenotype relevance, and predictive performance. In addition to real data, we also evaluated models based on parametric simulations, where results can be found in the Supplemental Materials.

Statistical Significance

Inference at the sample level

CBEA provides significance testing at the sample level through a self-contained competitive null hypothesis. Generating random sets approximate the global null setting where within each sample, sets generated by randomly sampling taxa should not be significantly more enriched than remainder taxa.

Fig ?? demonstrates type I error of sample-level inference evaluated using the random set approach. The Wilcoxon rank sum test and unadjusted CBEA under mixture normal assumption demonstrated good type I error control at the appropriate α level. This fits with our expectations since the mixture normal distribution has much better fit than the normal distribution especially at the tails (Fig ??). However, other variants




Fig 2. Random taxa set analyses for inference at the sample level of CBEA under different parametric assumptions compared against a Wilcoxon rank-sum test. Type I error (y -axis) was evaluated by generating random sets of different sizes (x -axis) (500 replications per size) and computing the fraction of samples where the set was found to be significantly enriched at $\alpha = 0.05$. Error bars represent the mean type I error \pm sample standard error computed across 500 replications of the experiment. Only the unadjusted CBEA with the mixture normal distribution and the Wilcoxon rank sum test were able to control for type I error at 0.05. All approaches are invariant to set sizes.

of CBEA demonstrated inflated type I error, especially correlation adjusted variants compared to their unadjusted counter parts. Encouragingly, all methods demonstrate consistent performance across all set sizes, with a slight increase in type I error at the highest levels.

Interestingly, simulation results (??) showed an opposite pattern. Adjusted approaches were good at controlling for type I error, especially under the low inter-taxa correlation values within the set (similar to generating random sets where the natural correlation structure is disrupted). In these simulations, unadjusted approaches and the Wilcoxon rank sum test had significant type I error inflation with increasing correlation. All approaches seems to be invariant to the level of data sparsity.

Inference at the population level

Similar to other single sample approaches to gene set testing such as GSVA [?], we can perform inference at the population level by utilizing a two-sample difference in means test. Here, we evaluate using CBEA scores generated under different settings with Welch's t-test in a supervised manner to assess whether a set is enriched across case/control status.

Fig ?? shows results for this scenario using both random sample label and random set evaluations. The random sample label approach (Fig ??A) provides a controlled setting where we can estimate type I error rate controlled at $\alpha = 0.05$. Across all replications, CBEA methods were able to control for type I error at the nominal threshold of 0.05,




Fig 3. Random sample label (A) and random set (B) analyses for population level inference. (A) Type I error (x -axis) was estimated as the overall fraction of sets found to be enriched $\alpha = 0.05$ using randomly generated sample labels (500 permutations). Error bars represent the mean type I error \pm sample standard error. (B) Proportion of significant sets (y -axis) using 100 randomly generated sets of different set sizes (x -axis). Confidence intervals computed using Agresti-Couli method for binomial proportions. For sample label permutation (A), all CBEA approaches were able to control for type I error but not for corncob and DESeq2. For random set analyses (B), all approaches demonstrate similar rate of accepting significant sets and were invariant to overall set size.

with CBEA raw scores being the most performant. Neither output types, correlation adjustment, nor distributional assumption improved performance values. Surprisingly, DESeq2 and corncob both exhibit significantly inflated type I error.

We also assessed the impact of set-size on the inference procedure by testing for enrichment using the original sample labels but with randomly sampled sets of different sizes (Fig ??B). Overall we observed very similar values across CBEA as well as corncob and DESeq2, suggesting that no individual method is systematically identifying too many significant sets. Additionally, similar to analogous analyses at the sample level, no

approach were significantly sensitive to changes in set sizes.

377

Phenotype Relevance

378

Inference at the sample level

379




Fig 4. Statistical power (A) and score rankings (B) to assess phenotype relevance. (A) Power (x -axis) was estimated as the overall fraction of aerobic microbes found to be enriched in supragingival samples at $\alpha = 0.05$. 95% confidence intervals were computed using the Agresti-Couli approach for binomial proportions. (B) Score rankings were evaluated by comparing computed scores against true values using AUROC (x -axis). DeLong 95 % confidence intervals for AUROC were computed.

In Fig ??, we evaluate whether sets found to be significant by CBEA are relevant to the phenotype of interest. We leveraged the gingival data set as stated in ?? section where we know beforehand that aerobic microbes are more likely to be enriched in supragingival subsite samples and vice versa.

380
381
382
383

We estimated statistical power using this data set as the fraction of supragingival samples where the set representing aerobic microbes were significantly enriched. We observed that adjusted CBEA approaches demonstrate much lower power compared to the Wilcoxon rank-sum test and unadjusted variants. This is surprising given the fact that in statistical significance analyses, the adjusted CBEA approach provides inflated type I error, especially if the normal distribution assumption was chosen, which indicates a mismatch in estimating the null distribution since a high type I error did not result in increased power.

384
385
386
387
388
389
390
391

We also evaluated phenotype relevance by assessing whether enriched sets according to ground truth are preferentially ranked higher using assigned continuous scores (instead of performing a hypothesis test). This aspect is captured through computing AUROC values comparing computed enrichment scores and true labels. Consistent with prior approaches, adjusting for correlation did not improve performance, where the AUROC values are equivalent to using the Wilcoxon Rank sum statistic at around 0.5.

392
393
394
395
396
397

Unadjusted methods were much better at ranking true enriched sets, however the mean AUROC values are lower than alternate single sample enrichment methods (GSVA [?] and ssGSEA [?]) even though this difference is not significant due to overlapping confidence intervals.

The above results were replicated in simulation studies where we observed that adjusted approaches was very conservative and demonstrated significantly lower power (??) with increasing correlation even at the highest evaluated effect sizes. When assessing score rankings, the performance of CBEA was closer to ssGSEA and GSVA compared to real data evaluations, however all single sample approaches were much better than using the W statistic from the Wilcoxon Rank Sum test.

Inference at the population level




Fig 5. Statistical power to assess phenotype relevance of inference tasks at the population level. Power (*x*-axis) was estimated as the overall fraction of sets representing genera that are aerobic or anaerobic microbes found to be differentially enriched across sample type (supragingival or subgingival). 95% confidence intervals were computed using the Agresti-Couli approach for binomial proportions.

We also assessed statistical power for population level inference scenarios using a similar approach. Here, enrichment scores for sets representing all identified genera were computed, and estimated power as the fraction of sets found to be differentially enriched across sample site labels (supragingival or subgingival). We compared these results against performing a differential abundance test of genus level features generated via sum-based approaches. Results are shown in Fig ???. Some CBEA variants, such as CDF outputs for the mixture normal distributional assumption, did not correctly detect as many significant sets as DESeq2 or corncob despite very close performance values. Using raw CBEA scores was best approach, however it did not exceed values obtained from DESeq2 and corncob.

Disease Prediction

Since CBEA can generate informative scores that can discriminate between samples with inflated counts for a set (Fig ??), we want to assess whether they can also act as useful inputs to predictive models. In this section we assessed the predictive performance of a standard baseline random forest model [?] with different single sample enrichment scoring methods as inputs (CBEA, ssGSEA, and GSVA). Additionally, we also compared predictive performance of using these scores against the a standard approach of using the centered log ratio transformation (CLR) on taxon sets aggregated via abundance summations.

We fit our model to two data sets with a similar disease classification task of discriminating patients who are diagnosed with IBD (includes both Crohn’s disease and ulcerative colitis) using only microbiome taxonomic composition. The two data sets represent different microbiome sequencing appraoches: the Gevers et al. [?] data set uses 16S rRNA gene sequencing, while the Nielsen et al [?] data set uses whole genome shotgun sequencing.

Fig ?? illustrates the performance of our model with AUROC as the evaluation criteria. In the 16S rRNA data set, the best performing CBEA variant (CDF values computed from an unadjusted mixture normal distribution) outperforms both GSVA and ssGSEA but not the standard CLR approach. Interestingly, in the whole genome sequencing data set, CBEA outperforms CLR, but were more similar in performance to GSVA. However, due to large confidence intervals, no method were significantly out-performing other approaches evaluated. As such, these results indicated that CBEA generated scores are can be informative, providing competitive performance when acting as inputs to disease predictive models. Simulation studies (??) showed similar results, however CBEA was more consistently underperforming compared to CLR across all scenarios. Interestingly, the performance gap decreases with increasing sparsity levels.

Discussion

Inference with CBEA

CBEA is a microbiome-specific approach to generate sample specific enrichment scores for taxonomic sets defined apriori. The formulation of CBEA as a comparison between taxa within the set and its complement corresponds to the competitive null hypothesis in the gene set testing literature [?]. Since this null hypothesis is self-contained per sample, this allows users perform enrichment testing at the sample level. Additionally, in combination with a difference in means test, CBEA can also test for enrichment at the population level across case/control status similar to GSVA [?].

For single-sample analyses, we demonstrated that the CBEA approach (unadjusted with mixture normal parametric assumption) was able to control for type I error at the nominal level of 0.05 under the global null (Fig ??) while also demonstrating respectable power (Fig ??). This performance is consistent across different set sizes as well as our prior distributional fit analyses (Fig ??), where the mixture normal displayed superior fit to the null distribution. Unfortunately, other variants of CBEA demonstrated neither good type I error control nor power. Interestingly, while the adjusted methods showed poor performance in real data evaluations (Fig ??), in simulation studies (??, ??) these approaches were able to control for type I error well with the trade-off of much lower power.




Fig 6. Predictive performance of a naive random forest model trained on CBEA, ssGSEA, GSVA generated scores as well as the standard CLR approach on predicting patients with inflammatory bowel disease versus controls using genus level taxonomic profiles. Data sets used span both 16S rRNA gene sequencing (Gevers et al. [?]) and whole-genome shotgun sequencing (Nielsen et al. [?]). CBEA performs better than GSVA and ssGSEA but not as well as CLR, with the exception of the whole genome sequencing data set.

For population-level inference task, CBEA also performed very well. Under the permutation global null, representing genera abundance using CBEA scores in combination with Welch's t-test controls for type I error the correct α threshold while also keeping respectable power. Since population level enrichment test is equivalent to differential abundance using set-based features, we compared CBEA approach against using element-wise summations with corncob [?] and DESeq2 [?] to test for set-level differential abundance. We chose DESeq2 because it is an older approach from the bulk RNA-seq literature that has strong support for usage in microbiome taxonomic data [?]. Alternately, corncob is a newer method developed specifically for microbiome taxonomic data sets, where taxonomic counts are modeled directly using a beta-binomial distribution instead of relying on normalization via size factor estimation. We observed that using this approach resulted in an inflated type I error compared to all variants of CBEA (Fig ??), yet did not improve power (Fig ??). Results for CBEA approaches were replicated in simulation analyses, however for corncob and DESeq2 we observed an opposite effect: in simulation experiments, both methods show good type I error control but low power (??, ??).

We hypothesized that the discrepancy between simulation and real data evaluations can arise due to differences in our assumptions regarding the data generating process that informed our simulation schema. For the non-zero component of each taxa, we sampled from the same negative binomial distribution where designated enriched taxa were generated with inflated means (but the same dispersion). These marginals were simulated to account for block exchangeable correlation within the enriched set only. This might affect our results in two ways: First, our simulation scenario ensures that any designated non-enriched taxa is identical to each other. This is not the case for real data, where our null scenarios involves randomly sampled sets that might by chance all have taxa with inflated means compared to remainder taxa. This is represented in ??, where the distribution of type I error across our 500 replications are right skewed for underperforming CBEA variants, indicating that these approaches are much more sensitive compared to the Wilcoxon rank sum test or unadjusted CBEA with mixture normal distribution. Second, as described in the ?? section, we did not consider taxon-specific biases that distort the observed relative abundance of taxa compred to true values [?]. In the context of sum-based aggregations, the resulting bias of the aggregated taxon are dependent on the relative abundances of the contributing taxa (Appendix I in [?]). Conceptually, this means that measurement error for a taxon aggregate is different across samples as relative abundance of contributing changes, leading to issues when attempting to perform inference. As such, we expect methods like corncob or DESeq2 when performed on such sum aggregates in the presence of taxon-specific biases to have inflated type I error compared to our taxon-ratio based approach. Conversely, in simulation studies, where taxon-specific biases are absent, corncob and DESeq2 should perform better.

Downstream analysis using predictive models

The sample-level enrichment scores generated by the CBEA method can be used in downstream analyses such as disease prediction. We evaluated whether CBEA can be used to generate set-based features for disease prediction models.

We fitted a basic random forest model [?] to predict continuous and binary outcomes using CBEA generated scores as inputs. Similar to our inference analysis, we compared CBEA against both ssGSEA and GSVA. Additionally, we also evaluated CBEA with the approach where counts of a set were aggregated using sums and applied the centered log-ratio transformation (CLR). This is because CLR is considered standard practice in using microbiome variables as predictors for a model [?]. Results showed that CBEA generate scores perform well across both real data settings and simulation scenarios. Since predictive models consider the effect of variables jointly (and in the case of random forest, consider interactions as well), good performance indicates that CBEA scores can capture joint distribution of sets, enabling both uniset and multi-set type analyses. Comparatively, CBEA generated scores outperformed other enrichment score methods (GSVA and ssGSEA), suggesting that it is more tailored for microbiome taxonomic data sets. This is consistent with our sample ranking analysis (Fig ??), where CBEA scores are on average more informative when used to rank samples based on their propensity to have inflated counts. However, CBEA did not outperform the CLR approach across our simulation studies, and only marginally performed better in the real data analysis with WGS data. Fortunately, in simulation studies, this performance gap between CLR and CBEA decreases with higher sparsity and correlation, especially in low effect saturation scenarios.

Limitations and future directions

These above results demonstrate the applicability of CBEA under different data analysis scenarios. If researchers are interested in perform inference, they can decide between an unsupervised sample level approach (i.e. screen samples for enrichment of certain characteristics) or a supervised population level approach (i.e. identifying characteristics that are differentially abundant across case/control status). For the unsupervised approach, utilizing the unadjusted CBEA with the mixture normal distribution provides a good initial starting point. In the case where researchers only wants to screen samples with mean-inflated taxon sets (instead of additionally detecting taxon sets with increased correlation), they can apply the adjusted approach, which can be effective at conserving type I error even at high correlation scenarios. However, the trade off for this adjustment is power that decreases with increasing correlation. For the supervised analysis, all CBEA variants control well for type I error and provides adequate statistical power. However, using raw CBEA scores with difference-in-means test such as Welch's t-test is preferable since it requires the least amount of computing resource (no estimation process) while still outperforming using sum-based approach with a standard differential abundance test.

Beyond inference, CBEA scores are flexible and can be useful for downstream analysis. We demonstrated that for a given number of set-based features, CBEA can produce informative scores that contribute to competitive performance of prediction models even in low signal-to-noise ratios with high inter-taxa correlation and sparsity. This is especially true for whole genome sequencing data sets, where CBEA outperforms the standard approach of applying a CLR transformation. Researchers might find CBEA useful under situations of high sparsity and inter-taxa correlation, or if the property of a singular covariance matrix (a byproduct of the CLR transformation [?]) is undesired. Even though we only evaluated prediction models in this manuscript, researchers can benchmark their own usage of CBEA for other downstream tasks such as sample ordination.

However, there are various limitations to our evaluation of CBEA. First, our simulation analysis might not capture the appropriate data-generating distributions underlying microbiome taxonomic data. There is strong evidence to suggest that our zero-inflated negative binomial distribution is representative [?], however other distributions such as the Dirichlet multinomial distribution [?] have been used in the evaluation of prior studies. Second, we were not able to evaluate the phenotypical relevance of enrichment results as in Geistlinger et al. [?] due to limited consistent annotations for microbiome signatures in health and disease, especially those that are experimentally verified (and not just from differential abundance studies). We attempted to perform this evaluation by leveraging the gingival data set similar to [?]. However, we acknowledge that this is not a perfect solution, since oxygen usage label of each microbe in the data set is only available at the genus level, and the difference in counts for obligate aerobes and anaerobes across the supragingival and subgingival sites might not be as clear cut. As such, results from power analyses using this data set is only relative between the comparison methods instead of treated as absolute measures of power or phenotype relevance. Finally, we assumed that taxa within a set are all equally associated with the outcome. This limits our ability to evaluate the performance of CBEA when only a small number of taxa within the set is associated with the outcome, or if there are variability in effect sizes or association direction of taxa within a set.

Our evaluation also showed various drawbacks of the CBEA method itself. First, inference with CBEA at the sample level is limited, and can be affected by inter-taxa correlation if users wish to only detect mean-inflated sets. Second, for downstream

analyses, CBEA might not always perform better than competing methods, especially when being used to generate inputs to predictive models. We hypothesized that this might be due to the lack of fit for the underlying null distribution in high correlation settings, especially the identifiability problem associated the estimation procedure associated with adjusting the mixture normal distribution. As such, we hope to refine the null distribution estimating procedure by either choosing a better distributional form, or to further constrain the optimization procedure of the mixture normal distribution by fixing the third and fourth moments.

In addition, CBEA itself did not consider other aspects of microbiome data. First, across all analyses, we relied on adding a pseudocount to ensure log operations are valid. Users can directly addressing this by incorporating model-based zero correction methods prior to modelling such as in [?] or [?]. However, in our simulation studies, sparsity seems to not have a significant impact on the overall performance of our approach. Second, CBEA also treated all taxa within the set as equally contributing to the set. Incorporation of taxa-specific weights (similar to PhILR [?]) could reduce the influence of outliers, such as rare or highly invariant taxa. Finally, even though for a given set of apriori annotations CBEA can generate useful summary scores, such values are limited in their utility if the annotation themselves are not meaningful. As such, curating and validating sets (similar to MSigDB [?]) based on physiological or genomic characteristics of microbes [?] or their association with human disease (in beta BugSigDB https://bugsigdb.org/Main_Page) can allow for incorporating functional insights into microbiome-outcome analyses.

Conclusion

Gene set testing, or pathway analysis is an important tool in the analysis of high-dimensional genomics data sets. However, limited work has been done developing set based methods specifically for microbiome relative abundance data. We introduced a new microbiome-specific method to generate set-based enrichment scores at the sample level. We demonstrated that our method can control for type I error for significance testing at the sample level, while generated scores are also valid inputs in downstream analyses, including disease prediction and differential abundance.

Acknowledgments

The authors thank Becky Lebeaux, Modupe Coker, Erika Dade, Jie Zhou, and Weston Viles for insightful comments and suggestions that greatly improved the paper.

Supporting information

S1 Fig. Simulation results for type I error evaluation for CBEA sample-level inference. Placeholder text.

S2 Fig. Simulation results for type I error evaluation for CBEA population-level inference. Placeholder.

S3 Fig. Simulation results for phenotype relevance evaluation for CBEA sample-level inference. Placeholder text.

S4 Fig. Simulation results for phenotype relevance evaluation for CBEA population-level inference. Placeholder text.

S5 Fig. Simulation results for predictive performance evaluation for CBEA. Placeholder text.

S6 Fig. Computational performance of CBEA. Placeholder text.

S7 Fig. Distribution of type I error values across all replications in real data random set evaluations for CBEA inference at the sample-level. Placeholder text.

S1 File. **Supplemental derivations.** Includes additional details on addressing variance inflation due to correlation in CBEA, simulation analyses, and run-time performance.