

QRU • HEALTH



BOOK

QRU Factory Acceptance Test — Understanding Sleep and Rest — Book

A Sleep & Rest learning product

Foundational • General public

**QRU Factory Acceptance Test - Understanding
Sleep and Rest - Book**

QRU PRESS™

Edition 1

TREASURE STANDARD(TM) CERTIFIED

QRU Book - Health

QRU Factory Acceptance Test - Understanding Sleep and Rest - Book

Manufactured by QRU Factory(TM). QRU simplifies the path to understanding the truth.

The Sleep Repair Principle

Why Rest Helps-and Why Sleep Matters

A QRU Companion Book for Understanding Sleep, Rest, Recovery, and Real-Life Sleep Challenges

Medical and Safety Disclaimer

This book is for general education only. It is not medical advice, diagnosis, or treatment. Sleep needs and sleep problems vary by person. Do not use this book to delay or replace care from a qualified healthcare professional.

Seek professional guidance if you have ongoing insomnia, excessive daytime sleepiness, loud snoring, gasping or choking during sleep, observed breathing pauses, restless legs, unusual nighttime behaviors, pregnancy-related sleep concerns, mental health concerns, medication questions, pain affecting sleep, or any medical condition that affects sleep.

If you feel dangerously sleepy while driving, operating machinery, caregiving, supervising children, working in healthcare, or performing any safety-sensitive task, stop as soon as it is safe and seek support. Drowsy driving can be dangerous.

Plain-Language Summary

Sleep is not wasted time. Sleep is active recovery time.

Rest is helpful because it lowers strain. A quiet break, prayer, meditation, stretching, calm breathing, or lying down may help you feel steadier. But rest does not fully replace sleep.

During sleep, your brain and body support memory, emotional balance, immune function, hormone regulation, metabolism, heart health, physical recovery, and brain maintenance. When sleep is repeatedly too short or poor quality, people often notice changes in focus, mood, reaction time, appetite, energy, and safety.

The QRU Sleep Repair Principle is:

Rest can pause the drain, but sleep protects the repair window.

This book will help you understand sleep, respect your biology, and build practical habits that support better rest and better sleep.

Table of Contents

1. [Introduction: The QRU Sleep Repair Principle](introduction-the-qru-sleep-repair-principle)
 2. [How to Use This Book](how-to-use-this-book)
 3. [Chapter 1: Sleep Is Not "Doing Nothing"](chapter-1-sleep-is-not-doing-nothing)
 4. [Chapter 2: The Body's Nightly Maintenance System](chapter-2-the-bodys-nightly-maintenance-system)
 5. [Chapter 3: How Much Sleep Do We Need?](chapter-3-how-much-sleep-do-we-need)
 6. [Chapter 4: What Happens When Sleep Runs Short](chapter-4-what-happens-when-sleep-runs-short)
 7. [Chapter 5: Rest Helps-but It Is Not Sleep](chapter-5-rest-helps-but-it-is-not-sleep)
 8. [Chapter 6: Building Better Sleep Hygiene](chapter-6-building-better-sleep-hygiene)
 9. [Chapter 7: Circadian Rhythm, Light, Caffeine, Alcohol, and Naps](chapter-7-circadian-rhythm-light-caffeine-alcohol-and-naps)
 10. [Chapter 8: Real-Life Sleep Barriers](chapter-8-real-life-sleep-barriers)
 11. [Chapter 9: When to Seek Help](chapter-9-when-to-seek-help)
 12. [Chapter 10: The One-Week QRU Sleep Reset](chapter-10-the-one-week-qru-sleep-reset)
 13. [Conclusion: Protect the Repair Window](conclusion-protect-the-repair-window)
 14. [Practical Tools](practical-tools)
 15. [References](references)
-

Introduction: The QRU Sleep Repair Principle

Sleep is often treated like a luxury, a weakness, or a reward we earn only after everything else is finished.

But sleep is not a bonus.

Sleep is a biological requirement.

The central idea of this book is simple:

Rest can pause the drain, but sleep protects the repair window.

Rest matters. A quiet break, a calm walk, a few minutes away from noise, prayer, meditation, or simply sitting still can reduce strain. Rest can help the body stop spending energy so aggressively.

But sleep is different.

Sleep is an active biological state. While you are asleep, your brain and body perform important work related to learning, memory, emotional regulation, immune function, metabolism,

hormones, cardiovascular health, and physical recovery.

Sleep does not magically solve every problem, and not every sleep difficulty can be fixed with habits alone. Many people face medical, work, caregiving, pregnancy, disability, pain, trauma, or environmental barriers. This book respects those realities.

The goal is not perfection.

The goal is understanding.

When you understand sleep, you can make better decisions about energy, recovery, safety, and support.

QRU Learning Outcomes

By the end of this book, you will be able to:

1. Explain why sleep is an active biological process.
2. Describe the difference between rest and sleep.
3. Identify general sleep duration recommendations by age group.
4. Explain why sleep needs vary by age, pregnancy, health status, activity level, schedule, and individual biology.
5. Recognize common effects of insufficient sleep on mood, thinking, reaction time, immune function, metabolism, and safety.
6. Use practical sleep hygiene strategies without treating them as a cure-all.
7. Explain how light, caffeine, alcohol, naps, stress, and schedules can affect sleep.
8. Identify sleep warning signs that may require professional evaluation.
9. Build a realistic one-week sleep reset plan.
10. Apply QRU tools to protect your sleep opportunity and rest needs.

How to Use This Book

You can read this book from beginning to end, or you can use it as a practical guide.

Each chapter includes:

- Key idea
- Plain-language summary
- Core teaching
- Practical exercise

- Memory reinforcement
- Chapter summary

For best results:

1. Read one chapter at a time.
2. Complete the exercise.
3. Notice patterns instead of judging yourself.
4. Use the tools at the end of the book.
5. Seek medical guidance when symptoms suggest a sleep disorder or health concern.

Chapter 1: Sleep Is Not "Doing Nothing"

Learning Outcomes

By the end of this chapter, you will be able to:

- Explain why sleep is biologically active.
- Identify major sleep stages.
- Describe why sleep quality matters, not only sleep quantity.

Key Idea

Sleep is not the absence of activity. Sleep is a different kind of activity.

Plain-Language Summary

When you sleep, your body is not "off." Your brain cycles through stages that support memory, emotion, physical recovery, and alertness the next day. Good sleep includes enough time and enough quality.

Sleep Is an Active State

From the outside, a sleeping person looks still. Inside, the body is busy.

During sleep, your body helps regulate:

- Brain activity
- Heart rate and blood pressure patterns
- Breathing

- Hormones
- Temperature
- Immune signaling
- Muscle recovery
- Memory processing
- Emotional regulation

Sleep is organized into repeating cycles. Each cycle usually lasts about 90 minutes, though this varies.

Major Sleep Stages

Sleep is often described in two broad categories:

1. Non-REM sleep
2. REM sleep

Non-REM Sleep

Non-REM sleep includes lighter sleep and deeper sleep.

Deep non-REM sleep is associated with physical restoration, immune processes, growth hormone release, and feeling physically refreshed. It is also involved in certain types of learning and memory.

REM Sleep

REM stands for rapid eye movement. REM sleep is often associated with vivid dreaming, emotional processing, and memory integration. During REM sleep, the brain is highly active, and most voluntary muscles are temporarily less active so the body does not act out dreams.

Sleep Quality Matters

Getting enough hours is important, but sleep quality also matters.

Poor sleep quality may include:

- Frequent awakenings
- Trouble falling back asleep
- Breathing interruptions
- Pain-related waking
- Restless legs
- Nightmares
- Irregular schedule

- Alcohol-disrupted sleep
- Environmental noise or light
- Waking unrefreshed despite enough time in bed

A person may spend eight hours in bed but still sleep poorly. Another person may sleep fewer hours for one night and recover, but repeated short sleep can become a problem.

Practical Exercise: Sleep Is Active Reflection

Answer these questions:

1. What do I usually assume sleep is: rest, shutdown, repair, escape, waste, or something else?
2. Do I treat sleep as a need or a reward?
3. What is one sign my body gives me when I have not slept enough?
4. What is one sign I slept well?

Memory Reinforcement

Remember the phrase:

Sleep is not "off." Sleep is organized recovery.

Chapter Summary

- Sleep is an active biological state.
 - Sleep includes non-REM and REM stages.
 - Different sleep stages support different functions.
 - Sleep quality matters as much as time in bed.
 - Repeated poor sleep can affect health, safety, and daily function.
-

Chapter 2: The Body's Nightly Maintenance System

Learning Outcomes

By the end of this chapter, you will be able to:

- Describe how sleep supports the brain and body.
- Explain sleep's relationship to memory, emotion, immunity, metabolism, and cardiovascular health.

- Discuss brain waste-clearance research accurately and cautiously.

Key Idea

Sleep supports maintenance across the whole body.

Plain-Language Summary

Sleep helps your brain and body reset for the next day. It supports memory, mood, immune defense, hormones, heart health, and physical recovery. Scientists are still studying exactly how all of these processes work.

Sleep and Learning

Sleep helps the brain process and organize information.

Research shows that sleep supports:

- Learning new information
- Strengthening memories
- Connecting ideas
- Problem-solving
- Skill learning
- Emotional processing

This does not mean studying and then sleeping guarantees perfect memory. It means sleep is one important part of learning.

Sleep and Emotional Regulation

Poor sleep can make emotions feel sharper and harder to manage.

After too little sleep, people may experience:

- Irritability
- Anxiety
- Lower patience
- More emotional reactivity
- Lower stress tolerance
- Difficulty reading social cues
- Reduced motivation

Sleep does not remove life stress. But it helps the brain respond to stress with more stability.

Sleep and Immune Function

Sleep and the immune system are connected. Sleep supports immune signaling and the body's response to infection and vaccination. Insufficient sleep may be linked with increased susceptibility to some infections and reduced immune response.

This does not mean sleep alone prevents illness. Health is shaped by many factors, including nutrition, vaccination, medical care, stress, environment, and underlying conditions.

Sleep and Metabolism

Sleep affects hormones involved in appetite, glucose regulation, and energy balance.

Short or disrupted sleep may be associated with:

- Increased hunger
- Cravings for high-energy foods
- Reduced insulin sensitivity
- Weight gain risk over time
- Higher risk of metabolic problems

These relationships are complex. Sleep is one factor among many, including genetics, food access, activity level, medications, stress, and medical conditions.

Sleep and Cardiovascular Health

Regularly insufficient or poor-quality sleep is associated with increased risk of high blood pressure, cardiovascular disease, and stroke. Sleep disorders such as obstructive sleep apnea can place extra strain on the cardiovascular system and should be evaluated.

Sleep and Brain Maintenance

You may hear that sleep "cleans the brain." This idea comes partly from research on the glymphatic system, a brain waste-clearance pathway studied in animals and humans.

Current evidence suggests sleep, especially deep sleep, may support waste clearance in the brain. Scientists are still studying the details, how this applies to different people, and what it means for long-term brain health.

A careful way to say it is:

Sleep appears to support important brain maintenance processes, including waste-clearance pathways, but research is ongoing.

Practical Exercise: Maintenance Map

Write one sentence for each area:

- When I sleep poorly, my memory feels?
- When I sleep poorly, my mood feels?
- When I sleep poorly, my food choices or appetite feel?
- When I sleep poorly, my body feels?
- When I sleep well, I notice?

Memory Reinforcement

Use the QRU "5 M's of Sleep Maintenance":

1. Memory
2. Mood
3. Metabolism
4. Muscles
5. Medical resilience

Chapter Summary

- Sleep supports learning, memory, and emotional regulation.
 - Sleep and immune function are connected.
 - Sleep influences metabolism and appetite regulation.
 - Sleep quality affects cardiovascular health.
 - Brain maintenance during sleep is an active area of research.
-

Chapter 3: How Much Sleep Do We Need?

Learning Outcomes

By the end of this chapter, you will be able to:

- Identify general sleep recommendations by age.
- Explain why individual sleep needs vary.

- Distinguish sleep opportunity from actual sleep.

Key Idea

Sleep needs change by age and life situation.

Plain-Language Summary

Most adults need at least seven hours of sleep per night. Children and teens need more. Some people need extra attention to sleep because of pregnancy, illness, caregiving, shift work, disability, stress, or demanding physical or mental work.

General Sleep Duration Recommendations

The following recommendations are based on guidance from organizations such as the CDC, American Academy of Sleep Medicine, American Academy of Pediatrics, and National Sleep Foundation.

| Age Group | Recommended Sleep Amount |

|---|---:|

| Newborns, 0-3 months | 14-17 hours per 24 hours |

| Infants, 4-12 months | 12-16 hours per 24 hours, including naps |

| Toddlers, 1-2 years | 11-14 hours per 24 hours, including naps |

| Preschoolers, 3-5 years | 10-13 hours per 24 hours, including naps |

| School-age children, 6-12 years | 9-12 hours per 24 hours |

| Teens, 13-18 years | 8-10 hours per 24 hours |

| Adults, 18-60 years | 7 or more hours per night |

| Adults, 61-64 years | 7-9 hours per night |

| Adults, 65+ years | 7-8 hours per night |

These are general ranges, not personal diagnoses.

Sleep Opportunity vs. Actual Sleep

Time in bed is not the same as time asleep.

If you get into bed at 10:00 p.m. and get out of bed at 6:00 a.m., you had an eight-hour sleep opportunity. But if it took an hour to fall asleep and you were awake several times, your actual sleep time may be much less.

Protecting sleep starts with protecting enough opportunity.

Why Sleep Needs Vary

Sleep needs may vary based on:

- Age
- Genetics
- Pregnancy
- Illness or recovery
- Physical activity
- Mental workload
- Stress
- Trauma
- Medications
- Pain
- Caregiving responsibilities
- Shift work
- Menopause
- Sleep disorders
- Environmental conditions
- Disability or chronic health conditions

Special Note for Teens

Teens often naturally feel sleepy later and wake later because of circadian rhythm changes during adolescence. Early school start times can make it difficult for teens to get the recommended 8-10 hours.

Teen sleep loss can affect:

- Mood
- Attention
- Learning
- Driving safety
- Sports performance
- Mental health
- Risk-taking behavior

Special Note for Older Adults

Older adults may experience lighter sleep, earlier sleep timing, more awakenings, pain,

medication effects, or medical conditions that affect sleep. However, poor sleep should not be dismissed as "just aging." Treatable sleep disorders and health issues can occur at any age.

Special Note for Pregnant People

Pregnancy can affect sleep through nausea, urination, discomfort, reflux, restless legs, anxiety, hormonal changes, and breathing changes. Loud snoring, breathing pauses, high blood pressure, severe insomnia, or extreme sleepiness during pregnancy should be discussed with a healthcare professional.

Practical Exercise: Sleep Opportunity Audit

For three days, record:

- Bedtime
- Time you estimate you fell asleep
- Wake time
- Number of awakenings
- Final out-of-bed time
- Estimated total sleep
- Energy level the next day

Then ask:

1. Am I giving myself enough sleep opportunity?
2. Is my sleep opportunity realistic for my life?
3. Do I need habit changes, support, medical evaluation, or schedule changes?

Memory Reinforcement

Remember:

You cannot sleep eight hours if you only allow six.

Chapter Summary

- Adults generally need at least seven hours of sleep.
- Children and teens need more.
- Sleep needs vary by age, body, health, and life circumstances.
- Sleep opportunity is the time you allow for sleep; actual sleep may be less.
- If you consistently wake unrefreshed despite enough time in bed, consider professional

guidance.

Chapter 4: What Happens When Sleep Runs Short

Learning Outcomes

By the end of this chapter, you will be able to:

- Identify short-term effects of sleep loss.
- Describe long-term risks associated with chronic insufficient sleep.
- Recognize safety concerns related to drowsiness.

Key Idea

Sleep loss affects the brain, body, mood, and safety.

Plain-Language Summary

Not sleeping enough can make it harder to think, react, regulate emotions, and stay safe. Long-term poor sleep is linked with higher risk for several health problems.

Short-Term Effects of Sleep Loss

After too little sleep, many people notice:

- Slower reaction time
- Reduced focus
- Memory lapses
- Irritability
- Lower motivation
- More mistakes
- Increased hunger
- Headaches
- Lower stress tolerance
- Microsleeps or nodding off
- Reduced coordination

Sleep Loss and Safety

Sleepiness can impair driving and work performance. Drowsy driving can be dangerous because reaction time, attention, and decision-making are reduced.

Warning signs include:

- Yawning repeatedly
- Heavy eyelids
- Drifting from your lane
- Missing exits or turns
- Not remembering the last few miles
- Nodding off
- Difficulty keeping your head up

If this happens, do not "push through." Stop as soon as it is safe.

Chronic Insufficient Sleep

Repeated insufficient sleep is associated with increased risk of:

- High blood pressure
- Heart disease
- Stroke
- Type 2 diabetes
- Obesity
- Depression
- Anxiety symptoms
- Impaired immune function
- Accidents and injuries
- Reduced quality of life

Association does not always mean direct cause in every individual, but the relationship between sleep and health is strong enough that sleep deserves protection.

Sleep Debt: A Helpful but Imperfect Idea

People often say they are "in sleep debt." This can be useful because it reminds us that repeated short sleep adds up.

But sleep is not a bank account that can be perfectly repaid. Sleeping longer on weekends may help some people feel better, but it may not fully erase the effects of chronic sleep restriction. Large weekend schedule shifts can also disrupt circadian rhythm.

Practical Exercise: Sleep Loss Pattern Check

Complete these sentences:

- When I sleep less than I need, I make more mistakes with?
- My mood changes by?
- My body feels?
- I am most at risk for drowsiness during?
- One safety step I can take is?

Memory Reinforcement

Use the QRU "Tired Brain Warning List":

- Slow
- Moody
- Hungry
- Forgetful
- Risky

Chapter Summary

- Sleep loss affects thinking, emotion, metabolism, and safety.
 - Drowsiness can be dangerous during driving, caregiving, and safety-sensitive work.
 - Chronic insufficient sleep is linked with several health risks.
 - Weekend catch-up sleep may help but does not fully replace consistent sleep.
-

Chapter 5: Rest Helps-but It Is Not Sleep

Learning Outcomes

By the end of this chapter, you will be able to:

- Define rest.
- Explain how rest differs from sleep.
- Use rest strategically without expecting it to replace sleep.

Key Idea

Rest reduces strain. Sleep supports deeper biological recovery.

Plain-Language Summary

Rest is valuable. It can calm your body, reduce stress, and help you get through the day. But rest does not provide all the same brain and body processes as sleep.

What Is Rest?

Rest may include:

- Sitting quietly
- Prayer
- Meditation
- Breathwork
- Gentle stretching
- A slow walk
- Listening to calming music
- Lying down
- Reducing stimulation
- Taking a break from screens
- Closing your eyes
- Quiet reflection

Rest can help lower arousal, reduce stress load, and conserve energy.

What Rest Can Do

Rest may help:

- Reduce tension
- Lower perceived stress
- Improve emotional steadiness
- Create a transition toward sleep
- Decrease sensory overload
- Support recovery during illness or fatigue
- Prevent burnout when used regularly

What Rest Cannot Fully Do

Rest does not fully replace sleep's role in:

- Sleep stage cycling
- REM sleep processes
- Deep sleep processes
- Memory consolidation
- Certain hormonal rhythms
- Full restoration of alertness
- Recovery from repeated sleep restriction

A quiet break is helpful. But if your body needs sleep, rest is not the same biological state.

The QRU Rest-and-Sleep Distinction

Use this phrase:

Rest is a pause. Sleep is a process.

Both matter.

Practical Exercise: Build a Rest Menu

Create a list of five rest options that do not require a major schedule change.

Examples:

1. Two minutes of slow breathing
2. Ten-minute walk without headphones
3. Prayer or quiet reflection
4. Stretching before bed
5. Sitting in natural light in the morning
6. Turning off notifications for 20 minutes
7. Lying down with eyes closed
8. Gentle music while doing a simple task

Choose one rest practice to try daily for one week.

Memory Reinforcement

Repeat:

Rest can lower the load. Sleep restores the system.

Chapter Summary

- Rest is valuable and should not be dismissed.
 - Rest reduces strain and supports emotional regulation.
 - Sleep is a distinct biological process.
 - Rest cannot fully replace sleep stage cycling or sleep-related recovery.
-

Chapter 6: Building Better Sleep Hygiene

Learning Outcomes

By the end of this chapter, you will be able to:

- Define sleep hygiene.
- Identify practical habits that support sleep.
- Understand the limits of sleep hygiene.

Key Idea

Sleep hygiene creates better conditions for sleep, but it is not a cure for every sleep problem.

Plain-Language Summary

Sleep hygiene means habits and environmental choices that support sleep. These habits can help, but if you have a sleep disorder, severe insomnia, pain, trauma, or medical condition, you may need professional care too.

What Sleep Hygiene Means

Sleep hygiene includes your routines, environment, and behaviors that affect sleep.

It is not about being perfect. It is about making sleep more likely.

Core Sleep Hygiene Practices

1. Keep a Consistent Wake Time

A consistent wake time helps anchor your circadian rhythm. If bedtime varies, wake time is often the best place to start.

2. Create a Wind-Down Routine

A wind-down routine tells the body that the day is ending.

Examples:

- Dim lights
- Take a warm shower
- Stretch gently
- Read something calming
- Pray or meditate
- Prepare clothes or lunch for tomorrow
- Write tomorrow's top three tasks
- Practice slow breathing

3. Make the Bedroom Sleep-Friendly

Aim for a bedroom that is:

- Dark
- Quiet
- Cool
- Comfortable
- Safe

Helpful tools may include:

- Blackout curtains
- Eye mask
- Earplugs
- White noise
- Fan
- Comfortable bedding
- Reduced clutter
- Phone outside reach

4. Limit Clock-Watching

Watching the clock can increase stress. If possible, turn the clock away.

5. Use the Bed Mostly for Sleep and Intimacy

For many people with insomnia, using the bed for work, scrolling, worrying, or watching intense content can train the brain to associate bed with wakefulness.

6. Get Out of Bed if You Cannot Sleep

If you are awake for a long time and becoming frustrated, consider getting out of bed and doing something quiet in low light until sleepy. This is a common behavioral strategy used in insomnia treatment.

This is general education, not personalized medical advice. People with mobility limitations, fall risk, caregiving demands, or medical conditions may need a modified approach.

The Limits of Sleep Hygiene

Sleep hygiene may not resolve:

- Chronic insomnia
- Sleep apnea
- Narcolepsy
- Restless legs syndrome
- Circadian rhythm disorders
- Depression or anxiety-related sleep disruption
- PTSD-related nightmares
- Pain-related sleep disruption
- Medication-related sleep problems
- Pregnancy-related sleep complications
- Menopause-related symptoms
- Unsafe housing or environmental noise
- Shift work sleep disorder

If sleep hygiene does not help, that does not mean you failed. It may mean you need a different level of support.

Practical Exercise: Bedroom Scan

Look at your sleep space and rate each area from 1 to 5:

| Sleep Factor | Rating 1-5 | One Improvement |

|---|---:|---|

| Darkness | | |

| Noise | | |

| Temperature | | |

| Comfort | | |

| Safety | | |

| Clutter/stimulation | | |

| Phone/screen control | | |

Choose one improvement this week.

Memory Reinforcement

Remember:

Make sleep easier to enter and harder to interrupt.

Chapter Summary

- Sleep hygiene supports sleep by improving routines and environment.
 - Consistent wake time is a powerful starting point.
 - A wind-down routine helps transition the body toward sleep.
 - Sleep hygiene is helpful but not a cure for all sleep disorders.
-

Chapter 7: Circadian Rhythm, Light, Caffeine, Alcohol, and Naps

Learning Outcomes

By the end of this chapter, you will be able to:

- Explain circadian rhythm in simple terms.
- Describe how light affects sleep timing.
- Use caffeine, alcohol, and naps more wisely.

Key Idea

Sleep is shaped by timing, chemistry, behavior, and environment.

Plain-Language Summary

Your body has an internal clock. Light, caffeine, alcohol, and naps can shift or disrupt sleep. Small timing changes can make a big difference.

Circadian Rhythm

Your circadian rhythm is an internal timing system that helps regulate sleep, alertness, body temperature, hormones, digestion, and other functions across roughly 24 hours.

Your rhythm is influenced by:

- Light exposure
- Meal timing
- Activity
- Social schedule
- Work schedule
- School schedule
- Travel
- Screen use
- Sleep timing

Light: The Main Clock Signal

Light is one of the strongest signals for the body clock.

Morning Light

Morning light helps signal daytime. Getting natural light soon after waking may support alertness and circadian alignment.

Evening Light

Bright light at night, especially close to the eyes, can delay sleep timing for some people. Screens can contribute, especially when they are bright, emotionally stimulating, or used in bed.

Practical steps:

- Get bright light earlier in the day.
- Dim lights in the evening.
- Reduce screen brightness at night.
- Use night mode if helpful.
- Avoid intense work or conflict on screens right before bed when possible.

Caffeine

Caffeine is a stimulant. It can improve alertness temporarily but may interfere with sleep.

Caffeine's half-life is often around 3-7 hours, but this varies by person due to genetics, pregnancy, medications, liver function, age, and sensitivity.

Common sources include:

- Coffee
- Tea
- Energy drinks
- Soda
- Chocolate
- Pre-workout supplements
- Some medications

General guidance:

- Avoid caffeine late in the day.
- Many people benefit from stopping caffeine at least 6-8 hours before bedtime.
- Sensitive people may need to stop earlier.
- Pregnant people should follow medical guidance on caffeine limits.

Alcohol

Alcohol may make people feel sleepy at first, but it can disrupt sleep quality.

Alcohol can:

- Fragment sleep
- Reduce REM sleep, especially earlier in the night
- Increase awakenings
- Worsen snoring
- Worsen obstructive sleep apnea
- Increase nighttime urination
- Reduce next-day alertness

Using alcohol as a sleep aid is not recommended. If you rely on alcohol to sleep or have difficulty reducing it, seek professional support.

Naps

Naps can help some people, especially when sleep opportunity is limited. But naps can also make nighttime sleep harder for others.

General nap guidance:

- Keep naps short, often 10-30 minutes.
- Nap earlier in the day, ideally before late afternoon.
- Avoid long or late naps if they worsen insomnia.
- Shift workers, caregivers, pregnant people, and people with medical needs may require

different nap strategies.

Practical Exercise: Timing Audit

Track for three days:

Factor	Day 1	Day 2	Day 3
--- --- --- ---			
Morning light time			
Last caffeine			
Alcohol? Amount/time			
Nap? Length/time			
Screen use in bed			
Bedtime			
Wake time			
Sleep quality 1-5			

Look for patterns.

Memory Reinforcement

Use the QRU "Clock Cues":

- Light early
- Dim late
- Caffeine with a cutoff
- Alcohol is not a sleep aid
- Nap with a plan

Chapter Summary

- Circadian rhythm helps regulate sleep timing.
- Morning light can support alertness and rhythm.
- Evening light and stimulating screen use may delay sleep.
- Caffeine can affect sleep for many hours.
- Alcohol can make sleep worse even if it causes drowsiness.
- Naps are useful when timed well but can interfere with nighttime sleep.

Chapter 8: Real-Life Sleep Barriers

Learning Outcomes

By the end of this chapter, you will be able to:

- Identify common life barriers to sleep.
- Apply realistic strategies for special situations.
- Use compassion instead of shame when sleep is difficult.

Key Idea

Sleep advice must fit real life.

Plain-Language Summary

Not everyone can simply "go to bed earlier." Work, caregiving, pregnancy, illness, stress, noise, pain, school, and safety issues can make sleep difficult. The goal is to improve what you can and get help where needed.

Shift Workers

Shift work can conflict with the body's natural circadian rhythm. This can cause sleepiness at work and difficulty sleeping during the day.

Helpful strategies may include:

- Keeping sleep and wake times as consistent as possible.
- Using bright light strategically during the work period.
- Wearing sunglasses on the commute home if morning light makes it harder to sleep.
- Making the daytime sleep environment dark, quiet, and cool.
- Using earplugs, white noise, or blackout curtains.
- Planning caffeine early in the shift, not near the end.
- Taking short planned naps when safe and appropriate.
- Avoiding drowsy driving after night shifts.

If severe sleepiness continues, consult a healthcare professional. Shift work sleep disorder is a recognized condition.

Caregivers and Parents

Caregivers may have interrupted sleep due to infants, children, elders, medical needs, disability support, or emergency responsibilities.

Helpful strategies:

- Sleep when there is a safe opportunity.
- Share night duties when possible.
- Use shifts with another adult if available.
- Prepare supplies before night.
- Protect one recovery block when possible.
- Accept practical help.
- Avoid judging yourself by routines designed for people without caregiving demands.

For infant sleep, follow safe sleep guidance from pediatric authorities.

Teens and Students

Teens often face biological late sleep timing plus early school demands.

Helpful strategies:

- Keep wake time as consistent as possible.
- Get morning light.
- Reduce late-night screen stimulation.
- Avoid caffeine late in the day.
- Plan homework earlier when possible.
- Protect sleep before exams; all-night studying can backfire.
- Avoid drowsy driving.

Older Adults

Older adults may experience earlier sleep timing, lighter sleep, more awakenings, medication effects, pain, or medical conditions.

Helpful strategies:

- Maintain daytime activity if medically safe.
- Get morning light.
- Review medications with a healthcare professional if sleep changes began after medication changes.
- Treat pain, urinary symptoms, breathing issues, or mood concerns.
- Avoid assuming poor sleep is unavoidable.

Pregnant People

Pregnancy can change sleep through discomfort, reflux, urination, leg cramps, restless legs, anxiety, and breathing changes.

Helpful strategies:

- Discuss sleep symptoms during prenatal care.
- Use pillows for comfort.
- Ask about safe positioning recommendations.
- Limit fluids close to bed if advised and safe.
- Discuss severe snoring, breathing pauses, high blood pressure, or extreme sleepiness promptly.
- Ask before using sleep supplements or medications.

People with Medical Conditions

Chronic pain, asthma, heart disease, reflux, diabetes, neurological conditions, mental health conditions, menopause symptoms, and many medications can affect sleep.

Helpful strategies:

- Track symptoms and sleep together.
- Ask whether medication timing affects sleep.
- Treat nighttime pain or reflux when possible.
- Discuss sleep changes with a clinician.
- Ask about screening for sleep apnea if symptoms match.

People in Unsafe or Noisy Environments

Some people cannot fully control light, noise, temperature, or safety.

Helpful strategies may include:

- Eye mask
- Earplugs
- White noise
- Fan
- Sleep headphones
- Door/window safety planning
- Community support
- Housing resources when available
- Rest breaks when full sleep is not possible

This is not a personal failure. Environment matters.

Practical Exercise: Barrier-to-Bridge Plan

Choose one barrier and one bridge.

| Barrier | What Makes It Hard? | One Realistic Bridge |

|---|---|---|

| Shift work | | |

| Caregiving | | |

| Stress | | |

| Pain | | |

| Noise | | |

| Screens | | |

| Caffeine | | |

| Medical symptoms | | |

Memory Reinforcement

Remember:

A sleep plan that ignores your life will not survive your life.

Chapter Summary

- Sleep barriers are real.
 - Shift workers, caregivers, teens, older adults, pregnant people, and people with medical conditions may need modified strategies.
 - Sleep support should be practical, compassionate, and safe.
 - Persistent symptoms may require professional care.
-

Chapter 9: When to Seek Help

Learning Outcomes

By the end of this chapter, you will be able to:

- Recognize signs of common sleep disorders.
- Identify when professional help is appropriate.
- Understand that sleep hygiene is not a substitute for treatment.

Key Idea

Some sleep problems need medical evaluation.

Plain-Language Summary

If you cannot sleep, cannot stay awake, snore loudly, stop breathing during sleep, or feel exhausted despite enough time in bed, get help. Many sleep disorders are treatable.

Seek Help Promptly If You Notice

- Loud, frequent snoring
- Breathing pauses during sleep
- Gasping or choking at night
- Excessive daytime sleepiness
- Falling asleep while driving or working
- Morning headaches with snoring or breathing symptoms
- High blood pressure with sleep apnea symptoms
- Restless legs that interfere with sleep
- Acting out dreams
- Sleepwalking that creates danger
- Panic, trauma symptoms, or nightmares affecting sleep
- Severe insomnia lasting weeks or months
- Sleep problems with depression, anxiety, mania, or suicidal thoughts
- Sudden sleep attacks
- Muscle weakness triggered by emotions
- Pregnancy-related severe snoring, breathing pauses, or extreme sleepiness
- Child or teen sleep problems affecting school, mood, behavior, or safety

If you have thoughts of self-harm or suicide, seek emergency help or contact local crisis support immediately.

Insomnia

Insomnia may include:

- Trouble falling asleep
- Trouble staying asleep
- Waking too early
- Sleep dissatisfaction

- Daytime impairment

For chronic insomnia, Cognitive Behavioral Therapy for Insomnia, or CBT-I, is recommended as a first-line treatment by major medical organizations. CBT-I is different from general sleep tips. It is a structured treatment that may include stimulus control, sleep restriction therapy, cognitive strategies, relaxation skills, and sleep education.

Do not start sleep restriction therapy on your own if you have bipolar disorder, seizure disorder, untreated sleep apnea, high fall risk, severe daytime sleepiness, or safety-sensitive duties unless guided by a qualified clinician.

Obstructive Sleep Apnea

Obstructive sleep apnea occurs when breathing repeatedly becomes partially or completely blocked during sleep.

Possible signs include:

- Loud snoring
- Gasping or choking
- Witnessed pauses in breathing
- Morning headaches
- Dry mouth
- Daytime sleepiness
- High blood pressure
- Waking unrefreshed
- Nighttime urination
- Mood or concentration problems

Sleep apnea can occur in people of any body size, age, or gender. It is more common with certain risk factors, but it should not be dismissed based on appearance.

Treatment may include positive airway pressure therapy, oral appliances, positional therapy, weight-related interventions when appropriate, surgery in selected cases, or treatment of contributing conditions. A clinician can guide evaluation and treatment.

Restless Legs Syndrome

Restless legs syndrome can cause uncomfortable leg sensations and an urge to move, often worse at rest and in the evening. It can disrupt sleep. Iron status, pregnancy, kidney disease, medications, and other factors may be involved. Medical evaluation can help.

Narcolepsy and Central Disorders of Hypersomnolence

If you experience overwhelming sleepiness despite enough sleep opportunity, sudden sleep attacks, vivid dream-like experiences as you fall asleep or wake, sleep paralysis, or sudden muscle weakness triggered by emotion, seek medical evaluation.

Medication and Supplement Caution

Sleep medications and supplements can have risks, interactions, and next-day effects. "Natural" does not always mean safe. Pregnant people, older adults, children, people with liver or kidney disease, people taking other medications, and people with breathing disorders should be especially cautious.

Discuss sleep aids with a qualified healthcare professional.

Practical Exercise: Help-Seeking Checklist

Check any that apply:

- ☐ I often snore loudly.
- ☐ Someone has seen me stop breathing during sleep.
- ☐ I wake gasping or choking.
- ☐ I feel sleepy while driving.
- ☐ I fall asleep unintentionally during the day.
- ☐ I have insomnia lasting more than a few weeks.
- ☐ I wake unrefreshed despite enough time in bed.
- ☐ My legs feel uncomfortable at night and movement helps.
- ☐ I act out dreams or move dangerously in sleep.
- ☐ Sleep problems are affecting mood, work, school, relationships, or safety.
- ☐ I am pregnant and having severe sleep symptoms.
- ☐ I have a medical condition or medication that may affect sleep.

If you checked any item, consider discussing sleep with a healthcare professional.

Memory Reinforcement

Remember:

Habits help. Symptoms need attention.

Chapter Summary

- Many sleep disorders are treatable.
 - Loud snoring, breathing pauses, severe sleepiness, and chronic insomnia deserve evaluation.
 - CBT-I is a first-line treatment for chronic insomnia.
 - Sleep aids and supplements should be used cautiously.
 - Safety concerns require immediate action.
-

Chapter 10: The One-Week QRU Sleep Reset

Learning Outcomes

By the end of this chapter, you will be able to:

- Build a realistic one-week sleep-support plan.
- Track sleep without obsessing over it.
- Apply behavior changes safely and appropriately.

Key Idea

A reset is not a cure. It is a structured way to notice patterns and support sleep.

Plain-Language Summary

The one-week sleep reset helps you improve sleep conditions. It is not medical treatment. If you have symptoms of a sleep disorder, severe insomnia, pregnancy concerns, or safety risks, seek professional guidance.

Important Boundary

The QRU Sleep Reset is general education and self-reflection. It is not a medical treatment plan.

Do not use this reset as a substitute for professional care if you have:

- Severe or chronic insomnia
- Suspected sleep apnea
- Narcolepsy symptoms
- Restless legs symptoms
- Bipolar disorder or mania symptoms
- Seizure disorder
- Severe depression or suicidal thoughts

- PTSD-related sleep symptoms
- Pregnancy-related severe sleep problems
- Dangerous daytime sleepiness
- Safety-sensitive work affected by sleepiness
- Complex medical conditions

Modify all steps for your safety, ability, medical needs, work schedule, caregiving responsibilities, and environment.

The Reset Goal

The goal is to protect your sleep opportunity and reduce sleep disruptors.

You are not trying to force sleep. You are creating better conditions for sleep.

Day 1: Establish Your Wake Anchor

Choose a wake time you can keep for most of the week.

Action steps:

- Wake within the same 30-60 minute window if possible.
- Get light soon after waking.
- Avoid judging the night. Start with the morning.

Reflection:

- What wake time is realistic?
- What makes it hard?
- What support would help?

Day 2: Create a Caffeine Cutoff

Choose a caffeine cutoff at least 6-8 hours before bedtime, or earlier if you are sensitive.

Action steps:

- Track all caffeine sources.
- Move caffeine earlier.
- Replace late caffeine with water, herbal tea, or a non-caffeinated routine.

Reflection:

- Did I notice cravings, headaches, or fatigue?
- Do I need to reduce caffeine gradually?

Day 3: Build a Wind-Down Routine

Create a 20-30 minute routine.

Possible routine:

1. Dim lights.
2. Prepare for tomorrow.
3. Wash face or shower.
4. Stretch gently.
5. Write worries or tasks on paper.
6. Read, pray, meditate, or breathe slowly.

Reflection:

- Which step felt calming?
- Which step felt unrealistic?

Day 4: Protect the Bedroom

Choose one environmental improvement.

Options:

- Darken the room.
- Reduce noise.
- Lower temperature if safe.
- Move phone away from bed.
- Clear clutter near the bed.
- Use an eye mask.
- Use white noise.

Reflection:

- What interrupted my sleep most?
- What can I control?
- What support do I need?

Day 5: Adjust Light and Screens

Action steps:

- Get morning light.
- Reduce bright light late at night.

- Avoid intense screen content in bed.
- Use night mode or dim settings if screens are necessary.
- Create a "last scroll" time.

Reflection:

- Did screens delay bedtime?
- Was the issue light, content, habit, loneliness, or stress?

Day 6: Plan Rest Before You Crash

Add one daily rest break.

Options:

- 3 slow breaths between tasks
- 5 minutes of silence
- 10-minute walk
- Prayer or meditation
- Eyes closed rest
- Gentle stretching
- Quiet cup of tea
- Sensory break

Reflection:

- Did rest reduce evening exhaustion?
- What type of rest actually helped?

Day 7: Review and Choose Your Keepers

Review your week.

Ask:

1. What improved my sleep opportunity?
2. What improved my sleep quality?
3. What made no difference?
4. What was unrealistic?
5. What symptoms may need professional help?
6. What two habits will I keep?

The QRU Sleep Reset Tracker

Day	Wake Time	Bedtime	Estimated Sleep	Caffeine Cutoff	Rest Break	Screen Boundary	Sleep Quality 1-5	Notes
1								
2								
3								
4								
5								
6								
7								

Practical Exercise: Choose Two Keepers

After seven days, choose two habits to continue.

My two keepers are:

- 1.
- 2.

My one support need is:

-

My one medical or professional question, if any, is:

-

Memory Reinforcement

Remember:

Do not chase perfect sleep. Build repeatable support.

Chapter Summary

- The one-week reset is educational, not medical treatment.
- Start with wake time, light, caffeine, wind-down, environment, screens, and rest.
- Track patterns without obsessing.
- Keep what works.
- Seek help for persistent or concerning symptoms.

Conclusion: Protect the Repair Window

Sleep is not laziness.

Sleep is not weakness.

Sleep is not wasted time.

Sleep is a biological requirement that supports how humans think, feel, heal, learn, work, drive, parent, create, regulate emotions, and recover.

Rest matters too. Rest can soften the load. Rest can give the nervous system a break. Rest can help you pause before you collapse.

But rest is not the same as sleep.

The QRU Sleep Repair Principle remains:

Rest can pause the drain, but sleep protects the repair window.

You do not need a perfect life to respect sleep. You need awareness, practical support, and honest attention to symptoms that deserve care.

Protect your sleep opportunity when you can.

Use rest wisely when sleep is limited.

Ask for help when symptoms persist.

Respect your biology.

Respect your limits.

Respect your repair window.

Practical Tools

Tool 1: QRU Sleep Opportunity Calculator

Use this to estimate whether you are allowing enough time for sleep.

| Question | Answer |

|---|---|

| What time do I need to wake up? | |

| How many hours of sleep do I need? | |

| How long do I usually take to fall asleep? | |

| How much time do I need for wind-down? | |

| What is my target bedtime? | |

Example:

- Wake time: 6:30 a.m.
- Sleep need: 8 hours
- Time to fall asleep: 20 minutes
- Wind-down: 30 minutes
- Target start of wind-down: 9:40 p.m.
- Target lights-out: 10:10 p.m.

Tool 2: QRU Sleepiness Safety Check

Before driving or safety-sensitive work, ask:

- ☐ Did I sleep less than usual?
- ☐ Am I yawning repeatedly?
- ☐ Are my eyelids heavy?
- ☐ Am I drifting, nodding, or losing focus?
- ☐ Have I used alcohol, sedating medication, or other substances?
- ☐ Am I working overnight or after a long shift?
- ☐ Would I want someone in my condition driving my family?

If safety is in doubt, stop and seek a safer option.

Tool 3: QRU Wind-Down Builder

Choose one item from each column.

| Body | Mind | Environment |

|---|---|---|

| Warm shower | Write tomorrow's top 3 tasks | Dim lights |

| Gentle stretch | Prayer | Cool room |

| Slow breathing | Gratitude list | Reduce noise |

| Relaxed walk | Read calming material | Move phone away |

| Comfortable clothes | Journal worries | Prepare bedding |

My wind-down plan:

- 1.
- 2.
- 3.

Tool 4: QRU Caffeine Map

Source	Amount	Time Used	Keep, Reduce, or Move Earlier?
Coffee			
Tea			
Soda			
Energy drink			
Chocolate			
Medication/supplement			

My caffeine cutoff goal:

-

Tool 5: QRU Rest Menu

Choose rest before exhaustion becomes a crash.

Time Available	Rest Option
30 seconds	One slow breath cycle
2 minutes	Relax shoulders, unclench jaw, breathe
5 minutes	Sit quietly with no phone
10 minutes	Walk outside or stretch
15 minutes	Prayer, meditation, or calming music
20 minutes	Eyes-closed rest or short nap
30 minutes	Planned nap or full decompression break

My top three rest options:

- 1.
- 2.
- 3.

Tool 6: QRU Sleep Symptom Notes for a Healthcare Visit

Bring this information to a healthcare professional if sleep is a concern.

Question	Notes
What is the main sleep problem?	
When did it start?	
How many nights per week?	

- | Bedtime and wake time? | |
- | Snoring, gasping, or breathing pauses? | |
- | Daytime sleepiness? | |
- | Drowsy driving or safety risks? | |
- | Restless legs? | |
- | Nightmares or acting out dreams? | |
- | Pain, reflux, urination, hot flashes, or other symptoms? | |
- | Medications, supplements, caffeine, alcohol? | |
- | Pregnancy, menopause, shift work, caregiving, stress? | |
- | Mental health symptoms? | |
- | What have you already tried? | |

Tool 7: QRU Sleep Support Checklist

Choose one action from each category.

Rhythm

- ☐ Keep a consistent wake time.
- ☐ Get morning light.
- ☐ Avoid large weekend schedule swings when possible.

Environment

- ☐ Make the room darker.
- ☐ Reduce noise.
- ☐ Adjust temperature.
- ☐ Improve comfort.
- ☐ Move phone away from bed.

Chemistry

- ☐ Set a caffeine cutoff.
- ☐ Avoid alcohol as a sleep aid.
- ☐ Review medications with a professional if needed.

Mind

- ☐ Write worries down before bed.
- ☐ Use prayer, meditation, or breathing.
- ☐ Plan tomorrow earlier in the evening.

Body

- ☐ Add daytime movement if medically safe.

- [] Stretch gently.
- [] Address pain, reflux, or breathing symptoms with professional help.

Safety

- [] Do not drive while dangerously sleepy.
- [] Plan transportation after night shifts if needed.
- [] Seek help for severe sleepiness.

References

American Academy of Pediatrics. (2016). American Academy of Pediatrics supports childhood sleep guidelines. <https://www.aap.org>

American Academy of Sleep Medicine. (2014). International classification of sleep disorders, third edition. American Academy of Sleep Medicine.

American Academy of Sleep Medicine. (2021). Sleep education and sleep duration recommendations. <https://aasm.org>

Centers for Disease Control and Prevention. (2022). How much sleep do I need? <https://www.cdc.gov/sleep/aboutsleep/howmuchsleep.html>

Centers for Disease Control and Prevention. (2024). Sleep and sleep disorders. <https://www.cdc.gov/sleep/>

Irish, L. A., Kline, C. E., Gunn, H. E., Buysse, D. J., & Hall, M. H. (2015). The role of sleep hygiene in promoting public health: A review of empirical evidence. *Sleep Medicine Reviews*, 22, 23-36. <https://doi.org/10.1016/j.smr.2014.10.001>

Krause, A. J., Simon, E. B., Mander, B. A., Greer, S. M., Saletin, J. M., Goldstein-Piekarski, A. N., & Walker, M. P. (2017). The sleep-deprived human brain. *Nature Reviews Neuroscience*, 18, 404-418. <https://doi.org/10.1038/nrn.2017.55>

Lo, J. C., Groeger, J. A., Santhi, N., Arbon, E. L., Lazar, A. S., Hasan, S., von Schantz, M., Archer, S. N., & Dijk, D. J. (2012). Effects of partial and acute total sleep deprivation on performance across cognitive domains. *PLoS ONE*, 7(9), e45987. <https://doi.org/10.1371/journal.pone.0045987>

Medic, G., Wille, M., & Hemels, M. E. H. (2017). Short- and long-term health consequences of sleep disruption. *Nature and Science of Sleep*, 9, 151-161. <https://doi.org/10.2147/NSS.S134864>

National Heart, Lung, and Blood Institute. (2022). Sleep deprivation and deficiency. National Institutes of Health. <https://www.nhlbi.nih.gov/health/sleep-deprivation>

National Heart, Lung, and Blood Institute. (2023). Sleep apnea. National Institutes of Health. <https://www.nhlbi.nih.gov/health/sleep-apnea>

National Institute of Neurological Disorders and Stroke. (2023). Brain basics: Understanding sleep. National Institutes of Health. <https://www.ninds.nih.gov/health-information/public-education/brain-basics/brain-basics-understanding-sleep>

National Sleep Foundation. (2015). National Sleep Foundation recommends new sleep times. *Sleep Health*, 1(1), 40-43. <https://doi.org/10.1016/j.sleh.2014.12.010>

Ramar, K., Malhotra, R. K., Carden, K. A., Martin, J. L., Abbasi-Feinberg, F., Aurora, R. N., Kapur, V. K., Olson, E. J., Rosen, C. L., Rowley, J. A., Shelgikar, A. V., & Trotti, L. M. (2021). Sleep is essential to health: An American Academy of Sleep Medicine position statement. *Journal of Clinical Sleep Medicine*, 17(10), 2115-2119. <https://doi.org/10.5664/jcsm.9476>

Sateia, M. J., Buysse, D. J., Krystal, A. D., Neubauer, D. N., & Heald, J. L. (2017). Clinical practice guideline for the pharmacologic treatment of chronic insomnia in adults. *Journal of Clinical Sleep Medicine*, 13(2), 307-349. <https://doi.org/10.5664/jcsm.6470>

Trauer, J. M., Qian, M. Y., Doyle, J. S., Rajaratnam, S. M. W., & Cunnington, D. (2015). Cognitive behavioral therapy for chronic insomnia: A systematic review and meta-analysis. *Annals of Internal Medicine*, 163(3), 191-204. <https://doi.org/10.7326/M14-2841>

Troxel, W. M. (2010). It's more than sex: Exploring the dyadic nature of sleep and implications for health. *Psychosomatic Medicine*, 72(6), 578-586. <https://doi.org/10.1097/PSY.0b013e3181de7ff8>

Watson, N. F., Badr, M. S., Belenky, G., Bliwise, D. L., Buxton, O. M., Buysse, D., Dinges, D. F., Gangwisch, J., Grandner, M. A., Kushida, C., Malhotra, R. K., Martin, J. L., Patel, S. R., Quan, S. F., & Tasali, E. (2015). Recommended amount of sleep for a healthy adult: A joint consensus statement of the American Academy of Sleep Medicine and Sleep Research Society. *Sleep*, 38(6), 843-844. <https://doi.org/10.5665/sleep.4716>

World Health Organization. (2020). Basic documents and health promotion resources. <https://www.who.int>

Xie, L., Kang, H., Xu, Q., Chen, M. J., Liao, Y., Thiyagarajan, M., O'Donnell, J., Christensen, D. J., Nicholson, C., Iliff, J. J., Takano, T., Deane, R., & Nedergaard, M. (2013). Sleep drives metabolite clearance from the adult brain. *Science*, 342(6156), 373-377. <https://doi.org/10.1126/science.1241224>

Continue your understanding at QRU:

