

2nd round:

Kuantum Metroloji

Metrology is the study of measurements.

Metrology is the study of measurements.

Quantum metrology uses quantum effects, i.e., entanglement, to make more precise measurements.

Metrology is the study of measurements.

Quantum metrology uses quantum effects, i.e., entanglement, to make more precise measurements.

More information = A high sensitivity of measurement.

(Quantum) Optical Coherence Tomography

(Quantum) Optical Coherence Tomography

PHYSICAL REVIEW A, VOLUME 65, 053817

Quantum-optical coherence tomography with dispersion cancellation

Ayman F. Abouraddy, Magued B. Nasr, Bahaa E. A. Saleh, Alexander V. Sergienko, and Malvin C. Teich*
*Quantum Imaging Laboratory,[†] Departments of Electrical & Computer Engineering and Physics, Boston University,
Boston, Massachusetts 02215*

(Received 27 November 2001; published 8 May 2002)

(Quantum) Optical Coherence Tomography

PHYSICAL REVIEW A, VOLUME 65, 053817

Quantum-optical coherence tomography with dispersion cancellation

A. Saleh, Alexander V. Sergienko, and Malvin C. Teich*
Electrical & Computer Engineering and Physics, Boston University,
Boston, Massachusetts 02215
(Received 12 October 2001; published 8 May 2002)




FIG. 1. Setup for optical coherence tomography (OCT). BS stands for beam splitter, D is a detector, and τ is a temporal delay introduced by moving the reference mirror.

(Quantum) Optical Coherence Tomography

PHYSICAL REVIEW A, VOLUME 65, 053817

Quantum-optical coherence tomography with dispersion cancellation




FIG. 1. Setup for optical coherence tomography (OCT). BS stands for beam splitter, D is a detector, and τ is a temporal delay introduced by moving the reference mirror.




FIG. 2. Setup for quantum-optical coherence tomography (QOCT). BS stands for beam splitter and τ is a temporal delay. D_1 and D_2 are single-photon-counting detectors that feed a coincidence circuit.

Advances in quantum metrology

Vittorio Giovannetti^{1*}, Seth Lloyd² and Lorenzo Maccone³

The statistical error in any estimation can be reduced by repeating the measurement and averaging the results. The central limit theorem implies that the reduction is proportional to the square root of the number of repetitions. Quantum metrology is the use of quantum techniques such as entanglement to yield higher statistical precision than purely classical approaches. In this Review, we analyse some of the most promising recent developments of this research field and point out some of the new experiments. We then look at one of the major new trends of the field: analyses of the effects of noise and experimental imperfections.

Quantum-Enhanced Measurements: Beating the Standard Quantum Limit

Vittorio Giovannetti,¹ Seth Lloyd,^{2*} Lorenzo Maccone³

Quantum mechanics, through the Heisenberg uncertainty principle, imposes limits on the precision of measurement. Conventional measurement techniques typically fail to reach these limits. Conventional bounds to the precision of measurements such as the shot noise limit or the standard quantum limit are not as fundamental as the Heisenberg limits and can be beaten using quantum strategies that employ "quantum tricks" such as squeezing and entanglement.

Measurement is a physical process, and the accuracy to which measurements can be performed is governed by the laws of phy-

sics. In particular, the behavior of systems at small scales is governed by the laws of quantum mechanics, which place limits on the accuracy to which measurements can be performed. These limits to accuracy take two forms. First, the Heisenberg uncertainty relation (I) imposes an intrinsic uncertainty on the values of measurement results of complementary observables such as position and momentum, or the different components of the angular momentum of a rotating object (Fig. 1). Second, every measurement apparatus is itself a quantum system: As a result, the uncertainty relations together with

other quantum constraints on the speed of evolution [such as the Margolus-Levitin theorem (2)] impose limits on how accurately we can measure quantities, given the amount of physical resources, such as energy, at hand to perform the measurement.

One important consequence of the physical nature of measurement is the so-called quantum back action: The extraction of information from a system can give rise to a feedback effect in which the system configuration after the measurement is determined by the measurement outcome. For example, the most extreme case (the so-called von Neumann or projective measurement) produces a complete determination of the post-measurement state. When performing successive measurements, quantum back action can be detrimental, because earlier measurements can negatively influence successive ones. A common strategy to get around the negative effect of back action

¹National Enterprise for nanoScience and nanoTechnology—Istituto Nazionale per la Fisica della Materia and Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126, Pisa, Italy. ²Research Laboratory of Electronics and Department of Mechanical Engineering, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA. ³Quantum Information Theory Group Dipartimento di Fisica "A. Volta," Università di Pavia, via A. Bassi 6 I-27100, Pavia, Italy.

*To whom correspondence should be addressed.
E-mail: slloyd@mit.edu




Characterizing Nonclassical Correlations via Local Quantum Uncertainty

Davide Girolami,¹ Tommaso Tufarelli,² and Gerardo Adesso¹

¹*School of Mathematical Sciences, The University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom*

²*Blackett Laboratory, QOLS, Imperial College London, London SW7 2BW, United Kingdom*

(Received 14 December 2012; revised manuscript received 1 April 2013; published 13 June 2013)






FIG. 1. **Conceptual scheme of a parameter estimation.** An initial probe is prepared (red box) in a state ρ_0 (eventually, from an initial state ρ_{in}). Then, it interacts with the unknown parameter λ through an evolution U_λ (green box). The state ρ_λ encoding the information on λ is measured by a POVM E_x (blue box) generating outcome x . Based on the outcomes x , a suitable estimator provides an estimate $\Lambda(x)$ of the parameter λ .




FIG. 1. **Conceptual scheme of a parameter estimation.** An initial probe is prepared (red box) in a state ρ_0 (eventually, from an initial state ρ_{in}). Then, it interacts with the unknown parameter λ through an evolution U_λ (green box). The state ρ_λ encoding the information on λ is measured by a POVM E_x (blue box) generating outcome x . Based on the outcomes x , a suitable estimator provides an estimate $\Lambda(x)$ of the parameter λ .




FIG. 2. **Conceptual scheme of a parallel parameter estimation.** The measurements here considered are separable. Indeed, employing entanglement in the measurement process does not allow to obtain better performances than the optimal separable strategy. Conversely, state preparation can lead to quantum enhancement by exploiting entanglement between probes^[9].

Photonic Quantum Metrology

Emanuele Polino,^{1, [a]} Mauro Valeri,^{1, [a]} Nicolò Spagnolo,¹ and Fabio Sciarrino^{1,2, [b]}

¹Dipartimento di Fisica, Sapienza Università di Roma, Piazzale Aldo Moro 5, I-00185 Roma, Italy

²Consiglio Nazionale delle Ricerche, Istituto dei Sistemi Complessi (CNR-ISC), Via dei Taurini 19, 00185 Roma, Italy

(Dated: 13 March 2020)

arXiv: 2002.05821



Resource Theory of Superposition

T. Theurer,¹ N. Killoran,^{1,2} D. Egloff,¹ and M. B. Plenio¹

¹*Institut für Theoretische Physik, Albert-Einstein-Allee 11, Universität Ulm, 89069 Ulm, Germany*

²*Department of Electrical and Computer Engineering, University of Toronto, Toronto, Canada*

(Received 10 April 2017; revised manuscript received 26 June 2017; published 5 December 2017)

.....



Resource Theory of Superposition

T. Theurer,¹ N. Killoran,^{1,2} D. Egloff,¹ and M. B. Plenio¹

¹*Institut für Theoretische Physik, Albert-Einstein-Allee 11, Universität Ulm, 89069 Ulm, Germany*

²*Department of Electrical and Computer Engineering, University of Toronto, Toronto, Canada*

(Received 10 April 2017; revised manuscript received 26 June 2017; published 5 December 2017)

PHYSICAL REVIEW A **103**, 032416 (2021)

Resource theory of superposition: State transformations

Gökhan Torun^{1,*}, Hüseyin Talha Şenyaşa^{2,†} and Ali Yıldız^{2,‡}

¹*Department of Physics, Boğaziçi University, 34342 Bebek, İstanbul, Turkey*

²*Department of Physics, İstanbul Technical University, 34469 Maslak, İstanbul, Turkey*



(Received 24 August 2020; accepted 2 March 2021; published 16 March 2021)

Channel Discrimination Task—a branch of quantum metrology

Channel Discrimination Task—a branch of quantum metrology

“Assume we have two players, say Alice and Bob. Alice performs a selective quantum operation (which is known to Bob) with outcomes $n = 0, 1, \dots, d < \infty$ on a state she received from Bob. If the result is $n = 0$, they start a new turn and Bob has to hand in a new state. In case the result was $n \neq 0$ she returns the post-measurement state to Bob, who is allowed to apply an arbitrary quantum operation on it. Then he has two choices: he either tells Alice his guess about the outcome n or he asks for a new turn. He has lost immediately if he gives a wrong answer and he wins if his answer is correct.” (PRL 119, 230401 (2017)).

Channel Discrimination Task—a branch of quantum metrology

“Assume we have two players, say Alice and Bob. Alice performs a selective quantum operation (which is known to Bob) with outcomes $n = 0, 1, \dots, d < \infty$ on a state she received from Bob. If the result is $n = 0$, they start a new turn and Bob has to hand in a new state. In case the result was $n \neq 0$ she returns the post-measurement state to Bob, who is allowed to apply an arbitrary quantum operation on it. Then he has two choices: he either tells Alice his guess about the outcome n or he asks for a new turn. He has lost immediately if he gives a wrong answer and he wins if his answer is correct.” (PRL 119, 230401 (2017)).

Identify outcomes $n = 1, \dots, d$ with the free Kraus operators

$$K_n = \sqrt{\frac{p}{d}} \sum_{j=1}^d e^{\frac{2\pi i j n}{d}} |c_j\rangle \langle c_j^\perp| \quad \left[p \in (0, 1], \sum_n K_n^\dagger K_n = p \sum_j |c_j^\perp\rangle \langle c_j^\perp| \leq \mathbb{I} \right]$$

1st case: Bob can hand in only free states—superposition-free states: $\rho_f = \sum_j |c_j\rangle\langle c_j|$

$$\left[K_n = \sqrt{\frac{p}{d}} \sum_{j=1}^d e^{\frac{2\pi i j n}{d}} |c_j\rangle\langle c_j^\perp| \right] \quad \rho_n = \frac{K_n \rho_f K_n^\dagger}{\text{tr}(K_n \rho_f K_n^\dagger)} = \rho_f \quad \left[p_n = \text{tr}(K_n \rho_f K_n^\dagger) = p/d \right]$$

The best choice is to make a random guess; since $p \leq 1$, Bob will lose with certainty for d against infinity.

2nd case: Bob can hand in only resource states—superposition states: $|\psi\rangle = \frac{1}{N} \sum_j |c_j\rangle$

$$|\psi_n\rangle = \frac{1}{N} \sum_k e^{\frac{2\pi i k n}{d}} |c_k\rangle$$

2nd case: Bob can hand in only resource states—superposition states: $|\psi\rangle = \frac{1}{N} \sum_j |c_j\rangle$

$$|\psi_n\rangle = \frac{1}{N} \sum_k e^{\frac{2\pi i k n}{d}} |c_k\rangle$$

These states are linearly independent. Since the free states ($|c_k\rangle$) are linearly independent,

$$\sum_n x_n |\psi_n\rangle = 0 \Rightarrow \sum_n x_n e^{\frac{2\pi i k n}{d}} = 0 \quad \forall k$$

2nd case: Bob can hand in only resource states—superposition states: $|\psi\rangle = \frac{1}{N} \sum_j |c_j\rangle$

$$|\psi_n\rangle = \frac{1}{N} \sum_k e^{\frac{2\pi i k n}{d}} |c_k\rangle$$

These states are linearly independent. Since the free states ($|c_k\rangle$) are linearly independent,

$$\sum_n x_n |\psi_n\rangle = 0 \Rightarrow \sum_n x_n e^{\frac{2\pi i k n}{d}} = 0 \quad \forall k \quad \left(\sum_n x_n u_n = 0 \quad \text{with} \quad u_n = \begin{bmatrix} e^{\frac{2\pi i 1 n}{d}} \\ e^{\frac{2\pi i 2 n}{d}} \\ \vdots \\ e^{\frac{2\pi i d n}{d}} \end{bmatrix} \right)$$

2nd case: Bob can hand in only resource states—superposition states: $|\psi\rangle = \frac{1}{N} \sum_j |c_j\rangle$

$$|\psi_n\rangle = \frac{1}{N} \sum_k e^{\frac{2\pi i k n}{d}} |c_k\rangle$$

These states are linearly independent. Since the free states ($|c_k\rangle$) are linearly independent,

$$\sum_n x_n |\psi_n\rangle = 0 \Rightarrow \sum_n x_n e^{\frac{2\pi i k n}{d}} = 0 \quad \forall k \quad \left(\sum_n x_n u_n = 0 \quad \text{with} \quad u_n = \begin{bmatrix} e^{\frac{2\pi i 1 n}{d}} \\ e^{\frac{2\pi i 2 n}{d}} \\ \vdots \\ e^{\frac{2\pi i d n}{d}} \end{bmatrix} \right)$$

Since $u_n^\dagger u_m = \delta_{nm} d$, the only solution is $x_n = 0$ for all n . Thus Bob can do unambiguous state discrimination on the states $\{|\psi_n\rangle\}$ and will, after enough repetitions, win with certainty (PRL 119, 230401 (2017)).

- M. Nielsen, and I. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, (2000).
- E.Chitambar and G.Gour, *Quantum Resource Theories*, *Rev. Mod. Phys*, **91**, 025001 (2019).
- R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Quantum Entanglement*, *Rev. Mod. Phys.* **81**, 865 (2009).
- M.Horodecki and J.Oppenheim, (*Quantumness in the context of*) *Resource Theories*, *Int. J. of M. Phys. B*, **27**, 1345019 (2013).
- H. Ollivier and W. H. Zurek, *Quantum Discord: A Measure of the Quantumness of Correlations*, *Phys. Rev. Lett.* **88**, 017901 (2001).

KUANTUM TEKNOLOJİLERİ & KAYNAK TEORİLERİ 〈 QSB | KU 〉

10 Nisan

**KUANTUM KAYNAK
TEORİLERİNE GİRİŞ**

14.00-14.50

Matematiksel Formalizm

14.50-15.40

**Kuantum Üst Üste Binme
Kuantum Eşevrelilik**

15.50-16.40

Kuantum Dolaşıklık

16.40-17.30

Kuantum Uyumsuzluk

17.30-18.15

QuTiP'e Giriş

11 Nisan

**KUANTUM TEKNOLOJİLERİNE
ÖRNEKLER**

14.00-14.50

**Kuantum Enformasyon
ve Hesaplama**

14.50-15.40

Kuantum Termodinamik

15.50-16.40

Kuantum Metroloji

16.40-17.30

Kuantum Biyoloji

Eğitimler Dr.Onur Pusuluk, Dr. Gökhan Torun ve Mohsen Izadyari tarafından verilecektir.

〈Q|Turkey〉

teşekkürler ...