

学号	姓名
23214321	陈宁浩
23214322	何昌烨
23214323	胡静静
23214324	黄项龙
23214326	刘润尧
23214329	宋珂
23214336	戴泳涛
23214338	杜冠男
23214339	段培明
23214345	黄瀚
23214346	黄樾
23214353	梁励
23214364	毛睿
23214369	钱甜奕
23214378	吴昊
23214383	徐博研
23214395	赵海洋
23214410	陈东
23214417	陈宁宁
23214421	陈腾跃
23214426	陈煜彦
23214427	崔铮浩
23214446	何鸿荣
23214449	何芷莹
23214452	洪桂航
23214460	黄泽林
23214466	赖柔成
23214474	李宏立
23214478	李茂锦
23214491	梁恒中
23214503	刘星宇
23214509	罗经周
23214534	苏达威
23214542	王辉
23214564	熊泽华
23214565	徐浩耀
23214573	杨坤业
23214576	杨子逸
23214578	杨沅旭
23214590	易钰淇
23214594	曾家洋
23214600	张珊
23214601	张晓逊
23214615	钟龙广
23214624	庄梓轩
23214625	邹国煌
23220055	李品律

Ex.1

(a) Since

$$\arg\left(\frac{i}{-2-2i}\right) = \arg i - \arg(-2-2i),$$

one value of $\arg\left(\frac{i}{-2-2i}\right)$ is $\frac{\pi}{2} - \left(-\frac{3\pi}{4}\right)$, or $\frac{5\pi}{4}$. Consequently, the principal value is

$$\frac{5\pi}{4} - 2\pi, \text{ or } -\frac{3\pi}{4}.$$

(b) Since

$$\arg(\sqrt{3}-i)^6 = 6\arg(\sqrt{3}-i),$$

one value of $\arg(\sqrt{3}-i)^6$ is $6\left(-\frac{\pi}{6}\right)$, or $-\pi$. So the principal value is $-\pi + 2\pi$, or π .

Ex.6

Proof : Let $z_1 = r_1 e^{i\theta_1}$, $z_2 = r_2 e^{i\theta_2}$, $\theta_1 = \text{Arg}(z_1)$, $\theta_2 = \text{Arg}(z_2)$. For $\text{Re}(z_1) > 0$, $\text{Re}(z_2) > 0$, then $-\frac{\pi}{2} < \theta_1, \theta_2 < \frac{\pi}{2}$.

$\text{Arg}(z_1 z_2) = \text{Arg}(r_1 r_2 e^{i(\theta_1 + \theta_2)})$, we have known that $-\pi < \theta_1 + \theta_2 < \pi$, then $\text{Arg}(z_1 z_2) = \theta_1 + \theta_2 = \text{Arg}(z_1) + \text{Arg}(z_2)$.

Ex.8

First of all, given two nonzero complex numbers z_1 and z_2 , suppose that there are complex numbers c_1 and c_2 such that $z_1 = c_1 c_2$ and $z_2 = c_1 \bar{c}_2$. Since

$$|z_1| = |c_1| |c_2| \quad \text{and} \quad |z_2| = |c_1| |\bar{c}_2| = |c_1| |c_2|,$$

it follows that $|z_1| = |z_2|$.

Suppose, on the other hand, that we know only that $|z_1| = |z_2|$. We may write

$$z_1 = r_1 \exp(i\theta_1) \quad \text{and} \quad z_2 = r_1 \exp(i\theta_2).$$

If we introduce the numbers

$$c_1 = r_1 \exp\left(i \frac{\theta_1 + \theta_2}{2}\right) \quad \text{and} \quad c_2 = \exp\left(i \frac{\theta_1 - \theta_2}{2}\right),$$

we find that

$$c_1 c_2 = r_1 \exp\left(i \frac{\theta_1 + \theta_2}{2}\right) \exp\left(i \frac{\theta_1 - \theta_2}{2}\right) = r_1 \exp(i\theta_1) = z_1$$

and

$$c_1 \bar{c}_2 = r_1 \exp\left(i \frac{\theta_1 + \theta_2}{2}\right) \exp\left(-i \frac{\theta_1 - \theta_2}{2}\right) = r_1 \exp \theta_2 = z_2.$$

That is,

$$z_1 = c_1 c_2 \quad \text{and} \quad z_2 = c_1 \bar{c}_2.$$

Ex.10

We know from de Moivre's formula that

$$(\cos \theta + i \sin \theta)^3 = \cos 3\theta + i \sin 3\theta,$$

or

$$\cos^3 \theta + 3 \cos^2 \theta (i \sin \theta) + 3 \cos \theta (i \sin \theta)^2 + (i \sin \theta)^3 = \cos 3\theta + i \sin 3\theta.$$

That is,

$$(\cos^3 \theta - 3 \cos \theta \sin^2 \theta) + i(3 \cos^2 \theta \sin \theta - \sin^3 \theta) = \cos 3\theta + i \sin 3\theta.$$

By equating real parts and then imaginary parts here, we arrive at the desired trigonometric identities:

$$(a) \cos 3\theta = \cos^3 \theta - 3 \cos \theta \sin^2 \theta; \quad (b) \sin 3\theta = 3 \cos^2 \theta \sin \theta - \sin^3 \theta.$$

page 29-31

Ex.2

(a) Since $-16 = 16 \exp[i(\pi + 2k\pi)]$ ($k = 0, \pm 1, \pm 2, \dots$), the needed roots are

$$(-16)^{1/4} = 2 \exp\left[i\left(\frac{\pi}{4} + \frac{k\pi}{2}\right)\right] \quad (k = 0, 1, 2, 3).$$

The principal root is

$$c_0 = 2e^{i\pi/4} = 2\left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4}\right) = 2\left(\frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}}\right) = \sqrt{2}(1 + i).$$

The other three roots are


$$c_1 = (2e^{i\pi/4})e^{i\pi/2} = c_0 i = \sqrt{2}(1 + i)i = -\sqrt{2}(1 - i),$$

$$c_2 = (2e^{i\pi/4})e^{i\pi} = -c_0 = -\sqrt{2}(1 + i),$$

and

$$c_3 = (2e^{i\pi/4})e^{i3\pi/2} = c_0(-i) = \sqrt{2}(1+i)(-i) = \sqrt{2}(1-i).$$

The four roots are shown below.



(b) First write $-8 - 8\sqrt{3}i = 16 \exp\left[i\left(-\frac{2\pi}{3} + 2k\pi\right)\right]$ ($k = 0, \pm 1, \pm 2, \dots$). Then

$$(-8 - 8\sqrt{3}i)^{1/4} = 2 \exp\left[i\left(-\frac{\pi}{6} + \frac{k\pi}{2}\right)\right] \quad (k = 0, 1, 2, 3).$$

The principal root is

$$c_0 = 2e^{-i\pi/6} = 2\left(\cos\frac{\pi}{6} - i\sin\frac{\pi}{6}\right) = 2\left(\frac{\sqrt{3}}{2} - \frac{i}{2}\right) = \sqrt{3} - i.$$


The others are

$$c_1 = (2e^{-i\pi/6})e^{i\pi/2} = c_0 i = 1 + \sqrt{3}i,$$

$$c_2 = (2e^{-i\pi/6})e^{i\pi} = -c_0 = -(\sqrt{3} - i),$$

$$c_3 = (2e^{-i\pi/6})e^{i3\pi/2} = c_0(-i) = -(1 + \sqrt{3}i).$$

These roots are all shown below.



Ex.4

The three cube roots of the number $z_0 = -4\sqrt{2} + 4\sqrt{2}i = 8\exp\left(i\frac{3\pi}{4}\right)$ are evidently

$$(z_0)^{1/3} = 2\exp\left[i\left(\frac{\pi}{4} + \frac{2k\pi}{3}\right)\right] \quad (k = 0, 1, 2).$$

In particular,

$$c_0 = 2\exp\left(i\frac{\pi}{4}\right) = \sqrt{2}(1+i).$$

With the aid of the number $\omega_3 = \frac{-1+\sqrt{3}i}{2}$, we obtain the other two roots:

$$c_1 = c_0\omega_3 = \sqrt{2}(1+i)\left(\frac{-1+\sqrt{3}i}{2}\right) = \frac{-(\sqrt{3}+1) + (\sqrt{3}-1)i}{\sqrt{2}},$$

$$c_2 = c_0\omega_3^2 = (c_0\omega_3)\omega_3 = \left[\frac{-(\sqrt{3}+1) + (\sqrt{3}-1)i}{\sqrt{2}}\right]\left(\frac{-1+\sqrt{3}i}{2}\right) = \frac{(\sqrt{3}-1) - (\sqrt{3}+1)i}{\sqrt{2}}.$$

Ex.5

(a) Let a denote any fixed real number. In order to find the two square roots of $a+i$ in exponential form, we write

$$A = |a+i| = \sqrt{a^2+1} \quad \text{and} \quad \alpha = \text{Arg}(a+i).$$

Since

$$a+i = A\exp[i(\alpha+2k\pi)] \quad (k = 0, \pm 1, \pm 2, \dots),$$

we see that

$$(a+i)^{1/2} = \sqrt{A}\exp\left[i\left(\frac{\alpha}{2} + k\pi\right)\right] \quad (k = 0, 1).$$

That is, the desired square roots are

$$\sqrt{A}e^{i\alpha/2} \quad \text{and} \quad \sqrt{A}e^{i\alpha/2}e^{i\pi} = -\sqrt{A}e^{i\alpha/2}.$$

(b) Since $a+i$ lies above the real axis, we know that $0 < \alpha < \pi$. Thus $0 < \frac{\alpha}{2} < \frac{\pi}{2}$, and this tells us that $\cos\left(\frac{\alpha}{2}\right) > 0$ and $\sin\left(\frac{\alpha}{2}\right) > 0$. Since $\cos\alpha = \frac{a}{A}$, it follows that

$$\cos\frac{\alpha}{2} = \sqrt{\frac{1+\cos\alpha}{2}} = \frac{1}{\sqrt{2}}\sqrt{1+\frac{a}{A}} = \frac{\sqrt{A+a}}{\sqrt{2}\sqrt{A}}$$

and

$$\sin\frac{\alpha}{2} = \sqrt{\frac{1-\cos\alpha}{2}} = \frac{1}{\sqrt{2}}\sqrt{1-\frac{a}{A}} = \frac{\sqrt{A-a}}{\sqrt{2}\sqrt{A}}.$$

Consequently,

$$\begin{aligned} \pm\sqrt{A}e^{i\alpha/2} &= \pm\sqrt{A}\left(\cos\frac{\alpha}{2} + i\sin\frac{\alpha}{2}\right) = \pm\sqrt{A}\left(\frac{\sqrt{A+a}}{\sqrt{2}\sqrt{A}} + i\frac{\sqrt{A-a}}{\sqrt{2}\sqrt{A}}\right) \\ &= \pm\frac{1}{\sqrt{2}}(\sqrt{A+a} + i\sqrt{A-a}). \end{aligned}$$

Ex.7

Let c be any n th root of unity other than unity itself. With the aid of the identity (see Exercise 9, Sec. 8),

$$1 + z + z^2 + \cdots + z^{n-1} = \frac{1 - z^n}{1 - z} \quad (z \neq 1),$$

we find that

$$1 + c + c^2 + \cdots + c^{n-1} = \frac{1 - c^n}{1 - c} = \frac{1 - 1}{1 - c} = 0.$$

Ex.9

Observe first that

$$(z^{1/m})^{-1} = \left[\sqrt[m]{r} \exp \frac{i(\theta + 2k\pi)}{m} \right]^{-1} = \frac{1}{\sqrt[m]{r}} \exp \frac{i(-\theta - 2k\pi)}{m} = \frac{1}{\sqrt[m]{r}} \exp \frac{i(-\theta)}{m} \exp \frac{i(-2k\pi)}{m}$$

and

$$(z^{-1})^{1/m} = \sqrt[m]{\frac{1}{r}} \exp \frac{i(-\theta + 2k\pi)}{m} = \frac{1}{\sqrt[m]{r}} \exp \frac{i(-\theta)}{m} \exp \frac{i(2k\pi)}{m},$$

where $k = 0, 1, 2, \dots, m-1$. Since the set

$$\exp \frac{i(-2k\pi)}{m} \quad (k = 0, 1, 2, \dots, m-1)$$

is the same as the set

$$\exp \frac{i(2k\pi)}{m} \quad (k = 0, 1, 2, \dots, m-1),$$

but in reverse order, we find that $(z^{1/m})^{-1} = (z^{-1})^{1/m}$.