

T: Triangle inequalities

Let $\alpha, \beta \in \mathbb{R}$. We then have that:

1. $|\alpha| + |\beta| \geq |\alpha + \beta|$
2. $||\alpha| - |\beta|| \leq |\alpha - \beta|$.

D: Supremum and infimum

Let $\alpha = \sup S$. Then:

1. $\forall s \in S; \alpha \geq s$
2. $\forall a \in \mathbb{R} : \forall s \in S; a \geq s;$
 $\quad \quad \quad \color{red}{a \geq \alpha}$

and similarly for infimum.

T: Approximation property

Consider bounded $E \subset \mathbb{R}$. Then:

$$\forall \epsilon > 0; \exists a \in E : \sup E - \epsilon < a \leq \sup E.$$

D: Completeness of \mathbb{R}

Every nonempty bounded subset of \mathbb{R} has an infimum and supremum.

T: Archimedean property

$$\forall a, b \in \mathbb{R}; a > 0; \exists n \in \mathbb{N} : na > b$$

D1.1: Nested intervals

A sequence of sets $(I_n)_{n \in \mathbb{N}}$ is nested if $I_1 \supset I_2 \supset I_3 \dots$.

T1.1: Nested interval property

Let $(I_n)_{n \in \mathbb{N}}$ be a sequence of nonempty, closed and bounded nested intervals. Then:

$$E = \bigcap_{n \in \mathbb{N}} I_n \neq \emptyset.$$

If $\lambda(I_n) \rightarrow 0$ then E contains one number, where λ denotes length.

T1.2

Let $E = [a, b]$ and that there exists an open collection of nested intervals $(I_\alpha)_{\alpha \in A}$ such that:

$$E \subset \bigcup_{\alpha \in A} I_\alpha.$$

Then $\exists \{\alpha_1, \alpha_2, \dots, \alpha_n\} \subset A$ such that:

$$E \subset I_{\alpha_1} \cup I_{\alpha_2} \cup \dots \cup I_{\alpha_n}.$$

D1.2: ϵ - N convergence

Let $\lim_{n \rightarrow \infty} x_n = a$. Then:

$$\forall \epsilon > 0; \exists N \in \mathbb{N} : \forall n \geq N \implies |x_n - a| < \epsilon.$$

D1.3: Cauchy sequences

The sequence (x_n) is Cauchy if:

$$\begin{aligned} \forall \epsilon > 0; \exists N \in \mathbb{N} : \forall n, m \geq N \\ \implies |x_n - x_m| < \epsilon. \end{aligned}$$

T1.3 and T1.4

$$\text{Cauchy} \iff \epsilon\text{-}N \text{ convergent.}$$

D1.4: Subsequences

The subsequence of $(x_n)_{n \in \mathbb{N}}$ is a sequence of form $(x_{n_k})_{k \in \mathbb{N}}$ and is a selection of the original sequence **taken in order**.

T1.5: Bolzano-Weierstrass

Every bounded real sequence has a convergent subsequence.

D1.5: Limit inferior and superior

Let (x_n) be a bounded real sequence. Then:

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \left(\sup_{k \geq n} x_k \right)$$

$$\liminf_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \left(\inf_{k \geq n} x_k \right).$$

T1.6

The real sequence (x_n) is convergent **iff**:

$$\limsup_{n \rightarrow \infty} x_n = \liminf_{n \rightarrow \infty} x_n.$$

D1.6: Convergence of infinite series

Series $S = \sum_{k=1}^{\infty} a_k$ is convergent if:

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n a_k < \infty.$$

Series S is **absolutely convergent** if

$$\sum_{k=1}^{\infty} |a_k| \text{ is also convergent.}$$

Otherwise S is conditionally convergent.

T1.7: Cauchy criterion for series

$$S = \sum_{k=1}^{\infty} a_k \text{ is convergent } \text{iff:}$$

$$\begin{aligned} \forall \epsilon > 0; \exists N \in \mathbb{N} : \forall m \geq n \geq N \\ \implies \left| \sum_{k=n+1}^m a_k \right| < \epsilon. \end{aligned}$$

T1.8

Let $S = \sum_{k=1}^{\infty} a_k$ be absolutely convergent. Let $z : \mathbb{N} \rightarrow \mathbb{N}$ be a bijection. Then:

$$\sum_{k=1}^{\infty} a_k = \sum_{k=1}^{\infty} a_{z(k)}.$$

T1.9: Riemann rearrangement

Let $S = \sum_{k=1}^{\infty} a_k$ be conditionally convergent. Then there exists rearrangements such that S can take on any value.

D1.7: Sequential continuity

Let $f : \text{dom}(f) \rightarrow \mathbb{R}$ where $\text{dom}(f) \subset \mathbb{R}$. f is continuous at $\alpha \in \text{dom}(f)$ if:

$$\begin{aligned} \forall (x_n)_{n \in \mathbb{N}} \subset \text{dom}(f) : \lim_{n \rightarrow \infty} x_n = \alpha \\ \implies \lim_{n \rightarrow \infty} f(x_n) = f(\alpha). \end{aligned}$$

T1.10

Let $\alpha \in \mathbb{R}$ and f, g continuous on D . Then $\alpha f, f + g, fg$ are continuous on D .

T1.11

Let f be continuous at $\alpha \in \mathbb{R}$ and g at $f(\alpha)$. Then $g \circ f$ is continuous at α .

D1.12: ϵ - δ continuity

Let $f : \text{dom}(f) \rightarrow \mathbb{R}$ where $\text{dom}(f) \subset \mathbb{R}$. Then f is continuous at $\alpha \in \text{dom}(f)$ if:

$$\begin{aligned} \forall \epsilon > 0; \exists \delta > 0 : |x - \alpha| < \delta \\ \implies |f(x) - f(\alpha)| < \epsilon. \end{aligned}$$

T1.13: Intermediate value theorem

Let f be continuous on $[a, b]$.

If $f(a)f(b) < 0$ then:

$$\exists c \in (a, b) : f(c) = 0.$$

T1.14: Extreme value theorem

Let f be continuous on $[a, b]$.

Then $\exists c, d \in [a, b]$ such that:

$$\begin{aligned} f(c) &= \inf\{f(x) : x \in [a, b]\} \\ f(d) &= \sup\{f(x) : x \in [a, b]\}. \end{aligned}$$

T: Mean value theorem

Let f be continuous on $[a, b]$ and differentiable on (a, b) . Then:

$$\exists c \in (a, b) : f'(c) = \frac{f(b) - f(a)}{b - a}.$$

D: Differentiability

f is differentiable at α if:

$$f'(\alpha) = \lim_{h \rightarrow 0} \frac{f(\alpha + h) - f(\alpha)}{h}.$$

T: Continuity test

f is continuous at α if:

$$\lim_{x \rightarrow \alpha} f(x) = f(\alpha)$$

where the limit from left/right must both exist and be equal to each other.

D2.1: Pointwise convergence

$f_n \rightarrow f$ pointwise on E if:

$$f(x) = \lim_{n \rightarrow \infty} f_n(x).$$

Here $f_n : E \rightarrow \mathbb{R}$ and:

$$\begin{aligned} \forall x \in E; \forall \epsilon > 0; \exists N \in \mathbb{N} : \forall n \geq N \\ \implies |f_n(x) - f(x)| < \epsilon. \end{aligned}$$

D2.2: Uniform convergence

$f_n \rightarrow f$ uniformly on E if:

$$\begin{aligned} \forall \epsilon > 0; \exists N \in \mathbb{N} : \forall n \geq N \text{ and } \forall x \in E \\ \implies |f_n(x) - f(x)| < \epsilon. \end{aligned}$$

P2.1

The following statements are equivalent.

1. $f_n \rightarrow f$ uniformly on E
2. $\lim_{n \rightarrow \infty} \sup_{x \in E} |f_n(x) - f(x)| = 0$
3. $\exists a_n \rightarrow 0$ s.t. $|f_n(x) - f(x)| \leq a_n$ for $\forall x \in E$.

T2.1

If f_n is continuous on E **and** $f_n \rightarrow f$ uniformly on E then f is continuous on E .

Remark

If f is not continuous on E then f_n cannot be uniform on E .

T2.5: Weierstrass M-test

Let $E \subset \mathbb{R}$ and $f_k : E \rightarrow \mathbb{R}$.

$$\exists M_k > 0 : \sum_{k=1}^{\infty} M_k < \infty.$$

If $\forall k \in \mathbb{N}$ and $\forall x \in E; |f_k(x)| \leq M_k$ then:

$$\sum_{k=1}^{\infty} f_k(x) \text{ converges uniformly on } E.$$

D: Power series

Let (a_n) be a real sequence and $c \in \mathbb{R}$. Then:

$$f_{PS}(x) = \sum_{n=0}^{\infty} a_n(x-c)^n$$

is a power series centered at c , with **radius of convergence**:

$$R = \sup\{r \geq 0 : (a_n r^n) \text{ is bounded}\}$$

where $R = \infty$ implying that series converges everywhere.

T3.1: Convergence of power series

Let $0 < R < \infty$. If $|x - c| < R$ then $f_{PS}(x)$ converges absolutely.

If $|x - c| > R$ then $f_{PS}(x)$ diverges.

T3.2: Continuity of power series

Let $0 < r < R$ where R is the radius of convergence of $f_{PS}(x)$.

Then for $|x - c| \leq r$, $f_{PS}(x)$ converges absolutely and uniformly to a continuous function $f(x)$.

L3.1

$\sum_{n=1}^{\infty} a_n(x-c)^n$ and $\sum_{n=1}^{\infty} n a_n(x-c)^{n-1}$ have the same radius of convergence.

T: Root and ratio tests

Let $S = \sum_{n=1}^{\infty} a_n$ and consider:

1. Ratio test: $\rho = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$
2. Root test: $\rho = \lim_{n \rightarrow \infty} |a_n|^{1/n}$.

Then:

•

T3.3

Let R be the radius of convergence of $f_{PS}(x)$. Then for $\forall x : |x - c| < R$, $f_{PS}(x)$ is **infinitely differentiable** and:

$$f_{PS}(x) = \sum_{n=0}^{\infty} a_n(x-c)^n$$

$$a_n = \frac{f^{(n)}(c)}{n!}.$$

T: Taylor's theorem**Elementary expansions**

- $E(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!}$
- $S(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$
- $C(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$

D: Characteristic functions

Let $E \subset \mathbb{R}$. The characteristic function is defined as a real function such that:

$$\chi_E(x) = \begin{cases} 1 & x \in E \\ 0 & \text{otherwise.} \end{cases}$$

D4.1 and D4.2: Step functions

The step function with respect to finite set $\{x_0, \dots, x_n\}$ for some $n \in \mathbb{N}$ is:

$$\phi(x) = \begin{cases} 0 & x < x_0 \text{ or } x > x_n \\ c_j & x \in (x_{j-1}, x_j); 1 \leq j \leq n \end{cases}$$

and its integral is defined as:

$$\int \phi = \sum_{j=1}^n c_j(x_j - x_{j-1}).$$

D4.3: Lebesgue integrable

$f : I \rightarrow \mathbb{R}$ is Lebesgue integrable on I if:

1. $\sum_{j=1}^{\infty} |c_j| \lambda(J_j) < \infty$
2. $\forall x \in I; f(x) = \sum_{j=1}^{\infty} |c_j| \chi_{J_j}(x) < \infty$

Here $c_j \in \mathbb{R}$, $J_i \subset I$ and is bounded for $j \in \{1, 2, 3, \dots\}$. Then:

$$\int_I f = \sum_{j=1}^{\infty} |c_j| \lambda(J_j).$$

T4.1**T4.2: Basic properties**

Let f, g be integrable on I and $\alpha, \beta \in \mathbb{R}$.

1. $\alpha f + \beta g$ is integrable on I and:

$$\int_I (\alpha f + \beta g) = \alpha \int_I f + \beta \int_I g.$$

2. If $f \geq g$ on I then $\int_I f \geq \int_I g$.
3. $|f|$ is integrable on I and:

$$\int_I |f| \geq \left| \int_I f \right|.$$

4. If f or g is bounded on I then fg is integrable on I .

5. If $f \geq 0$ and $\int_I f = 0$, then $\forall h$ such that $0 \leq h \leq f$ is also integrable on I .

T4.3