

Honours Differential Equations

Notes by Christopher Shen

Winter 2023

Contents

1	ODE systems	3
1.1	Integrating factors	3
1.2	Change of variables	3
1.3	Existence and uniqueness for IVPs	3
1.4	Homogeneous systems	4
1.5	Non-homogeneous systems	5
1.6	Critical points & linearisation	6
1.7	Stability of critical points	7
1.8	Lyapunov's theory and limit cycles	7
2	Fourier series	8
2.1	Real Fourier series	8
2.2	Complex Fourier series	10
2.3	Parseval's theorem	10
3	PDEs	11
3.1	Separation of variables	11
3.2	Heat equation	11
3.2.1	Standard boundary conditions	11
3.2.2	Fixed boundary temperatures	13
3.2.3	Insulated rod ends	14
3.3	Wave equation	15
3.4	Laplace's equation	16
3.4.1	Rectangular boundary conditions	16
3.4.2	Circular boundary conditions	17
4	Sturm-Liouville theory	19
4.1	Regular S-L problems	19
4.1.1	General 2nd order ODEs	20
4.1.2	Lagrange's identity	20
4.1.3	Series expansion	21
4.2	Non-homogeneous S-L problems	22
4.3	Singular S-L problems	23
4.4	S-L Series solutions	24
5	Laplace transforms	25

1 ODE systems

1.1 Integrating factors

Consider linear DE of form

$$y' + P(x)y = Q(x)$$

Our integrating factor for this DE is:

$$I(x) = \exp\left(\int P(x)dx\right)$$

Then our solution is

$$y(x) = \frac{1}{I(x)} \int Q(x)I(x)dx + \frac{\alpha}{I(x)}$$

where here α is a constant.

1.2 Change of variables

For higher order differential equations of form

$$y^{(n)} = F(y, y', \dots, y^{(n-1)}, t),$$

consider **change of variables** $x_{i+1} = y^{(i)}$ for $i \in \{0, 1, \dots, n-1\}$. Then take derivative with respect to time to obtain a first order ODE system, which takes the form:

$$x'_j = F_j(t, x_1, \dots, x_n)$$

for $j = 1, \dots, n$. We either immediately write this as a matrix system or linearise near a critical point.

1.3 Existence and uniqueness for IVPs

An initial value problem (**IVP**) is defined as

$$\frac{dx}{dt} = f(x, t)$$

for **initial** condition $x(t_0) = x_0$. A solution $x : I \rightarrow \mathbb{R}$ is a differentiable function that satisfies the IVP. Similarly for a first order system

$$x'_i = F_i(t, x_1, \dots, x_n)$$

to have a **unique** solution, F_i and $\frac{\partial F_i}{\partial x_j}$ must be continuous in a region. Here $i, j \in \{1, \dots, n\}$.

1.4 Homogeneous systems

Now consider $\mathbf{x}' = \mathbf{A}\mathbf{x}$, where \mathbf{A} is a $n \times n$ matrix.

Substituting $\mathbf{x} = e^{rt}\boldsymbol{\xi}$ results in an eigenvector problem:

$$(\mathbf{A} - r_i \mathbf{I}_n)\boldsymbol{\xi}^{(i)} = \mathbf{0}.$$

If eigenvalues repeat we try $\mathbf{x} = te^{rt}\boldsymbol{\xi} + e^{rt}\boldsymbol{\eta}$ which gives

$$(\mathbf{A} - r_i \mathbf{I}_n)\boldsymbol{\eta}^{(i)} = \boldsymbol{\xi}^{(i)}.$$

The **fundamental matrix** satisfies $\boldsymbol{\Psi}' = \mathbf{A}\boldsymbol{\Psi}$ and contains the basis of solutions.

$$\boldsymbol{\Psi}(t) = [\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(n)}]$$

for $\mathbf{x}^{(i)} = \exp\{(r_i t)\}\boldsymbol{\xi}^{(i)}$ and hence our general solution is $\mathbf{x} = \boldsymbol{\Psi}\mathbf{c}$.

1.5 Non-homogeneous systems

Consider non-homogeneous ODE system:

$$\mathbf{x}' = \mathbf{A}\mathbf{x} + \mathbf{g}.$$

There are a couple of different approaches we can take to solve such a system.

- **Change of basis**

Let $\mathbf{x} = \mathbf{T}\mathbf{y}$, where \mathbf{T} is our eigenvector matrix from diagonalisation. So $\mathbf{A} = \mathbf{T}\mathbf{D}\mathbf{T}^{-1}$, and after some algebra we obtain:

$$\mathbf{y}' = \mathbf{D}\mathbf{y} + \mathbf{T}^{-1}\mathbf{g}$$

which can be solved by integrating factors. Finally revert back to \mathbf{x} .

- **Variation of parameters**

So $\mathbf{x}_H = \Psi\mathbf{c}$ solves the $\mathbf{x}' = \mathbf{A}\mathbf{x}$, where \mathbf{c} is a constant vector.

We then assume that the solution to our non-homogeneous system takes the form:

$$\mathbf{x} = \Psi\mathbf{u}$$

for here $\mathbf{u} = \mathbf{u}(t)$. We then get $\Psi\mathbf{u}' = \mathbf{g}$, which can be solved by eliminating variables and integrating.

- **Method of undetermined coefficients**

Our non-homogeneous ODE system has solutions of form:

$$\mathbf{x} = \mathbf{x}_H + \mathbf{x}_p$$

Solving the homogeneous ODE gives us \mathbf{x}_H .

On the other hand we just need to find a **particular solution** \mathbf{x}_p that satisfies our non-homogeneous ODE. Then our solution is complete.

Whilst the fastest, this method is not guaranteed to work.

1.6 Critical points & linearisation

Consider non-linear ODE system

$$x' = F(x, y),$$

$$y' = G(x, y).$$

We define $\mathbf{x}^0 = \begin{bmatrix} x^0 \\ y^0 \end{bmatrix}$ as a **critical point** when $F(\mathbf{x}^0) = G(\mathbf{x}^0) = 0$.

Non-linear systems may then be linearised by Taylor expanding them around a critical point \mathbf{x}^0 , and discarding higher order terms.

i.e. let $\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}$ where $u_1 = x - x^0$ and $u_2 = y - y^0$.

$$\therefore u'_1 = x'$$

$$\approx F(x^0, y^0) + \left(\frac{\partial F}{\partial x} \right)_{x^0} (x - x^0) + \left(\frac{\partial F}{\partial y} \right)_{y^0} (y - y^0)$$

$$\therefore u'_2 = y'$$

$$\approx G(x^0, y^0) + \left(\frac{\partial G}{\partial x} \right)_{x^0} (x - x^0) + \left(\frac{\partial G}{\partial y} \right)_{y^0} (y - y^0)$$

Then we end up with the following linear system:

$$\mathbf{u}' = \mathbf{A}\mathbf{u}$$

where $\mathbf{A} = \begin{bmatrix} \partial F/\partial x & \partial F/\partial y \\ \partial G/\partial x & \partial G/\partial y \end{bmatrix}_{\mathbf{x}=\mathbf{x}^0}$ and is a 2×2 Jacobian matrix.

Our critical points \mathbf{x}^0 may also be classified:

Eigenvalues	Critical Points	Stability
$r_1 > r_2 > 0$	Node (source)	unstable
$r_1 < r_2 < 0$	Node (sink)	asympt. stable
$r_2 < 0 < r_1$	saddle	unstable
$r_1 = r_2 > 0$	Proper/Improper node	unstable
$r_1 = r_2 < 0$	Proper/Improper node	asympt. stable
$r_1, r_2 = \lambda \pm i\mu$ ($\lambda > 0$)	focus	unstable
$r_1, r_2 = \lambda \pm i\mu$ ($\lambda < 0$)	focus	asympt. stable
$r_1 = i\mu, r_2 = -i\mu$	center	stable

Linearisation preserves critical point behaviour **except** when eigenvalues are of $r = \pm i\mu$ form, for which then classification is unknown.

1.7 Stability of critical points

Stable critical points \mathbf{x}^0 : All solutions start and stay near \mathbf{x}^0 .

$$\forall \epsilon > 0; \exists \delta > 0; \forall \mathbf{x}_{\text{solution}} \text{ to } \mathbf{x}' = \mathbf{F}(\mathbf{x}, t): \\ |\mathbf{x}(0) - \mathbf{x}^0| < \delta \implies |\mathbf{x}(t) - \mathbf{x}^0| < \epsilon \text{ for } \forall t \geq 0$$

Attracting critical points \mathbf{x}^0 : All solutions tends to \mathbf{x}^0 .

$$\forall \delta > 0 : |\mathbf{x}(0) - \mathbf{x}^0| < \delta \implies \lim_{t \rightarrow \infty} \mathbf{x}(t) = \mathbf{x}^0$$

Asymptotically stable critical points \mathbf{x}^0 : Attracting **and** stable

1.8 Lyapunov's theory and limit cycles

In this section $\dot{\mathbf{x}}$ means its first time derivative. So consider:

$$\dot{x} = F(x, y),$$

$$\dot{y} = G(x, y)$$

defined in \mathbb{R}^2 . Let $\mathbf{x}^0 \in D$ be a critical point.

The function $E : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ is a Lyapunov function where $E(x^0, y^0) = 0$, whenever it exists. Note that the time derivative of E is:

$$\frac{dE}{dt} = \frac{\partial E}{\partial x} F + \frac{\partial E}{\partial y} G$$

- Let $E > 0$ for $\forall \mathbf{x} \neq \mathbf{x}^0$.
If $\frac{dE}{dt} \leq 0$ then \mathbf{x}^0 is stable.
If $\frac{dE}{dt} < 0$ then \mathbf{x}^0 is asymptotically stable.
- If every neighbourhood of \mathbf{x}^0 contains \mathbf{x}^* such that $E(\mathbf{x}^*) > 0$
and if $\frac{dE}{dt} > 0$ then \mathbf{x}^0 is unstable.

Now **limit cycles** are defined as periodic solutions such that at least one other **non-closed trajectory** approaches the limit cycle as $t \rightarrow \infty$.

2 Fourier series

2.1 Real Fourier series

Let $f(x)$ and $f'(x)$ be **piecewise continuous** in $[-L, L]$ with **period** $2L$.
i.e. $f(x) = f(x + 2L)$ for $\forall x$. Then the Fourier series for $f(x)$ is

$$f_{FS}(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right).$$

The **convergence** of our Fourier series depends on the continuity of $f(x)$:

- If $f(x)$ is continuous then $f_{FS}(x) = f(x)$.
- If $f(\alpha)$ is discontinuous then at point α we have

$$f_{FS}(\alpha) = \frac{f(\alpha^+) + f(\alpha^-)}{2}.$$

Note that $f(x)$ is continuous at α if $f(\alpha) = \lim_{x \rightarrow \alpha} f(x)$ and we define:

$$f(\alpha^-) = \lim_{x \rightarrow \alpha^-} f(x)$$

and

$$f(\alpha^+) = \lim_{x \rightarrow \alpha^+} f(x),$$

i.e. limits from left and right respectively. It is important to also note that the derivative of a Fourier series is **not necessarily convergent**.

Now consider $S_n = \sin \frac{n\pi x}{L}$ and $C_n = \cos \frac{n\pi x}{L}$. We then have the following **orthogonality relations**:

$$\langle S_n, S_m \rangle = \langle C_n, C_m \rangle = L\delta_{mn}$$

and

$$\langle S_n, C_m \rangle = 0$$

which are used to find a_n and b_n in $f_{FS}(x)$. So:

$$a_0 = \frac{1}{L} \int_{-L}^L f(x) dx,$$

$$a_n = \frac{1}{L} \int_{-L}^L \cos \frac{n\pi x}{L} f(x) dx$$

and

$$b_n = \frac{1}{L} \int_{-L}^L \sin \frac{n\pi x}{L} f(x) dx.$$

Note that δ_{mn} is the **Kronecker delta** and is defined as:

$$\delta_{mn} = \begin{cases} 1 & m = n \\ 0 & m \neq n \end{cases}$$

These trigonometric identities are useful for finding Fourier coefficients:

$$\cos A \cos B = \frac{1}{2} [\cos(A - B) + \cos(A + B)],$$

$$\sin A \sin B = \frac{1}{2} [\cos(A - B) - \cos(A + B)],$$

$$\sin A \cos B = \frac{1}{2} [\sin(A + B) + \sin(A - B)].$$

The **inner product** of two functions is also defined as:

$$\langle u(x), v(x) \rangle = \int_{-L}^L u(x)v(x)dx.$$

A shortcut is to note that:

- The Fourier series of **even** functions contains only cosines.
- The Fourier series of **odd** functions contains only sines.

Even functions are defined $f(-x) = f(x)$, and:

$$\int_{-L}^L f_{even} dx = 2 \int_0^L f_{even} dx.$$

Similarly **odd** functions are defined $f(-x) = -f(x)$, and:

$$\int_{-L}^L f_{odd} dx = 0.$$

2.2 Complex Fourier series

Our Fourier series may be rewritten as a **complex Fourier series**

$$f_{FS}(x) = \sum_{n=-\infty}^{\infty} c_n \exp\left(\frac{in\pi}{L}x\right)$$

using Euler's formula $e^{i\theta} = \sin \theta + i \cos \theta$. Then:

$$c_n = \begin{cases} (a_n - ib_n)/2 & n > 0 \\ (a_0)/2 & n = 0 \\ (a_n + ib_n)/2 & n < 0 \end{cases}$$

and therefore negative n values are included. By orthogonality:

$$c_n = \frac{1}{2L} \int_{-L}^L \exp\left(-\frac{in\pi}{L}x\right) f(x) dx$$

for $\forall n \in \mathbb{Z}$. Here we define the **inner product** for complex functions as

$$\langle f, g \rangle = \int_{\alpha}^{\beta} f^*(x) g(x) dx$$

where $f^*(x)$ is the complex conjugate of $f(x)$. Then:

$$\begin{aligned} \left\langle \exp\left(\frac{im\pi}{L}x\right), \exp\left(\frac{in\pi}{L}x\right) \right\rangle &= \int_{-L}^L \exp\left(-\frac{im\pi}{L}x\right) \exp\left(\frac{in\pi}{L}x\right) dx \\ &= 2L\delta_{mn} \end{aligned}$$

and since $f(x) = f_{FS}(x)$ we obtain our formula.

2.3 Parseval's theorem

Parseval's theorem states that given a periodic $f(x)$ with convergent Fourier series we have that

$$\begin{aligned} \langle f, f \rangle &= \int_{-L}^L |f(x)|^2 dx \\ &= 2L \sum_{n=-\infty}^{\infty} |c_n|^2 \\ &= L \left[\frac{|a_0|^2}{2} + \sum_{n=1}^{\infty} (|a_n|^2 + |b_n|^2) \right] \end{aligned}$$

and is derived by orthogonality and direct calculation.

3 PDEs

3.1 Separation of variables

The only methodology considered is separation of variables. So for PDE:

$$\hat{D}[u(x_1, \dots, x_n)] = 0$$

where \hat{D} is our differential operator, we look for solutions of form:

$$u(x_1, \dots, x_n) = X_1(x_1) \cdots X_n(x_n)$$

subject to **initial** and **boundary** conditions.

3.2 Heat equation

The heat equation is an equation of the following form:

$$\frac{\partial u}{\partial t} = \alpha^2 \frac{\partial^2 u}{\partial x^2}.$$

where α^2 is the thermal diffusivity constant.

3.2.1 Standard boundary conditions

We firstly define:

- **Initial condition:** $u(x, 0) = f(x)$ for $0 \leq x \leq L$
- **Boundary condition:** $u(0, t) = u(L, t) = 0$ for $\forall t > 0$

Let solutions be of form:

$$\begin{aligned} u(x, t) &= X(x) \cdot T(t) \\ \therefore X(x) \cdot \dot{T}(t) &= \alpha^2 X''(x) \cdot T(t) \end{aligned}$$

Only a constant function may satisfy the first equality:

$$\frac{1}{\alpha^2} \frac{\dot{T}}{T} = \frac{X''}{X} = -\lambda.$$

Writing this as two ODEs:

$$\begin{aligned} \dot{T} + \alpha^2 \lambda T &= 0 \\ X'' + \lambda X &= 0. \end{aligned}$$

The first one we can directly integrate, yielding:

$$T(t) = a_1 \exp(-\alpha^2 \lambda t).$$

The second ODE is a spring system, hence it has solution of form:

$$X(x) = b_1 \cos \lambda^{1/2} x + b_2 \sin \lambda^{1/2} x.$$

However this time before proceeding we need to consider boundary conditions:

$$X(0) = X(L) = 0.$$

We find $X(0) = b_1 = 0$ and $X(L) = b_2 \sin \lambda^{1/2} L = 0$.

The second equation implies that λ must of the following form:

$$\lambda_n = \left(\frac{n\pi}{L}\right)^2 \quad \text{for } \forall n \in \mathbb{N}$$

and so

$$X'' + \lambda X = 0 \implies X_n = b_2 \sin \lambda_n^{1/2} x.$$

Since λ is discretised:

$$\therefore T_n = a_1 \exp(-\alpha^2 \lambda_n t).$$

Our general solution must then be:

$$u(x, t) = \sum_{n=1}^{\infty} c_n \exp(-\alpha^2 \lambda_n t) \sin \lambda_n^{1/2} x$$

where $\lambda_n = \left(\frac{n\pi}{L}\right)^2$. Using initial condition $u(x, 0) = f(x)$:

$$f(x) = \sum_{n=1}^{\infty} c_n \sin \lambda_n^{1/2} x$$

and we recognise this as an odd Fourier series with period $2L$.

$$\begin{aligned} \therefore \int_{-L}^L \sin(\lambda_n^{1/2} x) f(x) dx &= \sum_{n=1}^{\infty} c_n \int_{-L}^L \left(\sin(\lambda_n^{1/2} x)\right)^2 dx \\ \therefore 2 \int_0^L \sin(\lambda_n^{1/2} x) f(x) dx &= c_n L \end{aligned}$$

The final step we split the integration range and use $x = -x^*$.

$$\therefore c_n = \frac{2}{L} \int_0^L \sin(\lambda_n^{1/2} x) f(x) dx$$

This is fine because we can extend $u(x, t)$ via reflection for negative x .

3.2.2 Fixed boundary temperatures

We reconsider the heat equation:

$$\frac{\partial u}{\partial t} = \alpha^2 \frac{\partial^2 u}{\partial x^2}$$

but now with the following **non-homogeneous boundary conditions**:

- $u(0, t) = T_1$
- $u(L, t) = T_2$
- $u(x, 0) = f(x)$

Physically our rod has fixed boundary temperatures, namely T_1 and T_2 .

We approach this problem with a change of variables:

$$v(x) = \lim_{t \rightarrow \infty} u(x, t).$$

Using our boundary conditions v must be linear:

$$\therefore v(x) = \frac{T_2 - T_1}{L}x + T_1$$

since $v'' = 0$, $v(0) = T_1$ and $v(L) = T_2$. We then deduce that:

$$u(x, t) = v(x) + \omega(x, t)$$

for $\omega(x, t)$ satisfies the same heat equation with initial conditions:

- $\omega(0, t) = \omega(L, t) = 0$
- $\omega(x, 0) = f(x) - v(x)$

Recognising this as our initial example:

$$\omega(x, t) = \sum_{n=1}^{\infty} c_n \exp(-\alpha^2 \lambda_n t) \sin \lambda_n^{1/2} x$$

where again $\lambda_n = \left(\frac{n\pi}{L}\right)^2$ and because $\omega(x, t)$ is a Fourier series with period $2L$:

$$c_n = \frac{2}{L} \int_0^L \sin(\lambda_n^{1/2} x) \{f(x) - v(x)\} dx.$$

3.2.3 Insulated rod ends

For the final example we consider:

$$\frac{\partial u}{\partial t} = \alpha^2 \frac{\partial^2 u}{\partial x^2}$$

and define the following conditions:

- $\frac{\partial}{\partial x} u(0, t) = \frac{\partial}{\partial x} u(L, t) = 0$
- $u(x, 0) = f(x)$

We begin again with a separation of variables:

3.3 Wave equation

3.4 Laplace's equation

Laplace's equation takes the form $\nabla^2 u = 0$. In two dimensions:

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

and we only consider boundary conditions. (Dirichlet conditions)

3.4.1 Rectangular boundary conditions

We open with the following example:

- **Boundary for y:** $u(x, 0) = u(x, b) = 0$
- **Boundary for x:** $u(0, y) = 0$ and $u(a, y) = f(y)$

where $x \in [0, a]$ and $y \in [0, b]$. Begin by separation of variables:

$$\begin{aligned} u(x, y) &= X(x) \cdot Y(y) \\ \therefore \frac{X''}{X} &= -\frac{Y''}{Y} = \lambda. \end{aligned}$$

Recognising the previous statement as two ODEs:

$$X'' - \lambda X = 0 \text{ for } X(0) = 0$$

$$Y'' + \lambda Y = 0 \text{ for } Y(0) = Y(b) = 0$$

The second ODE we have already solved in the heat equation. It has solution:

$$Y_n = a_1 \sin(\lambda_n^{1/2} y) \text{ for } \lambda_n = \frac{n^2 \pi^2}{b^2}.$$

The first ODE has solutions of form:

$$X_n = a_2 \cosh(\lambda_n^{1/2} x) + a_3 \sinh(\lambda_n^{1/2} x)$$

where these are the hyperbolic functions:

$$\sinh x = \frac{1}{2}(e^x - e^{-x})$$

$$\cosh x = \frac{1}{2}(e^x + e^{-x}).$$

Using our boundary condition $X(0) = 0$ gives:

$$X_n = a_3 \sinh(\lambda_n^{1/2} x).$$

Now putting all of this together we get:

$$u(x, y) = \sum_{n=1}^{\infty} c_n \sinh(\lambda_n^{1/2} x) \sin(\lambda_n^{1/2} y)$$

To find coefficients c_n we use $u(a, y) = f(y)$.

$$\therefore f(y) = \sum_{n=1}^{\infty} c_n \sinh(\lambda_n^{1/2} a) \sin(\lambda_n^{1/2} y)$$

Since we have a Fourier series with period $2b$:

$$\begin{aligned} \int_{-b}^b \sin(\lambda_n^{1/2} y) f(y) dy &= \sum_{n=1}^{\infty} c_n \sinh(\lambda_n^{1/2} a) \int_{-b}^b \sin(\lambda_n^{1/2} y) dy \\ &= c_n \sinh(\lambda_n^{1/2} a) \cdot b \end{aligned}$$

We can split the first integral to give us:

$$c_n = \frac{2}{b \sinh(\lambda_n^{1/2} a)} \int_0^b \sin(\lambda_n^{1/2} y) f(y) dy$$

where $\lambda_n = \left(\frac{n\pi}{b}\right)^2$ and our solution is complete.

3.4.2 Circular boundary conditions

Now we solve Laplace's equation

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

but with a circular boundary. In polar coordinates (r, θ) :

- $u(a, \theta) = f(\theta)$
- $u(r, \theta)$ is bounded

where a is the radius of our circle and $\theta \in [0, 2\pi]$. Since $u = u(x, y)$:

$$u'_\theta = u'_x x'_\theta + u'_y y'_\theta$$

$$u''_{\theta\theta} = (u''_{xx} x'_\theta + u''_{xy} y'_\theta) x'_\theta + u'_x x''_{\theta\theta} + (u''_{yy} y'_\theta + u''_{xy} x'_\theta) y'_\theta + u'_y y''_{\theta\theta}$$

$$u'_r = u'_x x'_r + u'_y y'_r$$

$$u''_{rr} = (u''_{xx} x'_r + u''_{xy} y'_r) x'_r + u'_x x''_{rr} + (u''_{xy} x'_r + u''_{yy} y'_r) y'_r + u'_y y''_{rr}$$

and here we have used the chain rule.

Applying these derivatives we obtain the following equation:

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} + \frac{1}{r} \frac{\partial u}{\partial r} = 0.$$

Using separation of variables:

$$u(r, \theta) = R(r)\Theta(\theta)$$

$$\therefore r^2 \frac{R''}{R} + r \frac{R'}{R} = -\frac{\ddot{\Theta}}{\Theta} = \lambda$$

where λ is our separation constant.

$$\therefore \ddot{\Theta} + \lambda\Theta = 0$$

$$\therefore r^2 R'' + rR' = \lambda R$$

4 Sturm-Liouville theory

4.1 Regular S-L problems

Sturm-Liouville theory is a general theory for 2nd order ODEs.

Consider the following eigenvalue ODE:

$$-\frac{d}{dx} \left(p(x) \frac{dy}{dx} \right) + q(x)y = \lambda r(x)y$$

where $r(x)$ is our weight function. We define the following boundary conditions:

1. $a_1 y(0) + a_2 y'(0) = 0$
2. $b_1 y(1) + b_2 y'(1) = 0$.

This is a **regular Sturm-Liouville** problem, where $p(x)$, $p'(x)$, $q(x)$, $r(x)$ are continuous functions and $p(x)$, $r(x)$ are strictly positive functions for $\forall x \in [0, 1]$.

Eigenvalues λ_n yield **eigenfunctions** $\phi_n(x)$ which are nontrivial solutions to our S-L problem. Important consequences include:

- Eigenvalues λ_n of a S-L problem are **real**.

Furthermore each eigenvalue corresponds to one eigenfunction.

- Eigenfunctions $\phi_n(x)$ are orthogonal:

$$\langle \phi_m, \phi_n \rangle = \int_0^1 r(x) \phi_m(x) \phi_n(x) dx = \delta_{mn}$$

in Hilbert space $L^2([0, 1], r(x)dx)$.

Note that our eigenfunctions are orthonormal and so we define:

$$\phi_n(x) = k_n y_n(x)$$

where k_n is our scale factor. Since $\langle \phi_n, \phi_n \rangle = 1$:

$$\therefore \int_0^1 r(x) k_n^2 y_n^2(x) dx = 1$$

and so we have that:

$$\begin{aligned} k_n &= \frac{1}{\sqrt{\langle y_n, y_n \rangle}} \\ &= \left(\int_0^1 r(x) y_n^2(x) dx \right)^{-1/2}. \end{aligned}$$

4.1.1 General 2nd order ODEs

Consider the following general 2nd order eigenvalue ODE:

$$-P(x)\frac{d^2y}{dx^2} - \omega(x)\frac{dy}{dx} + q(x)y = \lambda r(x)y.$$

Multiply this by the following integrating factor:

$$F(x) = \exp\left[\int_0^x \frac{\omega(s) - p'(s)}{p(s)} ds\right]$$

yields an ODE of S-L form:

$$-\frac{d}{dx}\left[F(x)P(x)\frac{dy}{dx}\right] + F(x)q(x)y = \lambda F(x)r(x)y.$$

4.1.2 Lagrange's identity

Our previous definition is motivated by the **Lagrange's identity**:

$$\begin{aligned}\langle \mathcal{L}[u], v \rangle - \langle u, \mathcal{L}[v] \rangle &= -\left[p(u'v^* - u(v^*)')\right]_0^1 \\ &= -\left[p(x)\left(\frac{du}{dx} \cdot v^* - u \cdot \frac{dv^*}{dx}\right)\right]_0^1\end{aligned}$$

where $u = u(x)$, $v = v(x)$ are complex functions and

$$\mathcal{L}[u] = -\frac{d}{dx}\left[p(x)\frac{du}{dx}\right] + q(x)u.$$

Here the inner product is defined as

$$\langle u, v \rangle = \int_0^1 uv^* dx$$

and we have integrated by parts using the following identities:

$$[pu'v^*]' = (pu')'v^* + pu'(v^*)'$$

$$[pu(v^*)']' = (p(v^*)')'u + pu'(v^*)'.$$

For a S-L problem its properties follow from:

$$\langle \mathcal{L}[u], v \rangle = \langle u, \mathcal{L}[v] \rangle$$

where functions u and v satisfy its boundary conditions.

4.1.3 Series expansion

Now the set of orthonormal eigenfunctions $\{\phi_n(x)\}$ from a S-L problem with boundary conditions may be used to expand function $f(x)$:

$$f_\phi(x) = \sum_{n=1}^{\infty} c_n \phi_n(x)$$

for $\forall x \in [0, 1]$. Integrating this on both sides:

$$\begin{aligned} \int_0^1 r(x) \phi_m(x) f(x) dx &= \int_0^1 r(x) \phi_m(x) \sum_{n=1}^{\infty} c_n \phi_n(x) dx \\ &= \sum_{n=1}^{\infty} c_n \int_0^1 r(x) \phi_m(x) \phi_n(x) dx \\ &= \sum_{n=1}^{\infty} c_n \delta_{mn} \\ &= c_m \end{aligned}$$

and so we get a general formula for our coefficients:

$$c_n = \int_0^1 r(x) \phi_n(x) f(x) dx.$$

If $f(x)$ and $f'(x)$ are piecewise continuous on $x \in [0, 1]$ then:

$$\forall x \in (0, 1); \sum_{n=1}^{\infty} c_n \phi_n(x) = \frac{f(x^-) + f(x^+)}{2}$$

where these are the left and right handed limits.

4.2 Non-homogeneous S-L problems

Consider:

$$\mathcal{L}[y] = \mu r(x)y + f(x)$$

where $f(x)$ is our non-homogeneous term, and that we define

$$\mathcal{L}[y] = -\frac{d}{dx} \left[P(x) \frac{dy}{dx} \right] + q(x)y.$$

Given that this ODE satisfies the S-L boundary conditions:

$$\mathcal{L}[y] = \lambda r(x)y$$

is solved by a sum of eigenfunctions

$$y(x) = \sum_{n=1}^{\infty} b_n \phi_n(x).$$

Substituting this into our original ODE gives:

$$\begin{aligned} \sum_{n=1}^{\infty} b_n \mathcal{L}[\phi_n(x)] &= r(x) \sum_{n=1}^{\infty} b_n \lambda_n \phi_n(x) \\ &= \mu r(x) \sum_{n=1}^{\infty} b_n \phi_n(x) + f(x). \end{aligned}$$

Rearranging then gives:

$$\frac{f(x)}{r(x)} = \sum_{n=1}^{\infty} b_n (\lambda_n - \mu) \phi_n(x).$$

Now we can also expand $\frac{f(x)}{r(x)}$ in terms of our eigenfunctions:

$$\frac{f(x)}{r(x)} = \sum_{n=1}^{\infty} c_n \phi_n(x)$$

with

$$\begin{aligned} c_n &= \int_0^1 r(x) \phi_n(x) \frac{f(x)}{r(x)} dx \\ &= \int_0^1 \phi_n(x) f(x) dx \end{aligned}$$

and so equating yields the following relation

$$b_n = \frac{c_n}{\lambda_n - \mu}.$$

4.3 Singular S-L problems

quite a lot of theory not covered in this course

general definition of singular s-l problems

bessel's equation (order 0 example)

conditions for singular problems

4.4 S-L Series solutions

5 Laplace transforms