

## Vector products

$$\mathbf{a} \cdot \mathbf{b} = ab \cos \theta$$

$$\mathbf{a} \times \mathbf{b} = ab \sin \theta \hat{\mathbf{n}}$$

$$\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a} \text{ and } \mathbf{a} \times \mathbf{a} = \mathbf{0}$$

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b})$$

## Suffix notation

1. A suffix that appears twice implies a summation.
2. Any suffix cannot appear more than twice in any term.

We define the **Kronecker delta** as:

$$\delta_{ij} = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}$$

and the **Levi-Civita** as:

$$\epsilon_{ijk} = \begin{cases} +1 & 123, 312, 231 \\ -1 & 132, 213, 321 \\ 0 & \text{repeat indices.} \end{cases}$$

Consequently:

$$\begin{aligned} \epsilon_{ijk} &= \epsilon_{kij} = \epsilon_{jki} \\ &= -\epsilon_{jik} = -\epsilon_{ikj} = -\epsilon_{kji} \end{aligned}$$

and we have the following identities:

$$\mathbf{a} = \sum_{i=1}^3 a_i \mathbf{e}_i = a_i \mathbf{e}_i$$

$$A\mathbf{x} = a_{ij}x_j \mathbf{e}_i \text{ for } m \times n \text{ matrix } A$$

$$\delta_{ii} = 3$$

$$[\dots]_j \delta_{jk} = [\dots]_k$$

$$\mathbf{e}_i \cdot \mathbf{e}_j = \delta_{ij}$$

$$\mathbf{e}_i \times \mathbf{e}_j = \epsilon_{ijk} \mathbf{e}_k$$

$$\mathbf{a} \times \mathbf{b} = \epsilon_{ijk} a_j b_k \mathbf{e}_i$$

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \epsilon_{ijk} a_i b_j c_k$$

$$\epsilon_{ijk} \epsilon_{klm} = \delta_{il} \delta_{jm} - \delta_{im} \delta_{jl}$$

$$\epsilon_{ijk} \epsilon_{ijl} = 2\delta_{kl} \text{ and } \epsilon_{ijk} \epsilon_{ijk} = 6.$$

## Transformations

Let matrix  $L$  relate basis  $\{\mathbf{e}_i\}$  to basis  $\{\mathbf{e}'_i\}$  with rule:

$$\mathbf{e}'_i = \ell_{ij} \mathbf{e}_j \text{ where } (L)_{ij} = \ell_{ij}.$$

Then  $L^T L = L L^T = I$ , and:

$$\ell_{ik} \ell_{jk} = \ell_{ki} \ell_{kj} = \delta_{ij}$$

$$p'_i = \ell_{ij} p_j \text{ for } \mathbf{p} = p_i \mathbf{e}_i = p'_i \mathbf{e}'_i.$$

## Tensors

A rank 3 tensor is defined as:

$$T'_{ijk} = \ell_{ip} \ell_{jq} \ell_{kr} T_{pqr}$$

which relates frame  $S$  in  $\{\mathbf{e}_i\}$  to frame  $S'$  in  $\{\mathbf{e}'_i\}$  with rule  $\mathbf{e}'_i = \ell_{ij} \mathbf{e}_j$ , etc.

Properties of tensors:

1. The addition of two rank  $n$  tensors is also a rank  $n$  tensor.
2. The multiplication of a rank  $m$  tensor with a rank  $n$  tensor yields a rank  $m + n$  tensor.
3. If  $T_{ijk\dots s}$  is a rank  $m$  tensor then  $T_{\mathbf{ii}k\dots s}$  is a rank  $m - 2$  tensor.
4. If  $T_{ij}$  is a tensor then  $T_{ji}$  is also a tensor. Explicitly:

$$\begin{aligned} T'_{ij} &= \ell_{ip} \ell_{jq} T_{pq} \implies T' = L T L^T \\ T'_{\mathbf{ji}} &= \ell_{jp} \ell_{iq} T_{pq}. \end{aligned}$$

## Symmetric tensors

$T_{ij}$  is a symmetric tensor when  $T_{ij} = T_{ji}$  in frame  $S$ . Then  $T'_{ij} = T'_{ji}$  in frame  $S'$ .

Similarly  $T_{ij}$  is an anti-symmetric tensor if  $T_{ij} = -T_{ji}$  and  $T'_{ij} = -T'_{ji}$ .

Finally **any tensor** can be written as a sum of symmetric and anti-symmetric parts:

$$T_{ij} = \frac{1}{2}(T_{ij} + T_{ji}) + \frac{1}{2}(T_{ij} - T_{ji}).$$

## Quotient theorem

Consider 9 entities  $T_{ij}$  in frame  $S$  and  $T'_{ij}$  in frame  $S'$ . Let  $b_i = T_{ij} a_j$  where  $a_j$  is a vector. If  $b_i$  always transforms as a vector then  $T_{ij}$  is a rank 2 tensor.

Generalising, let  $R_{ijk\dots r}$  be a rank  $m$  tensor and  $T_{ijk\dots s}$  a set of  $3^n$  numbers where  $n > m$ . If  $T_{ijk\dots s} R_{ijk\dots r}$  is a rank  $n - m$  tensor then  $T_{ijk\dots s}$  is a rank  $n$  tensor.

## Matrices

We define a  $m \times n$  matrix  $A$  as  $(A)_{ij} = a_{ij}$  where  $i = 1, \dots, m$  and  $j = 1, \dots, n$ .

- $\text{Tr } A = a_{ii}$
- $(A^T)_{ij} = a_{ji}$
- $(AB)^T = B^T A^T$
- $(I)_{ij} = \delta_{ij}$

The determinant of a  $3 \times 3$  matrix  $A$  is:

$$\begin{aligned} \det A &= \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \\ &= \epsilon_{lmn} a_{1l} a_{2m} a_{3n} \\ &= \epsilon_{lmn} a_{l1} a_{m2} a_{n3}. \end{aligned}$$

Furthermore:

$$\begin{aligned} \epsilon_{ijk} \det A &= \epsilon_{lmn} a_{il} a_{jm} a_{kn} \\ \epsilon_{lmn} \det A &= \epsilon_{ijk} a_{il} a_{jm} a_{kn} \\ \det A &= \frac{1}{3!} \epsilon_{ijk} \epsilon_{lmn} a_{il} a_{jm} a_{kn}. \end{aligned}$$

Properties of determinants:

1. Adding rows to each other does not change the determinant.
2. Interchanging two rows changes determinant signs.
3.  $\det A = \det A^T$
4.  $\det(AB) = \det A \cdot \det B$

These also apply to columns. Finally:

$$\epsilon_{ijk} \epsilon_{lmn} \det A = \begin{vmatrix} a_{il} & a_{im} & a_{in} \\ a_{jl} & a_{jm} & a_{jn} \\ a_{kl} & a_{km} & a_{kn} \end{vmatrix}$$

and setting  $A = I$  yields:

$$\epsilon_{ijk} \epsilon_{lmn} = \begin{vmatrix} \delta_{il} & \delta_{im} & \delta_{in} \\ \delta_{jl} & \delta_{jm} & \delta_{jn} \\ \delta_{kl} & \delta_{km} & \delta_{kn} \end{vmatrix}.$$

## Linear equations

Let  $\mathbf{y} = A\mathbf{x}$ . Then  $x_i = A_{ij}^{-1} y_j$  with:

$$\begin{aligned} A_{ij}^{-1} &= \frac{1}{2 \det A} \epsilon_{imn} \epsilon_{jpk} a_{pm} a_{qn} \\ &= \frac{1}{\det A} C_{ij}^T \end{aligned}$$

where  $C$  is the cofactor matrix of  $A$ .

## Pseudotensors

A rank 2 pseudotensor is defined as:

$$T'_{ij} = (\det L) \ell_{ip} \ell_{jq} T_{pq}$$

where  $(L)_{ij} = \ell_{ij}$  and  $\det L = \pm 1$ .

Pseudovectors are rank 1 pseudotensors.

## Invariant tensors

Tensor  $T$  is invariant or isotropic if:

$$T_{ijk\dots} = \ell_{i\alpha} \ell_{j\beta} \ell_{k\gamma} \dots T_{\alpha\beta\gamma\dots}$$

for every orthogonal matrix  $L$ .

- If  $a_{ij}$  is a rank 2 invariant tensor then  $a_{ij} = \lambda \delta_{ij}$ .
- The most general rank 3 invariant pseudotensor is  $a_{ijk} = \lambda \epsilon_{ijk}$ . There are no rank 3 invariant true tensors.
- Invariant true tensors can only be even ranked.
- Invariant pseudotensors can only be odd ranked.

## Rotation tensors

The clockwise rotation of position vector  $\mathbf{x}$  to  $\mathbf{y}$  about unit vector  $\hat{\mathbf{n}}$  is given by:

$$y_i = R_{ij}(\theta, \hat{\mathbf{n}})x_j$$

$$R_{ij}(\theta, \hat{\mathbf{n}}) = \delta_{ij} \cos \theta + (1 - \cos \theta)n_i n_j - \epsilon_{ijk} n_k \sin \theta$$

and is the rotation tensor.

## Reflections and inversions

The reflection of vector  $\mathbf{x}$  to  $\mathbf{y}$  in plane with unit vector  $\hat{\mathbf{n}}$  is:

$$y_i = \sigma_{ij}x_j$$

$$\sigma_{ij} = \delta_{ij} - 2n_i n_j.$$

The inversion of vector  $\mathbf{x}$  to  $\mathbf{y}$  is given by  $\mathbf{y} = -\mathbf{x}$  and is defined as:

$$y_i = P_{ij}x_j$$

$$P_{ij} = \delta_{ij}.$$

## Projections

We define  $P$  to be a parallel projection operator to vector  $\mathbf{u}$  if:

$$P\mathbf{u} = \mathbf{u} \text{ and } P\mathbf{v} = \mathbf{0}$$

where  $\mathbf{u} \cdot \mathbf{v} = \mathbf{0}$ . Then:

$$P_{ij} = \frac{u_i u_j}{u^2}.$$

Similarly we define  $Q$  to be an orthogonal projection to vector  $\mathbf{u}$  if:

$$Q\mathbf{u} = \mathbf{0} \text{ and } Q\mathbf{v} = \mathbf{v}.$$

Here  $Q = I - P$ .

## Inertia tensors

Let  $\mathbf{L}$  denote the angular momentum of a rigid body about the origin of mass  $m$ , volume  $V$  and density  $\rho$  at position  $\mathbf{r}$  with velocity  $\mathbf{v}$ . Then:

$$\mathbf{L}_i = I_{ij}\omega_j$$

$$I_{ij} = I_{ij}(O) = \int_V \rho(r^2 \delta_{ij} - x_i x_j) dV$$

where  $I_{ij}(O)$  is the inertia tensor about the origin. The kinetic energy of such a body is:

$$T = \frac{1}{2} I_{ij} \omega_i \omega_j = \frac{1}{2} \mathbf{L} \cdot \boldsymbol{\omega}.$$

## Parallel axis theorem

Consider the same rigid body now with centre of mass  $G$  and let  $\overrightarrow{OG} = \mathbf{R}$ . Then:

$$I_{ij}(O) = I_{ij}(G) + M(R^2 \delta_{ij} - X_i X_j)$$

$$M = \int_V \rho'(\mathbf{r}') dV'.$$

## Diagonalisation

Let  $\mathbf{L} = I_{ij}\omega_j$  where  $I_{ij}$  is a rank 2 tensor and let  $\mathbf{L} = \lambda\boldsymbol{\omega}$ . Then:

$$(I_{ij} - \lambda \delta_{ij})\omega_j = 0 \implies \det(I_{ij} - \lambda \delta_{ij}) = 0$$

where expanding this gives:

$$P - Q\lambda + R\lambda^2 - \lambda^3 = 0$$

for  $P = \det I$ ,  $Q = \frac{1}{2}[(\text{tr } I)^2 - \text{tr}(I^2)]$  and  $R = \text{tr } I$  given tensor  $I$ .

## Real symmetric tensors

Let rank 2 real symmetric tensor  $T$  be diagonalisable with real eigenvalues  $\lambda^{(i)}$  and orthonormal eigenvectors  $\boldsymbol{\ell}^{(i)}$  where  $i = 1, 2, 3$ . Let transformation matrix be:

$$L_{ij} = \ell_j^{(i)} = \begin{pmatrix} \ell_1^{(1)} & \ell_2^{(1)} & \ell_3^{(1)} \\ \ell_1^{(2)} & \ell_2^{(2)} & \ell_3^{(2)} \\ \ell_1^{(3)} & \ell_2^{(3)} & \ell_3^{(3)} \end{pmatrix}_{ij}$$

and always defined such that  $\det L = +1$  which transforms frame  $S \rightarrow S'$ .

Then since  $T_{pq}\ell_q^{(i)} = \lambda^{(i)}\ell_p^{(i)}$ :

$$T'_{ij} = \ell_{ip}\ell_{jq}T_{pq} = \lambda^{(i)}\delta_{ij} = \begin{pmatrix} \lambda^{(1)} & 0 & 0 \\ 0 & \lambda^{(2)} & 0 \\ 0 & 0 & \lambda^{(3)} \end{pmatrix}_{ij}.$$

## Taylor expansions

In the one-dimensional case we have:

$$f(x) = \sum_{n=0}^{\infty} \frac{1}{n!} f^{(n)}(a)(x-a)^n$$

and is  $f$  expanded about  $x = a$ .

Trigonometric expansions are in radians!

$$\begin{aligned} \therefore f(x+a) &= \sum_{n=0}^{\infty} \frac{1}{n!} f^{(n)}(x)a^n \\ &= \exp\left(a \frac{d}{dx}\right) f(x) \end{aligned}$$

Then for three dimensions:

$$\begin{aligned} \phi(\mathbf{r} + \mathbf{a}) &= \sum_{n=0}^{\infty} \frac{1}{n!} (\mathbf{a} \cdot \boldsymbol{\nabla}_r)^n \phi(\mathbf{r}) \\ &= \exp(\mathbf{a} \cdot \boldsymbol{\nabla}_r) \phi(\mathbf{r}). \end{aligned}$$

## Curvilinear coordinates

Let  $x_i$  denote Cartesian coordinates and  $u_i$  denote curvilinear coordinates. Then:

$$(x_1, x_2, x_3) \rightarrow (u_1, u_2, u_3)$$

where each  $u_i = u_i(x_1, x_2, x_3)$  and:

$$\begin{aligned} \mathbf{r} &= x_1 \mathbf{e}_1 + x_2 \mathbf{e}_2 + x_3 \mathbf{e}_3 \\ &= u_1 \mathbf{e}_{u_1} + u_2 \mathbf{e}_{u_2} + u_3 \mathbf{e}_{u_3}. \end{aligned}$$

## Scale factors

Let  $u_1 \rightarrow u_1 + du_1$  in  $\mathbf{r} = \mathbf{r}(u_1, u_2, u_3)$ . Then  $d\mathbf{r}$  in  $\mathbf{r} \rightarrow \mathbf{r} + d\mathbf{r}$  is defined as:

$$d\mathbf{r} = \frac{\partial \mathbf{r}}{\partial u_1} du_1 := h_1 \mathbf{e}_1 du_1.$$

$h_1$  is the scale factor of unit vector  $\mathbf{e}_1$ :

$$h_1 = \left| \frac{\partial \mathbf{r}}{\partial u_1} \right| \text{ and } \mathbf{e}_1 = \frac{1}{h_1} \frac{\partial \mathbf{r}}{\partial u_1}.$$

If  $\mathbf{e}_i \cdot \mathbf{e}_j = \delta_{ij}$  then  $u_i$  is an **orthogonal** curvilinear coordinate system.

## Vector and arc length

The vector length  $d\mathbf{r}$  of  $\mathbf{r}$  is defined as:

$$d\mathbf{r} = \sum_{i=1}^3 h_i du_i \mathbf{e}_i$$

where  $u_i \rightarrow u_i + du_i$  for  $\forall i = 1, 2, 3$ .

Then the arc length  $ds$  is defined as:

$$\begin{aligned} (ds)^2 &= d\mathbf{r} \cdot d\mathbf{r} \\ &= g_{ij} du_i du_j \end{aligned}$$

where  $g_{ij}$  is the metric tensor:

$$\begin{aligned} g_{ij} &= g_{ji} = \frac{\partial x_k}{\partial u_i} \frac{\partial x_k}{\partial u_j} \\ &= h_i h_j (\mathbf{e}_i \cdot \mathbf{e}_j). \end{aligned}$$

## Area and volume

Let  $d\mathbf{r}_i = h_i \mathbf{e}_i du_i$  denote vector length when  $u_i \rightarrow u_i + du_i$ . (**No** sum!)

The infinitesimal vector area formed by  $d\mathbf{r}_1$  and  $d\mathbf{r}_2$  is:

$$d\mathbf{S} = (h_1 d\mathbf{u}_1 \mathbf{e}_1) \times (h_2 d\mathbf{u}_2 \mathbf{e}_2).$$

Similarly the infinitesimal volume formed by edges  $d\mathbf{r}_1$ ,  $d\mathbf{r}_2$  and  $d\mathbf{r}_3$  is:

$$\begin{aligned} dV &= |(d\mathbf{r}_1 \times d\mathbf{r}_2) \cdot d\mathbf{r}_3| \\ &= \sqrt{g} du_1 du_2 du_3 \end{aligned}$$

where  $g = \det(g_{ij})$ .

## Cylindrical coordinates

$(u_1, u_2, u_3) = (\rho, \phi, z)$  where  $\rho$  represents the radial distance from the origin and  $\phi$  is the anticlockwise rotation angle on the  $x$ - $y$  plane. In Cartesian unit vectors:

$$\begin{aligned} \mathbf{r} &= \rho \cos \phi \mathbf{e}_1 + \rho \sin \phi \mathbf{e}_2 + z \mathbf{e}_3 \\ h_\rho &= 1, \quad \mathbf{e}_\rho = \cos \phi \mathbf{e}_1 + \sin \phi \mathbf{e}_2 \\ h_\phi &= \rho, \quad \mathbf{e}_\phi = -\sin \phi \mathbf{e}_1 + \cos \phi \mathbf{e}_2 \\ h_z &= 1, \quad \mathbf{e}_z = \mathbf{e}_3 \end{aligned}$$

and forms an orthogonal set.

## Spherical coordinates

$(u_1, u_2, u_3) = (r, \theta, \phi)$  where  $\theta$  represents the clockwise rotation angle in  $y$ - $z$  plane and  $\phi$  the anticlockwise rotation angle in  $x$ - $y$  plane. In Cartesian unit vectors:

$$\mathbf{r} = r \sin \theta \cos \phi \mathbf{e}_1 + r \sin \theta \sin \phi \mathbf{e}_2 + r \cos \theta \mathbf{e}_3$$

$$h_r = 1, \quad h_\theta = r, \quad h_\phi = r \sin \theta$$

$$\mathbf{e}_r = \sin \theta \cos \phi \mathbf{e}_1 + \sin \theta \sin \phi \mathbf{e}_2 + \cos \theta \mathbf{e}_3$$

$$\mathbf{e}_\theta = \cos \theta \cos \phi \mathbf{e}_1 + \cos \theta \sin \phi \mathbf{e}_2 - \sin \theta \mathbf{e}_3$$

$$\mathbf{e}_\phi = -\sin \phi \mathbf{e}_1 + \cos \phi \mathbf{e}_2$$

and also forms an orthogonal set.



## Gradient

The gradient of a scalar field  $f(\mathbf{r})$  is:

$$df(\mathbf{r}) := \nabla f(\mathbf{r}) \cdot d\mathbf{r}$$

when  $\mathbf{r} \rightarrow \mathbf{r} + d\mathbf{r} \implies f \rightarrow f + df$ .  
Taking the total differential of  $f$  yields:

$$\nabla f = \sum_{i=1}^3 \frac{1}{h_i} \frac{\partial f}{\partial u_i} \mathbf{e}_i$$

where  $\{\mathbf{e}_i\}$  is orthogonal.

## Divergence

The divergence of a vector field  $\mathbf{F}$  is:

$$\nabla \cdot \mathbf{F} := \lim_{\delta V \rightarrow 0} \frac{1}{\delta V} \int_{\delta S} \mathbf{F} \cdot d\mathbf{S}$$

for surface  $\delta S$  bounds infinitesimal  $\delta V$ .  
In orthogonal curvilinear coordinates:

$$\nabla \cdot \mathbf{F} = \frac{1}{h_1 h_2 h_3} \left\{ \frac{\partial}{\partial u_1} (F_1 h_2 h_3) + \frac{\partial}{\partial u_2} (h_1 F_2 h_3) + \frac{\partial}{\partial u_3} (h_1 h_2 F_3) \right\}.$$

## Curl

The curl of a vector field  $\mathbf{F}$  in the direction of unit vector  $\hat{\mathbf{n}}$  is:

$$\hat{\mathbf{n}} \cdot (\nabla \times \mathbf{F}) := \lim_{\delta S \rightarrow 0} \frac{1}{\delta S} \oint_{\delta C} \mathbf{F} \cdot d\mathbf{r}$$

where curve  $\delta C$  encloses plane  $\delta S$ .

In orthogonal curvilinear coordinates:

$$\nabla \times \mathbf{F} = \frac{1}{h_1 h_2 h_3} \begin{vmatrix} h_1 \mathbf{e}_1 & h_2 \mathbf{e}_2 & h_3 \mathbf{e}_3 \\ \frac{\partial}{\partial u_1} & \frac{\partial}{\partial u_2} & \frac{\partial}{\partial u_3} \\ h_1 a_1 & h_2 a_2 & h_3 a_3 \end{vmatrix}.$$

## Laplacian

The Laplacian of a scalar field  $f$  is:

$$\nabla^2 f = \nabla \cdot (\nabla f)$$

and in orthogonal curvilinear coordinates:

$$\nabla^2 f = \frac{1}{h_1 h_2 h_3} \left\{ \frac{\partial}{\partial u_1} \left( \frac{h_2 h_3}{h_1} \frac{\partial f}{\partial u_1} \right) + \frac{\partial}{\partial u_2} \left( \frac{h_3 h_1}{h_2} \frac{\partial f}{\partial u_2} \right) + \frac{\partial}{\partial u_3} \left( \frac{h_1 h_2}{h_3} \frac{\partial f}{\partial u_3} \right) \right\}.$$

The Laplacian of a vector field  $\mathbf{F}$  is:

$$\nabla^2 \mathbf{F} = \nabla(\nabla \cdot \mathbf{F}) - \nabla \times (\nabla \times \mathbf{F}).$$

## Vector calculus identities

Particularly for Cartesian coordinates we can apply the suffix notation:

$$\nabla f = \frac{\partial f}{\partial x_i} \mathbf{e}_i$$

$$(\mathbf{u} \cdot \nabla) \mathbf{F} = u_j \frac{\partial}{\partial x_j} F_i$$

$$\nabla \cdot \mathbf{F} = \frac{\partial F_i}{\partial x_i}$$

$$\nabla \times \mathbf{F} = \epsilon_{ijk} \frac{\partial F_k}{\partial x_j} \mathbf{e}_i$$

$$\frac{\partial x_i}{\partial x_j} = \delta_{ij} \text{ and } \frac{\partial x_i}{\partial x_i} = \delta_{ii} = 3.$$

If  $\psi$  is a scalar field and  $\mathbf{v}$  a vector field:

$$\nabla \times (\nabla \psi) = \mathbf{0}$$

$$\nabla \cdot (\nabla \times \mathbf{v}) = 0$$

$$\nabla \cdot (\psi \mathbf{v}) = \nabla \psi \cdot \mathbf{v} + \psi \nabla \cdot \mathbf{v}$$

$$\nabla \times (\psi \mathbf{v}) = \nabla \psi \times \mathbf{v} + \psi \nabla \times \mathbf{v}.$$

Let  $\mathbf{r} = x_i \mathbf{e}_i$  and  $r = (x_i^2)^{1/2}$ . Then:

- $\nabla r = \frac{\mathbf{r}}{r}$
- $\nabla r^n = n r^{n-2} \mathbf{r}$
- $\nabla \cdot \mathbf{r} = 3$  and  $\nabla \times \mathbf{r} = \mathbf{0}$
- $\nabla \times (\mathbf{c} \times \mathbf{r}) = 2\mathbf{c}$
- $\nabla \cdot (\mathbf{c} \times \mathbf{r}) = 0$  for constant  $\mathbf{c}$ .

## Divergence theorem

Let surface  $S$  enclose volume  $V$ . Then:

$$\iiint_V \nabla \cdot \mathbf{F} dV = \oint_S \mathbf{F} \cdot d\mathbf{S}$$

where  $\mathbf{F}$  is a vector field.

## Stokes' theorem

Let closed curve  $C$  bound open surface  $S$  and let  $\mathbf{F}$  be a vector field. Then:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S}$$

for  $C$  is traversed in anticlockwise sense.

## Dirac delta function

The Dirac delta in 1D is defined as:

$$\delta(x - a) := \begin{cases} \infty & x = a \\ 0 & \text{otherwise.} \end{cases}$$

In three dimensions this becomes:

$$\begin{aligned} \delta^{(3)}(\mathbf{r} - \mathbf{r}_0) &:= \delta(\mathbf{r} - \mathbf{r}_0) \\ &= \delta(x - x_0) \delta(y - y_0) \delta(z - z_0) \end{aligned}$$

where  $(x, y, z)$  are Cartesian coordinates.

In orthogonal curvilinear coordinates:

$$\begin{aligned} \delta(\mathbf{r} - \mathbf{a}) &= \frac{1}{h_1 h_2 h_3} \delta(u_1 - a_1) \\ &\quad \cdot \delta(u_2 - a_2) \delta(u_3 - a_3). \end{aligned}$$