

Q 1. Let γ be a real constant with $\gamma^2 \neq 1$ for the parametric functional

$$S[x, y] = \int_0^1 dt \left[\sqrt{\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2} - \lambda(\dot{x}y - x\dot{y}) \right], \quad \lambda > 0,$$

with the boundary conditions $x(0) = y(0) = 0, x(1) = R > 0$ and $y(1) = 0$.

(a) In order to show that the stationary paths of this parametric functional are given by the solutions of the equations

$$\frac{dx}{ds} + \gamma \frac{dy}{ds} = 2(c - \lambda y), \text{ and}$$

$$\gamma \frac{dx}{ds} + \frac{dy}{ds} = 2(d + \lambda x),$$

where c and d are constants and

$$s(t) = \int_0^t dt \sqrt{\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2} \quad (1.1)$$

consider the following.

From the parametric functional,

$$\Phi = (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{\frac{1}{2}} - \lambda(\dot{x}y - x\dot{y}), \quad \lambda > 0,$$

$$\frac{\partial \Phi}{\partial \dot{x}} = \frac{1}{2} (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (2\dot{x} + 2\gamma\dot{y}) + \lambda y.$$

$$\frac{\partial \Phi}{\partial x} = -\lambda \dot{y}.$$

The Euler-Lagrange equation is given by

$$\frac{d}{dt} \left(\frac{\partial \Phi}{\partial \dot{x}} \right) - \frac{\partial \Phi}{\partial x} = 0.$$

Substituting into the Euler-Lagrange equation for the derivatives obtained above gives,

$$\frac{d}{dt} \left(\frac{1}{2} (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (2\dot{x} + 2\gamma\dot{y}) + \lambda y \right) + \lambda \dot{y} = 0,$$

$$\frac{d}{dt} \left((\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (\dot{x} + \gamma\dot{y}) + \lambda y \right) + \lambda \dot{y} = 0,$$

and integrating the above express with respect to t gives,

$$\left[(\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (\dot{x} + \gamma\dot{y}) + \lambda y \right] + \lambda y = 2c,$$

where c is an arbitrary constant.

$$(\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (\dot{x} + \gamma\dot{y}) + 2\lambda y = 2c,$$

So,

$$(\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (\dot{x} + \gamma\dot{y}) = 2(c - \lambda y). \quad (1.2)$$

Now, from (1.1) the definition of s ,

$$\frac{ds}{dt} = (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{\frac{1}{2}},$$

and

$$\frac{dt}{ds} = \left(1 / \frac{ds}{dt}\right) = (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}}. \quad (1.3)$$

Substituting (1.3) into (1.2) gives,

$$\frac{dt}{ds} (\dot{x} + \gamma\dot{y}) = 2(c - \lambda y),$$

and noting that

$$\frac{dx}{dt} = \dot{x} \quad \text{and} \quad \frac{dy}{dt} = \dot{y},$$

then

$$\begin{aligned} \frac{dt}{ds} \left(\frac{dx}{dt} + \gamma \frac{dy}{dt} \right) &= 2(c - \lambda y) \\ \frac{dt}{ds} \cdot \frac{dx}{dt} + \gamma \frac{dt}{ds} \cdot \frac{dy}{dt} &= 2(c - \lambda y), \end{aligned}$$

thus, by the chain rule,

$$\boxed{\frac{dx}{ds} + \gamma \frac{dy}{ds} = 2(c - \lambda y)}. \quad (1.4)$$

Similarly for \dot{y}

$$\Phi = (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{\frac{1}{2}} - \lambda(x\dot{y} - \dot{x}y), \quad \lambda > 0,$$

$$\frac{\partial \Phi}{\partial \dot{y}} = \frac{1}{2} (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (2\gamma\dot{x} + 2\dot{y}) - \lambda x.$$

$$\frac{\partial \Phi}{\partial y} = \lambda \dot{x}.$$

The Euler-Lagrange equation is given by

$$\frac{d}{dt} \left(\frac{\partial \Phi}{\partial \dot{y}} \right) - \frac{\partial \Phi}{\partial y} = 0.$$

Substituting into the Euler-Lagrange equation for the derivatives obtained above gives,

$$\frac{d}{dt} \left[\frac{1}{2} (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (2\gamma\dot{x} + 2\dot{y}) - \lambda x \right] - \lambda \dot{x} = 0,$$

$$\frac{d}{dt} \left[(\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (\gamma\dot{x} + \dot{y}) - \lambda x \right] - \lambda\dot{x} = 0,$$

and integrating the above express with respect to t gives,

$$(\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (\gamma\dot{x} + \dot{y}) - \lambda x - \lambda x = 2d,$$

where d is an arbitrary constant. So,

$$(\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (\gamma\dot{x} + \dot{y}) - 2\lambda x = 2d,$$

and thus,

$$(\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}} (\gamma\dot{x} + \dot{y}) = 2(d + \lambda x).$$

Recall, (1.3) from above,

$$\frac{dt}{ds} = (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{-\frac{1}{2}},$$

and hence,

$$\begin{aligned} \frac{dt}{ds} (\gamma\dot{x} + \dot{y}) &= 2(d + \lambda x), \\ \frac{dt}{ds} \left(\gamma \frac{dx}{dt} + \frac{dy}{dt} \right) &= 2(d + \lambda x), \\ \gamma \frac{dt}{ds} \cdot \frac{dx}{dt} + \frac{dt}{ds} \cdot \frac{dy}{dt} &= 2(d + \lambda x), \end{aligned}$$

Thus, by the chain rule,

$$\boxed{\gamma \frac{dx}{ds} + \frac{dy}{ds} = 2(d + \lambda x).} \quad (1.5)$$

Hence, the stationary paths of the given parametric functional are given by the solutions to equations (1.4) and (1.5).

(b) It can be shown that

$$\left(\frac{dx}{ds} \right)^2 + 2\gamma \frac{dx}{ds} \cdot \frac{dy}{ds} + \left(\frac{dy}{ds} \right)^2 = 1,$$

as follows.

From part (a)

$$\frac{ds}{dt} = (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2)^{\frac{1}{2}},$$

and squaring both sides of the above expression gives,

$$\frac{ds}{dt} \cdot \frac{ds}{dt} = (\dot{x}^2 + 2\gamma\dot{x}\dot{y} + \dot{y}^2), \quad (1.6)$$

Again, recall that

$$\frac{dx}{dt} = \dot{x} \quad \text{and} \quad \frac{dy}{dt} = \dot{y},$$

so that (1.6) becomes

$$\frac{ds}{dt} \cdot \frac{ds}{dt} = \left(\frac{dx}{dt} \cdot \frac{dx}{dt} + 2\gamma \frac{dx}{dt} \cdot \frac{dy}{dt} + \frac{dy}{dt} \cdot \frac{dy}{dt} \right), \quad (1.7)$$

multiplying both sides of equation (1.7) by

$$\frac{dt}{ds} \cdot \frac{dt}{ds}$$

gives,

$$\frac{dt}{ds} \cdot \frac{dt}{ds} \cdot \frac{ds}{dt} \cdot \frac{ds}{dt} = \frac{dt}{ds} \cdot \frac{dt}{ds} \cdot \frac{dx}{dt} \cdot \frac{dx}{dt} + 2\gamma \frac{dt}{ds} \cdot \frac{dt}{ds} \cdot \frac{dx}{dt} \cdot \frac{dy}{dt} + \frac{dt}{ds} \cdot \frac{dt}{ds} \cdot \frac{dy}{dt} \cdot \frac{dy}{dt}, \quad (1.8)$$

then by the chain rule (1.8) becomes

$$1 = \left(\frac{dx}{ds} \right)^2 + 2\gamma \frac{dx}{ds} \cdot \frac{dy}{ds} + \left(\frac{dy}{ds} \right)^2, \quad (1.9)$$

as required.

(c) From equation (1.4)

$$\frac{dx}{ds} = 2(c - \lambda y) - \gamma \frac{dy}{ds}$$

and from equation (1.5)

$$\frac{dy}{ds} = 2(d + \lambda x) - \gamma \frac{dx}{ds}$$

Substituting into equation (1.4) for dy/ds gives,

$$\begin{aligned} \frac{dx}{ds} + \gamma \left(2(d + \lambda x) - \gamma \frac{dx}{ds} \right) &= 2(c - \lambda y), \\ \frac{dx}{ds} + 2\gamma(d + \lambda x) - \gamma^2 \frac{dx}{ds} &= 2(c - \lambda y), \\ \frac{dx}{ds} (1 - \gamma^2) + 2\gamma(d + \lambda x) &= 2(c - \lambda y), \\ \frac{dx}{ds} (1 - \gamma^2) &= 2(c - \lambda y) - 2\gamma(d + \lambda x), \\ \frac{dx}{ds} (1 - \gamma^2) &= 2c - 2\lambda y - 2\gamma d - 2\gamma \lambda x, \\ \frac{dx}{ds} (1 - \gamma^2) &= -2\lambda(y + \gamma x) + 2(c - \gamma d), \\ x' (1 - \gamma^2) &= -X + D = D - X. \end{aligned}$$

Therefore,

$$\boxed{(1 - \gamma^2)x' = D - X,} \quad (1.10)$$

as required.

Equation (1.5) is

$$\gamma \frac{dx}{ds} + \frac{dy}{ds} = 2(d + \lambda x).$$

Substituting into equation (1.5) for dx/ds from equation (1.4) gives,

$$\begin{aligned} \gamma \left(2(c - \lambda y) - \gamma \frac{dy}{ds} \right) + \frac{dy}{ds} &= 2(d + \lambda x), \\ 2\gamma(c - \lambda y) - \gamma^2 \frac{dy}{ds} + \frac{dy}{ds} &= 2(d + \lambda x), \\ \frac{dy}{ds} (1 - \gamma^2) &= 2(d + \lambda x) - 2\gamma(c - \lambda y), \\ \frac{dy}{ds} (1 - \gamma^2) &= 2d + 2\lambda x - 2\gamma c + 2\gamma \lambda y, \\ \frac{dy}{ds} (1 - \gamma^2) &= 2(d - \gamma c) + 2\lambda(x + \gamma y). \end{aligned}$$

$$\therefore \boxed{(1 - \gamma^2)y' = C + Y.} \quad (1.11)$$

as required.

Substituting for x' and y' from equations (1.10) and (1.11), respectively, into equation (1.9)

$$\frac{(D - X)^2}{(1 - \gamma^2)^2} + 2\gamma \frac{(D - X)(C + Y)}{(1 - \gamma^2)(1 - \gamma^2)} + \frac{(C + Y)^2}{(1 - \gamma^2)^2} = 1$$

Expanding the above and cross multiplying by $(1 - \gamma^2)^2$ gives,

$$\begin{aligned} D^2 - 2DX + X^2 + 2\gamma(D - X)(C + Y) + C^2 + 2CY + Y^2 &= (1 - \gamma^2)^2, \\ D^2 - 2DX + X^2 + (2\gamma D - 2\gamma X)(C + Y) + C^2 + 2CY + Y^2 &= (1 - \gamma^2)^2, \\ D^2 - 2DX + X^2 + 2\gamma CD + 2\gamma DY - 2\gamma CX - 2\gamma XY + C^2 + 2CY + Y^2 &= (1 - \gamma^2)^2, \\ X^2 + Y^2 - 2\gamma XY + C^2 + D^2 + 2\gamma CD - 2DX + 2\gamma DY - 2\gamma CX + 2CY &= (1 - \gamma^2)^2, \end{aligned}$$

Collecting terms in just X and just Y gives,

$$X^2 + Y^2 - 2\gamma XY + C^2 + D^2 + 2\gamma CD - 2X(D + \gamma C) + 2Y(C + \gamma D) = (1 - \gamma^2)^2. \quad (1.12)$$

Where,
 $D = 2(c - \gamma d)$,
 $X = 2\lambda(x - y)$ and
 $x' = \frac{dy}{ds}$.

Where,
 $C = 2(d - \gamma c)$,
 $Y = 2\lambda(x + \gamma y)$ and
 $y' = \frac{dy}{ds}$.

Now, from above (see margin notes)

$$\begin{aligned}(D + \gamma C) &= 2(c - \gamma d) + \gamma(2(d - \gamma c)), \\ &= 2c - 2\gamma d + 2\gamma d - 2\gamma^2 c, \\ &= 2c(1 - \gamma^2).\end{aligned}$$

$$\begin{aligned}(C + \gamma D) &= 2(d - \gamma c) + \gamma(2(c - \gamma d)), \\ &= 2d - 2\gamma c + 2\gamma c - 2\gamma^2 d, \\ &= 2d(1 - \gamma^2).\end{aligned}$$

So,

$$\begin{aligned}D + \gamma C &= 2c(1 - \gamma^2) \quad \text{and} \\ C + \gamma D &= 2d(1 - \gamma^2).\end{aligned}$$

Substituting these expressions into equation (1.12) gives,

$$X^2 + Y^2 - 2\gamma XY + C^2 + D^2 + 2\gamma CD - 4c(1 - \gamma^2)X + 4d(1 - \gamma^2)Y = (1 - \gamma^2)^2,$$

as required.

The boundary conditions are

$$x(0) = y(0) = 0, \quad x(1) = R > 0, \quad y(1) = 0.$$

Now, applying these boundary conditions to equation (1.12) and from the margin notes above we have:

$$X(t) = 2\lambda(\gamma x(t) + y(t)).$$

$$Y(t) = 2\lambda(x(t) + \gamma y(t)).$$

At $t = 0$,

$$X(0) = 2\lambda(\gamma x(0) + y(0)) = 0.$$

$$Y(0) = 2\lambda(x(0) + \gamma y(0)) = 0.$$

Thus, equation (1.12) at $t = 0$ reduces to

$$C^2 + D^2 + 2\gamma CD = (1 - \gamma^2)^2.$$

At $t = 1$,

$$X(1) = 2\lambda(\gamma x(1) + y(1)) = 2\lambda\gamma R.$$

$$Y(1) = 2\lambda(x(1) + \gamma y(1)) = 2\lambda R.$$

Thus, when applied to equation (1.12),

$$\begin{aligned}4\lambda^2\gamma^2 R^2 + 4\lambda^2 R^2 - 2\gamma(2\lambda\gamma R)(2\lambda R) - 4c(1 - \gamma^2)(2\lambda\gamma R) \\ + 4d(1 - \gamma^2)(2\lambda R) + C^2 + D^2 + 2\gamma CD = (1 - \gamma^2)^2.\end{aligned}$$

$$\begin{aligned}4\lambda^2\gamma^2 R^2 + 4\lambda^2 R^2 - 8\lambda^2\gamma^2 R^2 - 8c(1 - \gamma^2)\lambda\gamma R \\ + 8d(1 - \gamma^2)\lambda R + C^2 + D^2 + 2\gamma CD = (1 - \gamma^2)^2.\end{aligned}$$

Note from above that $C^2 + D^2 + 2\gamma CD = (1 - \gamma^2)^2$, thus

$$4\lambda^2\gamma^2R^2 + 4\lambda^2R^2 - 8\lambda^2\gamma^2R^2 - 8c(1 - \gamma^2)\lambda\gamma R + 8d(1 - \gamma^2)\lambda R = 0,$$

which, after dividing through by $4\lambda R$, gives

$$\lambda\gamma^2R + \lambda R - 2\lambda\gamma^2R - 2c(1 - \gamma^2)\gamma + 2d(1 - \gamma^2) = 0, \quad \lambda, R \neq 0 \text{ and } \gamma^2 \neq 1$$

$$\lambda R - \lambda\gamma^2R - 2c(1 - \gamma^2)\gamma + 2d(1 - \gamma^2) = 0,$$

$$\lambda R(1 - \gamma^2) - 2c(1 - \gamma^2)\gamma + 2d(1 - \gamma^2) = 0,$$

and dividing through by $(1 - \gamma^2)$, gives

$$\lambda R - 2c\gamma + 2d = 0.$$

Simplifying,

$$\lambda R - 2(c\gamma + d) = 0,$$

and noting that the quantity $2(c\gamma + d)$ is equal to C , then

$$\lambda R + C = 0.$$

Finally,

$$\boxed{C = -\lambda R, \quad R > 0,} \quad (1.13)$$

as required.

Recall that $C^2 + D^2 + 2\gamma CD = (1 - \gamma^2)^2$ and $C = -\lambda R$, $R > 0$.

Thus,

$$\lambda^2R^2 + D^2 + 2\gamma(-\lambda R)D = (1 - \gamma^2)^2,$$

$$D^2 - 2\lambda\gamma RD + \lambda^2R^2 - (1 - \gamma^2)^2 = 0.$$

Solving for D in the above quadratic,

$$D = \frac{2\lambda\gamma R \pm \sqrt{4\lambda^2\gamma^2R^2 - 4(\lambda^2R^2 - (1 - \gamma^2)^2)}}{2},$$

$$D = \lambda\gamma R \pm \sqrt{\lambda^2\gamma^2R^2 - \lambda^2R^2 + (1 - \gamma^2)^2},$$

$$D = \lambda\gamma R \pm \sqrt{\lambda^2R^2(\gamma^2 - 1) + (1 - \gamma^2)^2},$$

$$D = \lambda\gamma R \pm \sqrt{-\lambda^2R^2(1 - \gamma^2) + (1 - \gamma^2)^2}$$

It is seen that D will always have two real roots if $\gamma^2 > 1$, as in the expression below

$$D = \lambda\gamma R \pm \sqrt{-\lambda^2R^2(1 - \gamma^2) + (1 - \gamma^2)^2}$$

$(1 - \gamma^2)^2 > 0$ and $(1 - \gamma^2) < 0$ and consequently $-\lambda^2R^2(1 - \gamma^2) > 0$ making the entire quantity inside the radical positive.

Also, if $\gamma^2 < 1$ then $(1 - \gamma^2) > 0$ with $-\lambda^2 R^2 (1 - \gamma^2) < 0$ and the quantity $(1 - \gamma^2)^2 - \lambda^2 R^2 (1 - \gamma^2) > 0$ provided that $\lambda^2 R^2 (1 - \gamma^2) < (1 - \gamma^2)^2$, that is if $\lambda R < \sqrt{1 - \gamma^2}$, there will be two real roots of D .

Conversely, if $(1 - \gamma^2)^2 - \lambda^2 R^2 (1 - \gamma^2) < 0$ then $\lambda^2 R^2 (1 - \gamma^2) > (1 - \gamma^2)^2$, that is $\lambda R > \sqrt{1 - \gamma^2}$, there will not be any real solutions of D .

Q 2.

(a)

(b)

(c)

Q 3.

(a)

(b)

Q 4.

Q 5.

Q 6.

- (a) The model of tumour growth under radiotherapy from time $t = 0$ to $t = t_1 > 0$ is given as

$$\frac{dC}{dt} = -C \log \left(\frac{C}{C_{max}} \right) - \frac{DC}{1+D}, \quad (6.1)$$

where, at time t , $C(t)$ is the size of the tumour, $C_{max} > 0$ is the maximum size of the tumour, a constant, so that $0 < C(t) \leq C_{max}$, and $D(t) \geq 0$ is the rate at which the drug is administered.

Now, making a change of variable

$$x = \log \left(\frac{C}{C_{max}} \right),$$

then (6.1) becomes

$$\begin{aligned} \frac{dx}{dt} &= \frac{dC}{dt} \cdot \frac{dx}{dC}, \\ \frac{dx}{dt} &= \frac{dC}{dt} \cdot \frac{1}{C}, \\ \frac{dx}{dt} &= \left[-C \log \left(\frac{C}{C_{max}} \right) - \frac{DC}{1+D} \right] \cdot \frac{1}{C}, \\ \frac{dx}{dt} &= -\log \left(\frac{C}{C_{max}} \right) - \frac{D}{1+D}, \\ \text{thus, } \frac{dx}{dt} &= -x - \frac{D}{1+D}, \quad \text{where } -\infty < x \leq 0. \end{aligned}$$

- (b) We are told that:

It is required to reduce the size of the tumour from C_0 at $t = 0$ to C_1 at $t = t_1$. Let $x_0 = \log(C_0/C_{max})$ and $x_1 = \log(C_1/C_{max})$. For the health of the patient, it is desired to minimise the total amount of drug administered, which is given by the functional

$$S[D] = \int_0^{t_1} dt D(t).$$

So, by expressing D in terms of x and $\dot{x} = dx/dt$ it should be possible to show that $S[D]$ may be written as

$$S[x] = - \int_0^{t_1} dt \frac{\dot{x} + x}{1 + \dot{x} + x}.$$

From part (a)

$$\begin{aligned}\dot{x} &= -x - \frac{D}{1+D}, \\ \dot{x} + x &= -\frac{D}{1+D}, \\ (\dot{x} + x)(1+D) &= -D, \\ (\dot{x} + x) + D(\dot{x} + x) &= -D, \\ D(\dot{x} + x) + D &= -(\dot{x} + x), \\ D(\dot{x} + x + 1) &= -\dot{x} - x, \\ D &= \frac{-\dot{x} - x}{\dot{x} + x + 1}.\end{aligned}$$

So

$$S[x] = - \int_0^{t_1} dt \frac{\dot{x} + x}{\dot{x} + x + 1} \quad \text{as required.}$$

(c) To show that the Euler-Lagrange equation for $S[x]$ is given by

$$2\ddot{x} + 3\dot{x} + x = -1, \quad \text{where } \ddot{x} = d^2x/dt^2,$$

consider the following.

The Euler-Lagrange equation is given by

HB p17.

$$\frac{d}{dt} \left(\frac{\partial F}{\partial \dot{x}} \right) - \frac{\partial F}{\partial x} = 0, \quad y(a) = A, \quad y(b) = B.$$

$$\text{Let, } F = \frac{\dot{x} + x}{\dot{x} + x + 1},$$

then using the quotient rule to determine $\partial F/\partial \dot{x}$ and $\partial F/\partial x$

$$\frac{\partial}{\partial \dot{x}} \left(\frac{u}{v} \right) = \left[v \frac{\partial u}{\partial \dot{x}} - u \frac{\partial v}{\partial \dot{x}} \right] / v^2 \quad \text{where, } u = \dot{x} + x \text{ and } v = \dot{x} + x + 1.$$

$$\begin{aligned}\frac{\partial F}{\partial \dot{x}} &= \frac{(\dot{x} + x + 1) \cdot 1 - (\dot{x} + x) \cdot 1}{(\dot{x} + x + 1)^2} = \frac{1}{(\dot{x} + x + 1)^2}, \\ \frac{\partial F}{\partial x} &= \frac{(\dot{x} + x + 1) \cdot 1 - (\dot{x} + x) \cdot 1}{(\dot{x} + x + 1)^2} = \frac{1}{(\dot{x} + x + 1)^2}.\end{aligned}$$

$$\begin{aligned}\frac{d}{dt} \left(\frac{\partial F}{\partial \dot{x}} \right) &= \frac{d}{dt} (\dot{x} + x + 1)^{-2}, \\ &= -2(\dot{x} + x + 1)^{-3} (\ddot{x} + \dot{x}), \\ &= \frac{-2(\ddot{x} + \dot{x})}{(\dot{x} + x + 1)^3}.\end{aligned}$$

Therefore, the Euler-Lagrange equation is

$$\begin{aligned}
 -\frac{2(\ddot{x} + \dot{x})}{(\dot{x} + x + 1)^3} - \frac{1}{(\dot{x} + x + 1)^2} &= 0, \\
 \frac{2(\ddot{x} + \dot{x})}{(\dot{x} + x + 1)^3} &= -\frac{1}{(\dot{x} + x + 1)^2}, \\
 \frac{2(\ddot{x} + \dot{x})}{(\dot{x} + x + 1)} &= -1, \\
 2(\ddot{x} + \dot{x}) &= -(\dot{x} + x + 1), \\
 2(\ddot{x} + \dot{x}) + (\dot{x} + x) &= -1, \\
 2\ddot{x} + 3\dot{x} + x &= -1, \quad \text{as required.}
 \end{aligned} \tag{6.2} \quad \ddot{x} = d^2x/dt^2.$$

- (d) Now solving the second-order linear ordinary differential equation (6.2) with the boundary conditions $x(0) = x_0$ and $x(t_1) = x_1$ as follows.

The auxiliary equation is

$$2\lambda^2 + 3\lambda + 1 = 0.$$

This has roots $\lambda_1 = -1$ and $\lambda_2 = -\frac{1}{2}$. These roots are real and different so the complementary function is given by

$$y_c = c_1 e^{-t} + c_2 e^{-\frac{1}{2}t}, \quad \text{where } c_1 \text{ and } c_2 \text{ are arbitrary constants.}$$

Using the method of undetermined coefficients to determine the particular solution to (6.2) as follows.

The particular solution to

$$2\frac{d^2x(t)}{dt^2} + 3\frac{dx(t)}{dt} + x(t) = -1,$$

is of the form

$$x_p(t) = a_1,$$

where a_1 is an unknown constant to be determined as follows:

$$\frac{dx(t)}{dt} = \frac{d}{dt}(a_1) = 0,$$

and therefore

$$\frac{d^2x(t)}{dt^2} = 0.$$

Substituting the particular solution $x_p(t)$ in to (6.2) gives

$$2 \times 0 + 3 \times 0 + a_1 = -1.$$

Therefore, $a_1 = -1$ and the general solution is

$$x(t) = c_1 e^{-t} + c_2 e^{-\frac{1}{2}t} - 1. \tag{6.3}$$

Now using the boundary conditions to solve for the unknown constants c_1 and c_2 as follows.

At $t = 0$, $x(0) = x_0$ and substituting this condition into (6.3) gives,

$$c_1 + c_2 - 1 = x_0. \quad (6.4)$$

At $t = t_1$, $x(t_1) = x_1$ and substituting this condition into (6.3) gives,

$$c_1 e^{-t_1} + c_2 e^{-\frac{1}{2}t_1} - 1 = x_1 \quad (6.5)$$

From (6.4)

$$c_1 = (x_0 + 1) - c_2 \quad (6.6)$$

Substituting into (6.5) for c_1 from (6.6) and rearranging in terms of c_2

$$\begin{aligned} (x_0 + 1) e^{-t_1} - c_2 e^{-t_1} + c_2 e^{-\frac{1}{2}t_1} &= x_1 + 1, \\ (x_0 + 1) e^{-t_1} + c_2 \left(e^{-\frac{1}{2}t_1} - e^{-t_1} \right) &= (x_1 + 1), \\ c_2 \left(e^{-\frac{1}{2}t_1} - e^{-t_1} \right) &= (x_1 + 1) - (x_0 + 1) e^{-t_1}, \\ c_2 &= \frac{(x_1 + 1) - (x_0 + 1) e^{-t_1}}{\left(e^{-\frac{1}{2}t_1} - e^{-t_1} \right)}. \end{aligned} \quad (6.7)$$

Substituting for c_2 in (6.6) from (6.7)

$$\begin{aligned} c_1 &= (x_0 + 1) - \left[\frac{(x_1 + 1) - (x_0 + 1) e^{-t_1}}{\left(e^{-\frac{1}{2}t_1} - e^{-t_1} \right)} \right], \\ c_1 &= \frac{(x_0 + 1) \left(e^{-\frac{1}{2}t_1} - e^{-t_1} \right) - (x_1 + 1) + (x_0 + 1) e^{-t_1}}{\left(e^{-\frac{1}{2}t_1} - e^{-t_1} \right)} \\ c_1 &= \frac{(x_0 + 1) e^{-\frac{1}{2}t_1} - (x_0 + 1) e^{-t_1} - (x_1 + 1) + (x_0 + 1) e^{-t_1}}{\left(e^{-\frac{1}{2}t_1} - e^{-t_1} \right)} \\ c_1 &= \frac{(x_0 + 1) e^{-\frac{1}{2}t_1} - (x_1 + 1)}{\left(e^{-\frac{1}{2}t_1} - e^{-t_1} \right)} = \alpha. \end{aligned} \quad (6.8)$$

Therefore, from (6.4)

$$c_2 = (x_0 + 1) - \alpha. \quad (6.9)$$

Substituting into (6.3) for c_1 and c_2 from (6.8) and (6.9) respectively gives,

$$x(t) = \alpha e^{-t} + (x_0 + 1 - \alpha) e^{-\frac{1}{2}t} - 1 \quad (\alpha \text{ given in (6.8)}),$$

as required.
