

Performance of Empirical Models

Ravi Kothari, Ph.D.
ravi.kothari@ashoka.edu.in

“Correlation is not Causation..”

Aspects of Empirical Models

Aspects of Empirical Models

- Model complexity

Aspects of Empirical Models

- Model complexity
- Prediction error

Aspects of Empirical Models

- Model complexity
- Prediction error
 - ▶ Average case

Aspects of Empirical Models

- Model complexity
- Prediction error
 - ▶ Average case
 - ▶ Boosting

Aspects of Empirical Models

- Model complexity
- Prediction error
 - ▶ Average case
 - ▶ Boosting
 - ▶ Minimum Description Length

Aspects of Empirical Models

- Model complexity
- Prediction error
 - ▶ Average case
 - ▶ Boosting
 - ▶ Minimum Description Length
 - ▶ Uniform convergence


Aspects of Empirical Models

- Model complexity
- Prediction error
 - ▶ Average case
 - ▶ Boosting
 - ▶ Minimum Description Length
 - ▶ Uniform convergence
 - ▶ Worst case

Estimating the Prediction Error (1/2)


Estimating the Prediction Error (1/2)

- Sampling *without replacement* methods e.g. k -fold cross validation




Estimating the Prediction Error (1/2)

- Sampling *without replacement* methods e.g. k -fold cross validation




Estimating the Prediction Error (1/2)

- Sampling *without replacement* methods e.g. k -fold cross validation



Estimating the Prediction Error (1/2)


- Sampling *without replacement* methods e.g. k -fold cross validation



- Error estimate is the average of the k -error estimates

Estimating the Prediction Error (1/2)


- Sampling *without replacement* methods e.g. k -fold cross validation



- Error estimate is the average of the k -error estimates
- When $k = N$, we get the leave-one-out estimate

Estimating the Prediction Error (1/2)

- Sampling *without replacement* methods e.g. k -fold cross validation



- Error estimate is the average of the k -error estimates
- When $k = N$, we get the leave-one-out estimate
- Time consuming, unbiased estimate with large variance

Estimating the Prediction Error (2/2)

Estimating the Prediction Error (2/2)

- Sampling *with* replacement e.g. bootstrap

Estimating the Prediction Error (2/2)

- Sampling *with* replacement e.g. bootstrap
- Probability that a pattern is chosen is $(1 - (1 - 1/N)^N)$. For large N , this approaches $(1 - e^{-1}) = 0.632$

Estimating the Prediction Error (2/2)

- Sampling *with* replacement e.g. bootstrap
- Probability that a pattern is chosen is $(1 - (1 - 1/N)^N)$. For large N , this approaches $(1 - e^{-1}) = 0.632$
- Patterns not chosen in the i^{th} bootstrap sample become part of the i^{th} test set

Estimating the Prediction Error (2/2)

- Sampling *with* replacement e.g. bootstrap
- Probability that a pattern is chosen is $(1 - (1 - 1/N)^N)$. For large N , this approaches $(1 - e^{-1}) = 0.632$
- Patterns not chosen in the i^{th} bootstrap sample become part of the i^{th} test set
- Induce the model using the i^{th} bootstrap sample. Get the error ϵ_i from the i^{th} test set. Repeat b times

Estimating the Prediction Error (2/2)

- Sampling *with* replacement e.g. bootstrap
- Probability that a pattern is chosen is $(1 - (1 - 1/N)^N)$. For large N , this approaches $(1 - e^{-1}) = 0.632$
- Patterns not chosen in the i^{th} bootstrap sample become part of the i^{th} test set
- Induce the model using the i^{th} bootstrap sample. Get the error ϵ_i from the i^{th} test set. Repeat b times

$$J_{\text{boot}} = \frac{1}{b} \sum_{i=1}^b (0.632\epsilon_i + 0.368J_{\text{total}})$$

Average Case Analysis – The Bias-Variance Decomposition

Optimal Response of an Estimator

Optimal Response of an Estimator

- $y = f(x) + \epsilon$. So, the y obtained corresponding to t repeated observations of x are $y(1), y(2), \dots, y(t)$

Optimal Response of an Estimator

- $y = f(x) + \epsilon$. So, the y obtained corresponding to t repeated observations of x are $y(1), y(2), \dots, y(t)$
- What should the *optimal* estimator $\hat{f}^*(x; \theta)$ respond with?

Optimal Response of an Estimator

- $y = f(x) + \epsilon$. So, the y obtained corresponding to t repeated observations of x are $y(1), y(2), \dots, y(t)$
- What should the *optimal* estimator $\hat{f}^*(x; \theta)$ respond with?
-

$$J = \left(\hat{f}^*(x; \theta) - y(1) \right)^2 + \left(\hat{f}^*(x; \theta) - y(2) \right)^2 + \dots + \left(\hat{f}^*(x; \theta) - y(t) \right)^2$$

Optimal Response of an Estimator

- $y = f(x) + \epsilon$. So, the y obtained corresponding to t repeated observations of x are $y(1), y(2), \dots, y(t)$
- What should the *optimal* estimator $\hat{f}^*(x; \theta)$ respond with?

-

$$J = \left(\hat{f}^*(x; \theta) - y(1) \right)^2 + \left(\hat{f}^*(x; \theta) - y(2) \right)^2 + \dots + \left(\hat{f}^*(x; \theta) - y(t) \right)^2$$

- Minimum of J is achieved when $\hat{f}^*(x; \theta) = (y(1), y(2), \dots, y(t))/t$, i.e. $E[y|x]$

The Bias-Variance Decomposition

The Bias-Variance Decomposition

- Average case analysis of the prediction (generalization) error

The Bias-Variance Decomposition

- Average case analysis of the prediction (generalization) error

-

$$\begin{aligned} E_{\mathcal{X}} \left[\left(\hat{f}(x; \mathcal{X}) - E[y|x] \right)^2 \right] \\ &= E_{\mathcal{X}} \left[\left(\left(\hat{f}(x; \mathcal{X}) - E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] \right) + \right. \right. \\ &\quad \left. \left. \left(E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] - E[y|x] \right) \right)^2 \right] \\ &= \underbrace{\left(E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] - E[y|x] \right)^2}_{\text{Squared Bias}} + \underbrace{E_{\mathcal{X}} \left[\left(\hat{f}(x; \mathcal{X}) - E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] \right)^2 \right]}_{\text{Variance}} \end{aligned}$$

Bias

Bias

- The Bias term,

$$\underbrace{\left(E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] - E[y|x] \right)^2}_{\text{Squared Bias}}$$

Bias

- The Bias term,

$$\underbrace{\left(E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] - E[y|x] \right)^2}_{\text{Squared Bias}}$$

- Measures deviation of the averaged estimator output from the averaged system output

Bias

- The Bias term,

$$\underbrace{\left(E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] - E[y|x] \right)^2}_{\text{Squared Bias}}$$

- Measures deviation of the averaged estimator output from the averaged system output
- Bias is 0 even when a particular estimator has a large error which is canceled out by an opposite error generated by another model

Variance

Variance

- The Variance term,

$$\underbrace{E_{\mathcal{X}} \left[\left(\hat{f}(x; \mathcal{X}) - E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] \right)^2 \right]}_{\text{Variance}}$$

Variance

- The Variance term,

$$\underbrace{E_{\mathcal{X}} \left[\left(\hat{f}(x; \mathcal{X}) - E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] \right)^2 \right]}_{\text{Variance}}$$

- Measures the sensitivity of the estimator

Variance

- The Variance term,

$$\underbrace{E_{\mathcal{X}} \left[\left(\hat{f}(x; \mathcal{X}) - E_{\mathcal{X}} \left[\hat{f}(x; \mathcal{X}) \right] \right)^2 \right]}_{\text{Variance}}$$

- Measures the sensitivity of the estimator
- It is independent of the underlying system $f(x)$

Understanding Bias and Variance

Understanding Bias and Variance

- Let the estimator be k -nearest neighbor

Understanding Bias and Variance

- Let the estimator be k -nearest neighbor
- When $k = N$, the output is simply the average of the training set output i.e. $(1/N) \sum_{i=1}^N y^{(i)}$
 - ▶ Estimate is likely to be unchanged from one training data set to another. Bias is high but variance is low

Understanding Bias and Variance

- Let the estimator be k -nearest neighbor
- When $k = N$, the output is simply the average of the training set output i.e. $(1/N) \sum_{i=1}^N y^{(i)}$
 - ▶ Estimate is likely to be unchanged from one training data set to another. Bias is high but variance is low
- If $k = 1$, the output follows local changes (as opposed to the population behavior)
 - ▶ Indeed, when $N \rightarrow \infty$ then Bias $\rightarrow 0$. Bias is low but variance will be high

Understanding Bias and Variance

- Let the estimator be k -nearest neighbor
- When $k = N$, the output is simply the average of the training set output i.e. $(1/N) \sum_{i=1}^N y^{(i)}$
 - ▶ Estimate is likely to be unchanged from one training data set to another. Bias is high but variance is low
- If $k = 1$, the output follows local changes (as opposed to the population behavior)
 - ▶ Indeed, when $N \rightarrow \infty$ then Bias $\rightarrow 0$. Bias is low but variance will be high
- The best solution is usually some intermediate k

The Bias/Variance Trade-Off

The Bias/Variance Trade-Off

- Bias and Variance are complementary, i.e. reducing one (most often) increases the other

The Bias/Variance Trade-Off

- Bias and Variance are complementary, i.e. reducing one (most often) increases the other
- The trick is to allow one to increase if the other decreases more than the increase

The Bias/Variance Trade-Off

- Bias and Variance are complementary, i.e. reducing one (most often) increases the other
- The trick is to allow one to increase if the other decreases more than the increase
- Approaches like weight decay, pruning, growing, early stopping are based on the above premise (avoid overfitting, i.e. tolerate increased bias in the *hope* that variance reduces more)

The Bias/Variance Trade-Off

- Bias and Variance are complementary, i.e. reducing one (most often) increases the other
- The trick is to allow one to increase if the other decreases more than the increase
- Approaches like weight decay, pruning, growing, early stopping are based on the above premise (avoid overfitting, i.e. tolerate increased bias in the *hope* that variance reduces more)
- Other approaches are based on aggregation (e.g. bagging – bootstrap aggregating, boosting)

Boosting (1/2)

Boosting (1/2)

- AdaBoost for a two-class classification setting,

Boosting (1/2)

- AdaBoost for a two-class classification setting,
 - ▶ Create T models each trained with a different distribution on the training set,

$$F(x) = \text{sign} \left(\sum_{t=1}^T \alpha_t \hat{f}_t(x; \theta_t) \right)$$

Boosting (1/2)

- AdaBoost for a two-class classification setting,
 - ▶ Create T models each trained with a different distribution on the training set,

$$F(x) = \text{sign} \left(\sum_{t=1}^T \alpha_t \hat{f}_t(x; \theta_t) \right)$$

- ▶ Initially, weight $D_1^{(i)}$ on each pattern is $1/N$

Boosting (1/2)

- AdaBoost for a two-class classification setting,
 - ▶ Create T models each trained with a different distribution on the training set,

$$F(x) = \text{sign} \left(\sum_{t=1}^T \alpha_t \hat{f}_t(x; \theta_t) \right)$$

- ▶ Initially, weight $D_1^{(i)}$ on each pattern is $1/N$
 - ▶ Train using the distribution D_t

Boosting (2/2)

Boosting (2/2)

- Get error rate ϵ_t as,

$$\epsilon_t = \frac{1}{N} \sum_{i=1}^N I \left[\hat{f}_i(x^{(i)}; \Theta) \neq y^{(i)} \right]$$

$$\alpha_t = \log \left(\frac{1 - \epsilon_t}{\epsilon_t} \right)$$

- Reassign weightage on each pattern as,

$$D_{t+1}(i) = \frac{D_t}{Z_t} \times \begin{cases} e^{-\alpha_t} & \text{if } \hat{f}_i(x^{(i)}; \theta_i) = y^{(i)} \\ e^{\alpha_t} & \text{if } \hat{f}_i(x^{(i)}; \theta_i) \neq y^{(i)} \end{cases}$$

Minimum Description Length

Minimum Description Length

$$\underbrace{L(D)}_{\text{Description Length}} = \underbrace{L(D|H)}_{\text{Perturbation}} + \underbrace{L(H)}_{\text{Complexity (Nominal Model)}} \quad (1)$$

Minimum Description Length

$$\underbrace{L(D)}_{\text{Description Length}} = \underbrace{L(D|H)}_{\text{Perturbation}} + \underbrace{L(H)}_{\text{Complexity (Nominal Model)}} \quad (1)$$

- In ffnn, we could say the weights are $\mathcal{N}(0, \sigma_1)$ i.e. the model is $\exp(-\|w\|^2)/(2\sigma_1^2)$

Minimum Description Length

$$\underbrace{L(D)}_{\text{Description Length}} = \underbrace{L(D|H)}_{\text{Perturbation}} + \underbrace{L(H)}_{\text{Complexity (Nominal Model)}} \quad (1)$$

- In ffnn, we could say the weights are $\mathcal{N}(0, \sigma_1)$ i.e. the model is $\exp(-\|w\|^2)/(2\sigma_1^2)$
- Say, perturbation based on the data is $\mathcal{N}(0, \sigma_2)$ i.e. $\exp(-\|\epsilon\|^2)/(2\sigma_2^2)$

Minimum Description Length

$$\underbrace{L(D)}_{\text{Description Length}} = \underbrace{L(D|H)}_{\text{Perturbation}} + \underbrace{L(H)}_{\text{Complexity (Nominal Model)}} \quad (1)$$

- In ffnn, we could say the weights are $\mathcal{N}(0, \sigma_1)$ i.e. the model is $\exp(-\|w\|^2)/(2\sigma_1^2)$
- Say, perturbation based on the data is $\mathcal{N}(0, \sigma_2)$ i.e. $\exp(-\|\epsilon\|^2)/(2\sigma_2^2)$
- The total description length is,

$$\begin{aligned} L(D) &= -\log(D|H) - \log(H) \\ &= \frac{1}{2\sigma_2^2} (y^{(i)} - \hat{f}(x^{(i)}; \theta))^2 + \frac{1}{2\sigma_1^2} \|w\|^2 \end{aligned}$$

Worst Case Analysis – The VC Dimension

Empirical and True Risk

Empirical and True Risk

- Let $\mathbf{x} \in \{-1, +1\}^n$. Then,

$$J_{\text{emp}}(\theta) = \frac{1}{N} \sum_{i=1}^N \left(y^{(i)} - \hat{f}(\mathbf{x}^{(i)}; \theta) \right)^2$$

$$J(\theta) = E \left[\left(y - \hat{f}(\mathbf{x}; \mathcal{X}) \right)^2 \right]$$

Empirical and True Risk

- Let $x \in \{-1, +1\}^n$. Then,

$$J_{\text{emp}}(\theta) = \frac{1}{N} \sum_{i=1}^N \left(y^{(i)} - \hat{f}(x^{(i)}; \theta) \right)^2$$

$$J(\theta) = E \left[\left(y - \hat{f}(x; \mathcal{X}) \right)^2 \right]$$

- $J(\theta) \gg J_{\text{emp}}(\theta)$. From the law of large numbers,

$$\Pr[|J(\theta) - J_{\text{emp}}(\theta)| > \epsilon] \rightarrow 0, \text{ as } N \rightarrow \infty$$

Uniform Convergence

Uniform Convergence

- However the stronger result also holds,

$$\Pr \left[\sup_{\theta} |J(\theta) - J_{\text{emp}}(\theta)| > \epsilon \right] \rightarrow 0, \text{ as } N \rightarrow \infty$$

Uniform Convergence

- However the stronger result also holds,

$$\Pr \left[\sup_{\theta} |J(\theta) - J_{\text{emp}}(\theta)| > \epsilon \right] \rightarrow 0, \text{ as } N \rightarrow \infty$$

- This is the uniform convergence of the empirical error rate to the true error rate

Uniform Convergence

- However the stronger result also holds,

$$\Pr \left[\sup_{\theta} |J(\theta) - J_{\text{emp}}(\theta)| > \epsilon \right] \rightarrow 0, \text{ as } N \rightarrow \infty$$

- This is the uniform convergence of the empirical error rate to the true error rate
- What is the rate of uniform convergence?

Rate of Uniform Convergence

Rate of Uniform Convergence

- However the stronger result also holds,

$$\Pr \left[\sup_{\theta} |J(\theta) - J_{\text{emp}}(\theta)| > \epsilon \right] < 4 \Delta(2N) e^{-\frac{\epsilon^2 N}{8}}$$

Rate of Uniform Convergence

- However the stronger result also holds,

$$\Pr \left[\sup_{\theta} |J(\theta) - J_{\text{emp}}(\theta)| > \epsilon \right] < 4 \Delta(2N) e^{-\frac{\epsilon^2 N}{8}}$$

All realizations

Growth function

Rate of Uniform Convergence

- However the stronger result also holds,

$$\Pr \left[\sup_{\theta} |J(\theta) - J_{\text{emp}}(\theta)| > \epsilon \right] < 4 \Delta(2N) e^{-\frac{\epsilon^2 N}{8}}$$

All realizations

Growth function

- This is the uniform convergence of the empirical error rate to the true error rate


VC Dimension

VC Dimension

- $\Delta(2N)$ is identically equal to 2^N or bounded above by $\Delta(N) \leq N^{d_{\text{vc}}} + 1$ i.e. the machine can shatter upto d_{vc} points in a general position (realize all possible dichotomies)


VC Dimension

- $\Delta(2N)$ is identically equal to 2^N or bounded above by $\Delta(N) \leq N^{d_{\text{vc}}} + 1$ i.e. the machine can shatter upto d_{vc} points in a general position (realize all possible dichotomies)



VC Dimension

- $\Delta(2N)$ is identically equal to 2^N or bounded above by $\Delta(N) \leq N^{d_{VC}} + 1$ i.e. the machine can shatter upto d_{VC} points in a general position (realize all possible dichotomies)



- The estimator can generalize beyond d_{VC} – the VC dimension


VC Dimension – Example

VC Dimension – Example

- d_{VC} for a linear classifier in n dimensions is $(n + 1)$


VC Dimension – Example

- d_{VC} for a linear classifier in n dimensions is $(n + 1)$



VC Dimension – Example

- d_{VC} for a linear classifier in n dimensions is $(n + 1)$



- The estimator can generalize beyond d_{VC} – the VC dimension

Linear Classifier

Linear Classifier

$$\begin{aligned}\Pr \left[\sup_{\theta} |J(\theta) - J_{\text{emp}}(\theta)| > \epsilon \right] &< 4 \Delta(2N) e^{-\frac{\epsilon^2 N}{8}} \\ &< 4 \left[(2N)^{d_{\text{vc}}} + 1 \right] e^{-\frac{\epsilon^2 N}{8}} \\ &< 4 \left[(2N)^{n+1} + 1 \right] e^{-\frac{\epsilon^2 N}{8}}\end{aligned}$$

Linear Classifier

$$\begin{aligned}\Pr \left[\sup_{\theta} |J(\theta) - J_{\text{emp}}(\theta)| > \epsilon \right] &< 4 \Delta(2N) e^{-\frac{\epsilon^2 N}{8}} \\ &< 4 \left[(2N)^{d_{\text{vc}}} + 1 \right] e^{-\frac{\epsilon^2 N}{8}} \\ &< 4 \left[(2N)^{n+1} + 1 \right] e^{-\frac{\epsilon^2 N}{8}}\end{aligned}$$

- For right side to be small, approximately,

$$N > \frac{8n \log n}{\epsilon^2}$$

Some VC Dimensions

Some VC Dimensions

- Intervals in \mathcal{R} : 2

Some VC Dimensions

- Intervals in \mathcal{R} : 2
- Axis parallel rectangles in \mathcal{R}^2 : 4

Some VC Dimensions

- Intervals in \mathcal{R} : 2
- Axis parallel rectangles in \mathcal{R}^2 : 4
- Circle in \mathcal{R}^2 : 3

Some VC Dimensions

- Intervals in \mathcal{R} : 2
- Axis parallel rectangles in \mathcal{R}^2 : 4
- Circle in \mathcal{R}^2 : 3
- Triangle in \mathcal{R}^2 : 7

Conclusion

Conclusion

- Empirical models need to be carefully designed and used to realize their advantages

Conclusion

- Empirical models need to be carefully designed and used to realize their advantages
- B^2V , VC-Dimension or MDL are good points for designing/analyzing new algorithms