

Lab 8

CALCULUS FOR IT 501031

1 Exercises

Exercise 1: Find the critical numbers (C.N) of $f(x)$ for the following cases:

(a) $f(x) = 18x^2 - 9$

(c) $f(x) = -\frac{x^2}{3} + x^2 + 3x + 4$

(b) $f(x) = \frac{x+2}{2x^2}$

(d) $f(x) = \frac{5x^2+5}{x}$

Exercise 2: Find the relative extrema using the second derivative test for the following cases:

(a) $f(x) = 18x^2 - 9$

(c) $f(x) = -\frac{x^2}{3} + x^2 + 3x + 4$

(b) $f(x) = \frac{x+2}{2x^2}$

(d) $f(x) = \frac{5x^2+5}{x}$

Exercise 3: Given $f(x)$ over a closed interval $[a, b]$, find the absolute maximum and the absolute minimum for the following cases:

(a) $f(x) = x^3 - 27x, [0, 5]$

(c) $f(x) = \frac{1}{2}x^4 - 4x^2 + 5, [1, 3]$

(b) $f(x) = \frac{3}{2}x^4 - 4x^3 + 4, [0, 3]$

(d) $f(x) = \frac{5}{2}x^4 - \frac{20}{3}x^3 + 6, [-1, 3]$

Exercise 4: Determine the minima or maxima of the functions $f(x)$ following:

(a) $f(x) = x^2 - 2x - 5, a = 0, b = 2$

(d) $f(x) = e^{x^2} + 3x, a = -1, b = 1$

(b) $f(x) = 3x + x^3 + 5, a = -4, b = 4$

(e) $f(x) = x^3 - 3x, a = -3, b = 0$

(c) $f(x) = \sin(x) + 3x^2, a = -2, b = 2$

(f) $f(x) = x^3 - 3x, a = 0, b = 3$

Illustrate $f(x)$ and mark the maximum point on graph.

Exercise 5: Write a program to implement **Golden Search** and apply to determinate minimum value of $f(x) = x^2$ in $[-2, 1]$, with a tolerate $\epsilon = 0.3$, and illustrate on the graph/ table for each iteration.

Exercise 6: Implement **Fibonacci Search** and apply to determinate minimum value of $f(x) = x^2$ in $[-2, 1]$, with a tolerate $\epsilon = 0.3$, and illustrate on the graph/ table for each iteration.

Exercise 7: Determine m to $y = x^3 - 3mx^2 + 3(m^2 - 1)x - (m^2 - 1)$ maximize at $x_0 = 1$

Algorithm 1 Golden Search

Input: Objective function $f(x)$, boundaries a and b , and tolerance ϵ

```

 $d = b - a$ 
while  $b - a \geq \epsilon$  do
     $d \leftarrow 0.618 \times d$ 
     $x_1 \leftarrow b - d$ 
     $x_2 \leftarrow a + d$ 
    if  $f(x_1) \leq f(x_2)$  then
         $b \leftarrow x_2$ 
    else
         $a \leftarrow x_1$ 
    end if
end while

```

Output: Reduced interval $[a, b]$

Algorithm 2 Fibonacci Search

Input: Objective function $f(x)$, boundaries a and b , and tolerance ϵ

```

 $F_1 = 2, F_2 = 3$ 

 $n = 2$  while  $b - a \geq \epsilon$  do
     $d \leftarrow b - a$ 
     $x_1 \leftarrow b - d \frac{F_{n-1}}{F_n}$ 
     $x_2 \leftarrow a + d \frac{F_{n-1}}{F_n}$ 
    if  $f(x_1) \leq f(x_2)$  then
         $b \leftarrow x_2$ 
    else
         $a \leftarrow x_1$ 
    end if

```

```

 $n = n + 1$ 
 $F_n = F_{n-1} + F_{n-2}$  end while
Output: Reduced interval  $[a, b]$ 

```

Exercise 8: Find the minimum value of $f(x)$ using Golden Search and Fibonacci Search, respectively. Next, present on the graphs.

- (a) $f(x) = -2x^2 + x + 4$, in $[-5, 5]$, and $\epsilon = \frac{1}{9}$
- (b) $f(x) = -4x^2 + 2x + 2$, in $[-6, 6]$, and $\epsilon = \frac{1}{10}$
- (c) $f(x) = x^3 + 6x^2 + 5x - 12$, in $[-5, -2]$, and $\epsilon = \frac{1}{10}$
- (d) $f(x) = 2x - x^2$, in $[0, 3]$, and $\epsilon = \frac{1}{100}$
- (e) $f(x) = x^2 - x - 10$, in $[-10, 10]$, and $\epsilon = \frac{1}{5}$