

Understanding Analysis

Solution of exercise problems.

Quasar Chunawala

Version 1.0.0, last revised on 2021-05-13.

Abstract

This is a solution manual for Understanding Analysis, 2nd edition, by Stephen Abbott.

Chapter 3. Basic Topology of \mathbf{R}

Theorem. The open interval $(0, 1) = \{x \in \mathbf{R} : 0 < x < 1\}$ is uncountable.

Proof.

We proceed by contradiction and assume that there exists a function $f : \mathbf{N} \rightarrow (0, 1)$ that is $1-1$ and onto. $1-1$ implies that distinct elements have distinct images. Onto implies that every element in the co-domain has atleast one pre-image. For each $m \in \mathbf{N}$, $f(m)$ is a real number between 0 and 1, and we represent it using the decimal notation

$$f(m) = .a_{m1}a_{m2}a_{m3}a_{m4}a_{m5} \dots$$

What is meant here is that for each $m, n \in \mathbf{N}$, a_{mn} is the digit from the set $\{0, 1, 2, 3, \dots, 9\}$ that represents the n th digit in the decimal expansion of m th real number, $f(m)$. The $1-1$ correspondence between \mathbf{N} and $(0, 1)$ can be summarized in the doubly indexed array:

N			(0, 1)							
1	\longleftrightarrow	$f(1)$	=	.	a ₁₁	a_{12}	a_{13}	a_{14}	a_{15}	a_{16} ...
2	\longleftrightarrow	$f(2)$	=	.	a_{21}	a ₂₂	a_{23}	a_{24}	a_{25}	a_{26} ...
3	\longleftrightarrow	$f(3)$	=	.	a_{31}	a_{32}	a ₃₃	a_{34}	a_{35}	a_{36} ...
4	\longleftrightarrow	$f(4)$	=	.	a_{41}	a_{42}	a_{43}	a ₄₄	a_{45}	a_{46} ...
5	\longleftrightarrow	$f(5)$	=	.	a_{51}	a_{52}	a_{53}	a_{54}	a ₅₅	a_{56} ...
6	\longleftrightarrow	$f(6)$	=	.	a_{61}	a_{62}	a_{63}	a_{64}	a_{65}	a ₆₆ ...
\vdots		\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots

The key assumption about this correspondence is that **every** real number in $(0, 1)$ is assumed to appear somewhere on this list.

Now for the pearl of the argument. Define a real number $x \in (0, 1)$ with the decimal expansion $x = .b_1b_2b_3b_4 \dots$ using the rule

$$b_n = \begin{cases} 2 & \text{if } a_{nn} \neq 2 \\ 3 & \text{if } a_{nn} = 2 \end{cases}$$

Now, the real number $x = .b_1b_2b_3b_4 \dots$ cannot be $f(1)$, simply because its first digit b_1 differs from the first digit a_{11} of $f(1)$. Similarly, the second digit b_2 differs from the second digit a_{22} of $f(2)$. In general, the n th digit of x differs from the n th digit of $f(n)$. So, we have constructed a real number x that is not in the set $\{f(1), f(2), f(3), \dots, f(n)\}$. But, this is a contradiction. Hence, our initial assumption is false. The set of real numbers in $(0, 1)$ are uncountable.

Exercise. [Abbott, 1.6.4] Let S be the set consisting of all sequences of 0s and 1s. Observe that S is not a particular sequence, but rather a large set whose elements are sequences, namely:

$$S = \{(a_1, a_2, a_3, \dots) : a_n = 0 \text{ or } 1\}$$

As an example, the binary sequence $(1, 0, 1, 0, 1, 0, 1, 0, \dots)$ is an element of S as is the sequence $(1, 1, 1, 1, 1, 1, \dots)$. Give a rigorous argument showing that S is uncountable.

Proof.

Suppose that S - the set of all possible binary strings of infinite length is countable. Then, we can define a bijection $f: \mathbf{N} \rightarrow S$ between the natural numbers and S . For each $m \in \mathbf{N}$, $f(m)$ is a binary string in S . Let us enlist the first few elements of this correspondence.

N		(0, 1)								
1	\longleftrightarrow	$f(1)$	=	.	a₁₁	a_{12}	a_{13}	a_{14}	a_{15}	a_{16} ...
2	\longleftrightarrow	$f(2)$	=	.	a_{21}	a₂₂	a_{23}	a_{24}	a_{25}	a_{26} ...
3	\longleftrightarrow	$f(3)$	=	.	a_{31}	a_{32}	a₃₃	a_{34}	a_{35}	a_{36} ...
4	\longleftrightarrow	$f(4)$	=	.	a_{41}	a_{42}	a_{43}	a₄₄	a_{45}	a_{46} ...
5	\longleftrightarrow	$f(5)$	=	.	a_{51}	a_{52}	a_{53}	a_{54}	a₅₅	a_{56} ...
6	\longleftrightarrow	$f(6)$	=	.	a_{61}	a_{62}	a_{63}	a_{64}	a_{65}	a₆₆ ...
\vdots		\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\ddots

Define a binary sequence $x = (b_1, b_2, b_3, b_4, \dots)$ such that

$$b_i = \begin{cases} 1 & \text{if } a_{ii} = 0 \\ 0 & \text{if } a_{ii} = 1 \end{cases}$$

Thus, the binary sequence $x = (b_1, b_2, b_3, b_4, \dots)$ has atleast one bit that differs from all of the elements in S . Consequently, $x \notin S$. This is a contradiction, as S is supposed to contain all binary strings. Hence, S is not countable.

3.1 Discussion: The Cantor Set.

What follows is a fascinating mathematical construction, due to Georg Cantor, which is extremely useful for extending the horizons of our intuition about the nature of subsets of the real line. Cantor's name has already appeared in the first chapter in our discussion of uncountable sets. Indeed, Cantor's proof that \mathbf{R} is uncountable occupies another spot on the short list of the most significant contrubutions towards the understanding of the mathematical infinite. In the words of the mathematician David Hilbert, "No one shall expel us from the paradise that Cantor has created for us."

Let C_0 be the closed interval $[0, 1]$ and define C_1 to be the set that results when the open middle one third is removedl that is,


$$C_1 = C_0 \setminus \left(\frac{1}{3}, \frac{2}{3} \right) = \left[0, \frac{1}{3} \right] \cup \left[\frac{2}{3}, 1 \right]$$

Now, construct C_2 in a similar way by removing the open middle third of each of the two components of C_1 :

$$C_2 = \left(\left[0, \frac{1}{9} \right] \cup \left[\frac{2}{9}, \frac{3}{9} \right] \right) \cup \left(\left[\frac{6}{9}, \frac{7}{9} \right] \cup \left[\frac{8}{9}, 1 \right] \right)$$

If we continue this process inductively then for each $n = 0, 1, 2, \dots$ we get a set of C_n consisting of 2^n closed intervals each having length $1/3^n$. Finally, we define the Cantor set C to be the intersection

$$C = \bigcap_{n=0}^{\infty} C_n$$



Defining the Cantor set

It may be useful to understand C as the remainder of the interval $[0, 1]$ after the iterative process of removing open middle one thirds is taken to infinity.

$$C = [0, 1] - \left[\left(\frac{1}{3}, \frac{2}{3} \right) \cup \left(\frac{1}{9}, \frac{2}{9} \right) \cup \left(\frac{7}{9}, \frac{8}{9} \right) \cup \dots \right]$$

There is some initial doubt whether anything remains at all, but notice that because we are always removing open middle one thirds, then for every $n \in \mathbf{N}$, $0 \in C_n$ and hence $0 \in C$. The same argument shows that $1 \in C$. In fact, if y is the endpoint of some closed interval of some particular set C_n , then it is also an endpoint of one of the intervals of C_{n+1} . Because at each stage, the endpoints are never removed, it follows that $y \in C_n$ for all n . Thus, C at least contains the endpoints of all of the intervals that make up each of the sets C_n .

Is there anything else? Is C countable? Does C contain any intervals? Any irrational numbers? These are difficult questions at the moment. All of the endpoints mentioned earlier are rational numbers (they have the form $m/3^n$), which means that if it is true that C consists of only these endpoints, then C would be a subset of \mathbf{Q} and hence countable. We shall see about this. There is some strong evidence that not much is left in C if we consider the total length of the intervals

removed. To form C_1 , an open interval of length $1/3$ was taken out. In the second step, we removed two intervals of length $1/9$ and to construct C_n , we removed 2^{n-1} middle thirds of length $1/3^n$. There is some logic, then to defining the length of C to be 1 minus the total

$$\frac{1}{3} + 2\left(\frac{1}{9}\right) + 4\left(\frac{1}{27}\right) + \dots + 2^{n-1}\left(\frac{1}{3^n}\right) + \dots = \frac{\frac{1}{3}}{1 - \frac{2}{3}} = 1$$

The Cantor set has *zero length*.

To this point, the information we have collected suggests a mental picture of C as relatively small. For these reasons, the set C is often referred to as Cantor dust. But, there are some strong counterarguments that imply a very different picture. First C is actually *uncountable*, with cardinality equal to the cardinality of \mathbf{R} . One slightly intuitive but convincing way to see this is to create a 1 – 1 correspondence between C and the sequences of the form $(a_n)_{n=1}^{\infty}$, where $a_n = 0$ or 1. For each $c \in C$, set $a_1 = 0$ if c falls in the left-hand component and set $a_1 = 1$ if c falls in the right hand component. Having established where in C_1 , the point c is located, there are now two possible components of C_2 that might contain c . This time, we set $a_2 = 0$ or 1 depending on whether c falls in the left of right half of these two components of C_2 . Continuing in this way, we come to see that every element $c \in C$ yields a sequence (a_1, a_2, a_3, \dots) of zeroes and ones (a binary string), that acts as a set of directions for how to locate c within C . Because the set of sequences of zeroes and ones is uncountable, we must conclude that C is uncountable as well.

What does this imply? In the first place, because the end points of the approximating sets C_n form a countable set, we are forced to accept the fact that not only are there other points in C , but there are uncountably many of them, From the point of view of cardinality, C is quite large - as large as \mathbf{R} , in fact. This should be contrasted with the fact that from the point of view of length, C measures the same size as a single point. We conclude this discussion with a demonstration that from the point of view of *dimension*, C strangely falls somewhere in between.

There is a sensible agreement that a point has dimension zero, a line segment has dimension one, a square has dimension two, and a cube has dimension three. Without attempting a formal definition of dimension (of which there are several) we can nevertheless get a sense of how one might be defined by observing how the dimension affects the result of magnifying each particular set by a factor of 3. (The reason for the choice of 3 will become clear when we turn our attention back to the Cantor set). A single point undergoes no change at all, whereas a line segment triples in length. For the square, magnifying each length by a factor of 3 results in a larger square that contains 9 copies of the original square. Finally, the magnified cube yields a cube that contains 27 copies of the original cube within its volume. Notice that, in each case, to compute the size of the

new set, the dimension appears as the exponent of the magnification factor.

	dim	$\times 3$	new copies
point	0	\rightarrow	$1 = 3^0$
segment	1	\rightarrow	$3 = 3^1$
square	2	\rightarrow	$9 = 3^2$
cube	3	\rightarrow	$27 = 3^3$

Dimension of C ,

Now, apply this transformation to the Cantor set. The set $C_0 = [0, 1]$ becomes the interval $[0, 3]$. Deleting the middle one-third leaves $[0, 1] \cup [2, 3]$, which is where we started in the original construction except that we now stand to produce an additional copy of C in the interval $[2, 3]$. Magnifying the Cantor set by a factor of 3 yields two copies of the original set. Thus, if x is the dimension of C , we must have $3^x = 2$, or $x = \ln 2 / \ln 3 = 0.631$.

The notion of a non-integer or fractional dimension is the impetus behind the term fractal, coined in 1975 by Benoit Mandelbrot to describe a class of sets whose intricate structures have much in common with the Cantor set. Cantor's construction is over hundred years old and for us represents an invaluable testing ground for upcoming theorems and conjectures about the often elusive nature of subsets of the real line.

3.2 Open and Closed Sets.

Given $a \in \mathbf{R}$ and $\epsilon > 0$, recall that the ϵ -neighbourhood of a is the set

$$V_\epsilon(a) = \{x \in \mathbf{R} : |x - a| < \epsilon\}$$

In other words, $V_\epsilon(a)$ is the open interval $(a - \epsilon, a + \epsilon)$ centered at a with radius ϵ .

Definition. A set $O \subseteq \mathbf{R}$ is open if for all points $a \in O$ there exists an ϵ -neighbourhood $V_\epsilon(a) \subseteq O$.

Example.

(i) Perhaps, the simplest example of an open set is \mathbf{R} itself. Given an arbitrary element $a \in \mathbf{R}$, we are free to pick any ϵ -neighbourhood we like and it will always be true that $V_\epsilon(a) \subseteq \mathbf{R}$. It is also the case that the logical structure of the definition requires us to classify the empty set \emptyset as an open subset of the real line.

(ii) For a more useful collection of examples, consider the open interval

$$(c, d) = \{x \in \mathbf{R} : c < x < d\}$$

To see that (c, d) is open in the same sense just defined, let $x \in (c, d)$ be arbitrary. If we take $\epsilon = \min\{x - c, d - x\}$, then it follows that $V_\epsilon(x) \subseteq (c, d)$. It is important to see where this argument breaks down if the interval includes either one of its endpoints.

The union of open intervals is another example of an open set. This observation leads to the next result.

Theorem. (i) The union of an arbitrary collection of open sets is open.
(ii) The intersection of a countably finite collection of open sets is open.

Proof.

To prove (i), we let $\{O_\lambda : \lambda \in \Lambda\}$ be a collection of open sets and $O = \bigcup_{\lambda \in \Lambda} O_\lambda$. Here, Λ could be a countably infinite or an uncountable set. Let a be an arbitrary element of O . In order to show that O is open, the definition insists that we produce an ϵ -neighbourhood of a completely contained in O . But, $a \in O$ implies that a is an element of at least one particular $O_{\lambda'}$. Because we are assuming that $O_{\lambda'}$ is open, by definition we can assert, that there exists $V_\epsilon(a) \subseteq O_{\lambda'}$. The fact that $O_{\lambda'} \subseteq O$ allows us to conclude that $V_\epsilon(a) \subseteq O$ for all $a \in O$. This completes the proof of (i).

For (ii), let $\{O_1, O_2, \dots, O_N\}$ be a finite collection of open sets. Now, if $a \in \bigcap_{k=1}^N O_k$, then a is an element of each of the open sets. By the definition of an open set, we know that, for each, $1 \leq k \leq N$, there exists a $V_{\epsilon_k}(a) \subseteq O_k$. We are in search of a single neighbourhood of a that is contained in every O_k , so the trick is to take the smallest one. Letting

$\epsilon = \min\{\epsilon_1, \epsilon_2, \dots, \epsilon_N\}$, it follows that $V_\epsilon(a) \subseteq V_{\epsilon_k}(a)$ for all k , and hence $V_\epsilon(a) \subseteq \bigcap_{k=1}^N O_k$ as desired.

Closed Sets.

Definition. A point x is the **limit point** of a set A , if every ϵ -neighbourhood $V_\epsilon(x)$ intersects the set A at some point other than x .

Limit points are also referred to as cluster points or accumulation points, but the phrase, " x is a limit point of A " has the advantage of explicitly reminding us that x is quite literally the limit of the sequence in A .

Theorem. A point x is a limit point of a set A if and only if $x = \lim a_n$ for some sequence (a_n) contained in A , satisfying $a_n \neq x$ for all $n \in \mathbf{N}$.

Proof.

(\implies) Assume that x is a limit point of A . In order to produce a sequence (a_n) converging to x , we are going to consider the particular ϵ -neighbourhoods obtained using $\epsilon = 1/n$. By definition, every neighborhood of x intersects A in some point other than x . This means that, for each $n \in \mathbf{N}$, we are justified in picking a point

$$a_n \in V_{1/n}(x) \cap A$$

with the stipulation that $a_n \neq x$. It should not be too difficult to see why $(a_n) \rightarrow x$. Given an arbitrary $\epsilon > 0$, choose N such that $1/N < \epsilon$. Then for all $n \geq N$, we have $|a_n - x| < \epsilon$.

(\impliedby) For the reverse implication, we assume that $\lim a_n = x$ where $a_n \in A$ but $a_n \neq x$ and let $V_\epsilon(x)$ be an arbitrary neighbourhood of x . The definition of convergence assures us that there exists a term a_N in the sequence satisfying $a_N \in V_\epsilon(x)$. So, every ϵ -neighbourhood, $V_\epsilon(x)$ intersects A in some element other than x . Hence, $x = \lim a_n$. QED.

The restriction that $a_n \neq x$ in the above theorem deserves a comment. Given a point $a \in A$, it is always the case that a is the limit of a sequence in A , if we are allowed to consider the constant sequence (a, a, a, \dots) . There will be occasions where we will want to avoid this somewhat uninteresting situation, so it is important to have vocabulary that can distinguish limit points of a set from isolated points.

Definition. A point $a \in A$ is an isolated point of A if it is not a limit point of A .

As a word of caution, we need to be a little careful about how we understand the relationship between these concepts. Whereas an isolated point is always an element of the relevant set A , it is quite possible for a limit point of A not to belong to A . As an example, consider the endpoint of an open interval. This situation is the subject of the next important definition.

Definition. A set $F \subseteq \mathbf{R}$ is closed if it contains its limit points.

The adjective "closed" appears in several other mathematical contexts and is usually employed to

mean that an operation on the elements of a given set does not take us outside of the set. In linear algebra, for example, a vector space is a set that is closed under vector addition and scalar multiplication. In analysis, the operation we are concerned with is the limiting operation. Topologically speaking, a closed set is the one where convergent sequences within the set have limits that are also in the set.

Theorem. A set $F \subseteq \mathbf{R}$ is closed if and only if every Cauchy sequence contained in F has a limit that is also an element of F .

Proof. Exercise 3.2.5.

Example. (i) Consider

$$A = \left\{ \frac{1}{n} : n \in \mathbf{N} \right\}$$

Let's show that each point of A is isolated. Given $1/n \in A$, choose $\epsilon = \frac{1}{n} - \frac{1}{n+1}$. (Note that, $\frac{1}{n-1} - \frac{1}{n}$ is larger than $\frac{1}{n} - \frac{1}{n+1}$). Then,

$$V_\epsilon(1/n) \cap A = \frac{1}{n}$$

It follows from the definition that $1/n$ is not a limit point and so is isolated. Although all of the points of A are isolated, the set does have one limit point, namely 0. This is because every ϵ -neighbourhood of centered at zero, no matter how small, is going to contain points of A . Because $0 \notin A$, A is not closed. The set $F = A \cup \{0\}$ is an example of a closed set and is called the closure of A . (The closure of a set is discussed in a moment).

(ii) Let's prove that a closed interval

$$[c, d] = \{x \in \mathbf{R} : c \leq x \leq d\}$$

is a closed set using the definition. If x is a limit point of $[c, d]$, then by the previous theorem there exists a sequence (x_n) contained in $[c, d]$ such that $(x_n) \rightarrow x$ and $x_n \neq x$. We need to prove that x belongs to $[c, d]$.

The key to this argument is contained in the Order Limit Theorem, which summarizes the

relationship between inequalities and the corresponding limiting process. Because, $c \leq x_n \leq d$, it follows from the Order Limit Theorem, that $c \leq x \leq d$. Thus, $[c, d]$ is closed.

(iii) Consider the set $\mathbf{Q} \subseteq \mathbf{R}$ of rational numbers. An extremely important property of \mathbf{Q} is that the set of all limit points of \mathbf{Q} is actually all of \mathbf{R} . To see why this is so, recall the theorem, which is referred to as the density property of \mathbf{Q} in \mathbf{R} . \mathbf{Q} is dense in \mathbf{R} implies that \mathbf{Q} sits inside of \mathbf{R} . Between any two real numbers $a, b \in \mathbf{R}$, you can always find a rational number r satisfying, $a < r < b$.

Let $y \in \mathbf{R}$ be arbitrary and consider any neighbourhood $(y - \epsilon, y + \epsilon)$. The density theorem allows us to conclude that there exists a rational number $r \neq y$ that falls in this neighbourhood. Thus, y is a limit point of \mathbf{Q} .

The density property of \mathbf{Q} can now be reformulated in the following way.

Theorem.(Density of \mathbf{Q} in \mathbf{R}). For every $y \in \mathbf{R}$, there exists a sequence of the rational numbers that converges to y .

Proof.

From the above discussion, we know that for an arbitrary $y \in \mathbf{R}$, every ϵ -neighbourhood of y intersects \mathbf{Q} in some point other than y . y is a limit point of \mathbf{Q} . By the theorem on limit points of sets, there exists a sequence $(x_n) \subseteq \mathbf{Q}$, such that $(x_n) \rightarrow y$, and $x_n \neq y$ for all n .

The same argument can also be used to show that every real number is the limit of a sequence of irrational numbers. Although interesting, part of the allure of the rational numbers is that, in addition to being dense in \mathbf{R} , they are countable. As we will see, this tangible aspect of \mathbf{Q} makes it an extremely useful set, both for proving theorems and for producing interesting counterexamples.

Closure.

Definition. Given a set $A \subseteq \mathbf{R}$, let L be the set of all limit points of A . The closure A is defined to $cl(A) = A \cup L$.

In example 3.2.9 (i), we saw that if $A = \{1/n : n \in \mathbf{N}\}$, then the closure of $A = cl(A) = A \cup \{0\}$. Example 3.2.9 (iii) verifies that $cl(\mathbf{Q}) = \mathbf{R}$. If A is an open interval, then $cl(A) = [a, b]$. If A is a closed interval, then $cl(A) = A$. It is not for lack of imagination that in each of these examples $cl(A)$ is always a closed set.

Theorem. For any $A \subseteq \mathbf{R}$, the closure $cl(A)$ is a closed set and is the smallest closed set

containing A .

Proof.

If L is the set of limit points of A , then it is immediately clear that $cl(A)$ contains the limit points of A . There is still something more to prove, however because taking the union of L with A could potentially produce some new limit points of $cl(A)$. In exercise 3.2.7, we outline the argument that this does not happen.

Now, any closed set containing A must contain L as well. This shows that $cl(A) = A \cup L$ is the smallest such closed set containing A .

Complements.

The mathematical notions of open and closed are not antonyms the way they are in standard English. If a set is not open, that does not imply it must be closed. Many sets such as the half-open interval $(c, d] = \{x \in \mathbf{R} : c < x \leq d\}$ are neither open nor closed. The sets \mathbf{R} and \emptyset are simultaneously open and closed, although, thankfully, these are the only ones with this disorientating property. (Exercises 3.2.13). There is, however, an important relationship between open and closed sets. Recall that the complement of a set $A \subseteq \mathbf{R}$ is defined to be the set:

$$A^C = \{x \in \mathbf{R} : x \notin A\}$$

Theorem. A set O is open if and only if O^C is closed. Likewise, a set F is closed if and only if F^C is open.

Proof.

(\implies) Given an open set $O \subseteq \mathbf{R}$, let's first prove that O^C is a closed set. To prove that O^C is a closed, we need to show that it contains all of its limit points. If x is a limit point of O^C , then every ϵ -neighbourhood of x contains some point of O^C other than x . But that is enough to conclude that x cannot be in O , because if $x \in O$, then every open interval $(x - \epsilon, x + \epsilon)$ is contained in O . Consequently, $x \in O^C$. Thus, O^C is a closed set.

(\impliedby) For the converse statement, we assume that O^C is closed and argue that O is open. Thus, given an arbitrary point $x \in O$, we must produce an ϵ -neighbourhood $V_\epsilon(x) \subseteq O$. Because, $x \notin O^C$, we can be sure that x is not a limit point of O^C . If x is not a limit point of O^C , there exists an $\epsilon > 0$, such that $(x - \epsilon, x + \epsilon) \not\subseteq O^C$, or equivalently $(x - \epsilon, x + \epsilon) \subseteq O$. This is precisely what we needed to show.

Theorem. (i) The union of a countably finite collection of closed sets is closed.

(ii) The intersection of an arbitrary collection of closed sets is closed.

Proof.

De Morgan's Laws state that for any collection of sets $\{E_\lambda : \lambda \in \Lambda\}$ it is true that

$$\left(\bigcup_{\lambda \in \Lambda} E_\lambda \right)^c = \bigcap_{\lambda \in \Lambda} E_\lambda^c \quad \text{and} \quad \left(\bigcap_{\lambda \in \Lambda} E_\lambda \right)^c = \bigcup_{\lambda \in \Lambda} E_\lambda^c$$

The result follows directly from these statements and Theorem 3.2.3. The details are requested in Exercise 3.2.9.

Exercises.

1. [Abbot, 3.2.1] (a) Where in the proof of Theorem 3.2.3 part (ii) does the assumption that the collection of open sets be *finite* get used?

Solution. a is contained in each of the open sets O_1, O_2, \dots, O_N . Since, there are open sets, there exists an ϵ_k -neighbour of a , such that $V_{\epsilon_k}(x) \subseteq O_k, k = 1, \dots, N$.

We would like to select an ϵ -neighbourhood that is contained in $\bigcap_{k=1}^n O_k$. To do so, we choose ϵ to be the smallest distance in the collection $\{\epsilon_1, \dots, \epsilon_N\}$. This is where the assumption that the collection of open sets be finite gets used.

For if, the collection is countably infinite or uncountable, we risk $\inf\{\epsilon_1, \epsilon_2, \dots\} = 0$. We simply might not be able to find an ϵ -neighbourhood contained in every open set O_k .

(b) Give an example of a countable collection of open sets $\{O_1, O_2, O_3, \dots\}$ whose intersection $\bigcap_{n=1}^{\infty} O_n$ is closed, not empty and not all of \mathbf{R} .

Proof.

Consider $O_n = \left(-1 - \frac{1}{n}, 1 + \frac{1}{n}\right)$. O_n is an open set. $\bigcap_{n=1}^{\infty} O_n = [-1, 1]$ and this is a closed set.

2. [Abbott, 3.2.2] Let

$$A = \left\{ (-1)^n + \frac{2}{n} : n = 1, 2, 3, \dots \right\} \quad \text{and} \quad B = \{x \in \mathbf{Q} : 0 < x < 1\}$$

Answer the following questions for each set.

(a) What are the limit points?

The Set A :

Enlisting the first few elements of A :

$$A = \left\{ 1, 2, -\frac{1}{3}, \frac{3}{2}, -\frac{3}{5}, \frac{4}{3}, \dots \right\}$$

The subsequence (a_{2n}) converges to 1, whilst the subsequence (a_{2n-1}) converges to (-1) . Hence, the limit points of A are $L = \{1, -1\}$.

The Set B :

Note that \mathbf{Q} is dense in \mathbf{R} . Consider an arbitrary real number $y \in [0, 1]$. Pick $\epsilon = \min\{y - 0, 1 - y\}$. Every open interval $(y - \epsilon, y + \epsilon)$ intersects B in some point other than y , because we can find a rational x such that $y - \epsilon < x < y$. Consequently, y is a limit point of B .

So, the limit points of B is the closed interval $[0, 1]$.

(b) Is the set open? Closed?

The Set A :

Pick an arbitrary point $x \in A$, for example $x = \frac{3}{2}$. Let $\epsilon > 0$. Clearly, for all $\epsilon > 0$,

$\left(\frac{3}{2} - \epsilon, \frac{3}{2} + \epsilon\right)$ is not contained in A . Thus, there exists an $x \in A$, such that for all $\epsilon > 0$, $(x - \epsilon, x + \epsilon)$ is not in A . So, A is not open.

The limit point -1 does not belong to A . So, A is not closed.

The Set B :

Between any two rational numbers, there is an irrational. For example, consider $a, b \in B$, where $a < b$. Then,

$$a < a + \frac{b-a}{\sqrt{2}} < b$$

So, every open interval $(b - \epsilon, b + \epsilon)$ will contain irrational points, and is therefore not contained in B .

We have shown that $\forall \epsilon > 0$, we have $(b - \epsilon, b + \epsilon) \not\subseteq B$ for atleast one $b \in B$. Consequently, B is not open.

The set B does not contain its limit points. So, B is not closed.

(c) Does the set contain any isolated points?

The Set A :

Consider the distance $\left| \left((-1)^n + \frac{2}{n} \right) - \left((-1)^{n+2} + \frac{2}{n+2} \right) \right| = \frac{4}{n(n+2)}$. Since,

$\frac{4}{(n+2)^2} < \frac{4}{n(n+2)}$, pick $\epsilon = \frac{4}{(n+2)^2}$. For this choice of ϵ , the open interval $(a_n - \epsilon, a_n + \epsilon) \cap A = a_n$, for $n > 1$. Hence, all of the points except 1, are isolated points of A .

The Set B :

Let $x = \frac{p}{q}$ be any rational such that $0 < x < 1$. The sequence $x_n = \frac{p}{q} + \frac{1}{n}$ converges to $\frac{p}{q}$. So every rational number $x \in B$ is a limit point. B has no isolated points.

(d) Find the closure of the set.

The set A :

$$cl(A) = A \cup \{-1\}$$

The set B :

$$cl(B) = [0, 1]$$

3. [Abbott, 3.2.3] Decide whether the following sets are open, closed or neither. If a set is not open, find a point in the set for which there is no ϵ -neighbourhood contained in the set. If a set is not closed, find a limit point that is not contained in the set.

(a) \mathbb{Q} .

Given a rational number $q \in \mathbf{Q}$, there exists an irrational number $x \in \mathbf{I}$, such that $|x - q| < \epsilon$ for all $\epsilon > 0$. For example, if $q = \frac{m}{n}$ and $\epsilon = \frac{1}{n}$, then $\frac{m}{n} < \frac{m+1}{\sqrt{2}n} < \frac{m}{n} + \frac{1}{n}$. So, $(x - \epsilon, x + \epsilon)$ is not contained in \mathbf{Q} . Therefore, \mathbf{Q} is not open.

Also, let $y \in \mathbf{R}$. Every ϵ -neighbourhood of y intersects \mathbf{Q} , since \mathbf{Q} is dense in \mathbf{R} . So, we can always find a rational q such that $y - \epsilon < q < y$. Consequently, the limit points of \mathbf{Q} is all of \mathbf{R} . So, \mathbf{Q} is not closed.

(b) \mathbf{N} .

For all $\epsilon > 0$, $(n - \epsilon, n + \epsilon)$ is not contained in \mathbf{N} . To see this, consider $n + \epsilon/2 \in (n - \epsilon, n + \epsilon)$. Clearly, $(n + \epsilon/2) \notin \mathbf{N}$. So, \mathbf{N} is not open.

\mathbf{N} is unbounded and has no limit points. Therefore, \mathbf{N} is closed.

(c) $\{x \in \mathbf{R} : x \neq 0\}$

Let $x \in \mathbf{R} - \{0\}$. There exists an ϵ -neighbourhood $V_\epsilon(x) = (x - \epsilon, x + \epsilon)$, of x , that is contained entirely in the set $\{x \in \mathbf{R} : x \neq 0\}$ for all x . Hence, it is an open set.

Consider $x_n = \frac{1}{n}$. Clearly, $x_n \in \{x \in \mathbf{R} : x \neq 0\}$ and $\lim(x_n) = 0$. Thus, 0 is a limit point of the given set. Since, 0 is not in the given set, it is not a closed set.

(d) $\left\{1 + \frac{1}{4} + \frac{1}{9} + \dots + \frac{1}{n^2} : n \in \mathbf{N}\right\}$.

Let (s_n) be the partial sums of the infinite series $\sum_{n=1}^{\infty} \frac{1}{n^2}$. So, the elements of the set are:

$$\left\{s_n : s_n = \sum_{k=1}^n \frac{1}{k^2}\right\}$$

For all $\epsilon > 0$, $(s_n - \epsilon, s_n + \epsilon)$ is not contained in the given set. So, it is not an open set.

Moreover, (s_n) is a convergent sequence. So, $(s_n) \rightarrow s$ with $s_n \neq s$ for all n . Thus, the limit

point s is not an element of the given set. Consequently, $\left\{s_n : s_n = \sum_{k=1}^n \frac{1}{k^2}\right\}$ is not closed.

(e) $\{1 + 1/2 + 1/3 + \dots + 1/n : n \in \mathbf{N}\}$.

For all $\epsilon > 0$, open interval $(s_n - \epsilon, s_n + \epsilon)$ is not contained in the given set. So, it is not an open set. Further, the harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ is divergent, so the set has no limit points. Hence, it is closed.

4. [Abbott, 3.2.4] Let A be non-empty and bounded above so that $s = \sup A$ exists.

(a) Show that $s \in cl(A)$.

By the definition of the supremum, for all $\epsilon > 0$, there exists an $a_n \in A$, such that $s - \epsilon < a_n < s$. If we successively choose, $\epsilon = 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots, \frac{1}{n}$, we produce a sequence $(a_n) \subseteq A$ such that $\lim(a_n) = s$ and $a_n \neq s$. Thus, s is the limit point of A . Consequently, $s \in cl(A)$.

(b) Can an open set contain its supremum?

Let O be an open set and suppose $s \in O$. Then, by definition of open sets, there exists an $\epsilon > 0$, such that the open interval $(s - \epsilon, s + \epsilon) \subseteq O$. So, $s + \epsilon/2 \in O$. But this implies, s is not the supremum of O . This is a contradiction. Hence, our initial assumption is false, and $s \notin O$.

5. [Abbott, 3.2.5] Prove theorem 3.2.8

Theorem. A set $F \subseteq \mathbf{R}$ is closed if and only if every Cauchy sequence contained in F has a limit point that is also an element of F .

p : A set F is closed.

q : **Every** Cauchy sequence contained in F converges to some point in F .

$(p \implies q)$ Assume that the set F is closed. Let (x_n) be an arbitrary Cauchy sequence in F . Cauchy sequences are convergent, so (x_n) is a convergent sequence. Suppose $\lim x_n = x$. By definition of convergence, for all $\epsilon > 0$, there exists an $N \in \mathbf{N}$, such that $x_n \in (x - \epsilon, x + \epsilon)$ for all $n \geq N$.

Thus, for all $\epsilon > 0$, $(x - \epsilon, x + \epsilon)$ intersects F in some point other than x . Consequently, x is a limit point of F . Since, F is closed, $x \in F$.

Thus, every Cauchy sequence contained in F has a limit point in F .

($p \iff q$) Assume that every Cauchy sequence contained in F converges to some point in F . We must prove that F contains all its limit points. Let x be an arbitrary limit point of F .

By definition of a limit point, there exists a sequence $(x_n) \subseteq F$, such that $\lim x_n = x$, with $x_n \neq x$. Convergent sequences are Cauchy. So, (x_n) is a Cauchy sequence. Therefore, $x \in F$. Consequently, F contains all its limit points.

6. [Abbott, 3.2.6] Decide whether the following statements are true or false. Provide counterexamples for those that are false, and supply proofs for those that are true.

(a) An open set that contains every rational number must necessarily be all of \mathbf{R} .

This proposition is false.

Counterexample. Consider the set $\mathbf{R} - \{0\} = \mathbf{R}_+ \cup \mathbf{R}_-$. There exists an ϵ -neighbourhood, $V_\epsilon(x)$, of x , such that $V_\epsilon(x) \subseteq \mathbf{R} - \{0\}$ for all x belonging to the set. So, $\mathbf{R} - \{0\}$ is open, but it does not contain all of the real numbers.

(b) The nested interval property (NIP) remains true if the term **closed interval** is replaced by closed set.

This proposition is false.

Counterexample.

Consider $\bigcap_{n=1}^{\infty} [n, \infty)$.

(c) Every non-empty open set contains a rational number.

This proposition is true.

Proof. Let O be an open set and suppose $x \in O$. By definition, there exists an $\epsilon > 0$, such that $(x - \epsilon, x + \epsilon) \subseteq O$. As \mathbf{Q} is dense in \mathbf{R} , there exists a rational number q such that

$x - \epsilon < q < x$. So, every non-empty open set contains a rational number.

(d) Every bounded infinite closed set contains a rational number.

This proposition is false.

Counterexample.

Consider the set $A = \left\{ \pi + \frac{1}{n} : n \in \mathbf{N} \right\} \cup \{\pi\}$. π is a limit point of the set and it belongs to A .

So, A is a closed set. A is a bounded infinite set. A does not contain rational numbers.

(e) The Cantor set is closed.

C_n is a closed set for all $n \in \mathbf{N}$. The intersection of an arbitrary collection of closed sets is closed, so $\bigcap_{n=1}^{\infty} C_n$ is also a closed set.

7. [Abbott, 3.2.7] Given $A \subseteq \mathbf{R}$, let L be the set of all limit points of A .

(a) Show that the set L is closed.

Let x be a limit point of L . We are interested to prove that $x \in L$.

Since x is a limit point of L , every ϵ -neighbourhood $V_{\epsilon}(x)$, of x intersects L in some point y other than x .

So, for all $\epsilon > 0$, there exists a $y \in L$, such that

$$|y - x| < \frac{\epsilon}{2}$$

But, y is a limit point of A . So, for all $\epsilon > 0$, there exists a $z \in A$, such that

$$|z - y| < \frac{\epsilon}{2}$$

The distance $|z - x|$ can be bounded as follows:

$$\begin{aligned}
|z - x| &= |z - y + y - x| \\
&\leq |z - y| + |y - x| \\
&< \epsilon/2 + \epsilon/2 = \epsilon
\end{aligned}$$

So, every ϵ -neighbourhood of x intersects A in some point z other than x . Consequently, x is a limit point of A . Thus, $x \in L$. So, L is closed.

(b) Argue that if x is a limit point of $A \cup L$, then x is a limit point of A .

Suppose x is a limit point of $A \cup L$. Thus, either x is a limit point of A or x is a limit point of L . If x is a limit point of A , then $x \in L$. If x is a limit point of L , then since L is closed, $x \in L$. Also, $A \cup L$ is the smallest closed set containing A . So, the above two possibilities are exhaustive.

8. [Abbott, 3.2.8] Assume that A is an open set and B is a closed set. Determine if the following sets are definitely open, definitely closed, both or neither.