

HIERARCHICAL MODELING: MIXED EFFECTS VS BAYESIAN

SAGESURE DATA SCIENCE SYMPOSIUM 2022

W. D. BRINDA

QUANTITATIONS LLC

11 NOV 2022



LINEAR MODEL VS LINEAR MIXED MODEL

The general formulation of the multiple linear model with iid Normal errors:

$$Y_i = \beta_0 + \beta_1 x_i^{(1)} + \dots + \beta_d x_i^{(d)} + \epsilon_i$$

where $\epsilon_1, \dots, \epsilon_n$ are iid Normal with mean zero.

LINEAR MODEL VS LINEAR MIXED MODEL

The general formulation of the multiple linear model with iid Normal errors:

$$Y_i = \beta_0 + \beta_1 x_i^{(1)} + \dots + \beta_d x_i^{(d)} + \epsilon_i$$

where $\epsilon_1, \dots, \epsilon_n$ are iid Normal with mean zero.

The coefficients are considered *fixed*; that is, the model makes *absolutely no assumptions* about them.

LINEAR MODEL VS LINEAR MIXED MODEL

The general formulation of the multiple linear model with iid Normal errors:

$$Y_i = \beta_0 + \beta_1 x_i^{(1)} + \dots + \beta_d x_i^{(d)} + \epsilon_i$$

where $\epsilon_1, \dots, \epsilon_n$ are iid Normal with mean zero.

The coefficients are considered *fixed*; that is, the model makes *absolutely no assumptions* about them.

In contrast, the general formulation of the linear *mixed* model with iid Normal errors:

$$Y_i = \beta_0 + \beta_1 x_i^{(1)} + \dots + \beta_d x_i^{(d)} + \underbrace{b_1 z_i^{(1)} + \dots + b_m z_i^{(m)}} + \epsilon_i$$

with $\epsilon_1, \dots, \epsilon_n$ iid Normal and the “*random effects*” b_1, \dots, b_m are assumed to be drawn from a multivariate Normal distribution.

EXAMPLE: BLOOD PRESSURE MEASUREMENTS

Suppose that n_f females and n_m males each had their [systolic] blood pressures measured. Assume that the distributions of female and male blood pressures are both Normal with the same standard deviation σ_w . Let their expectations be denoted μ_f and μ_m respectively. With x_1, \dots, x_n representing sex (the only explanatory variable) and Y_1, \dots, Y_N representing blood pressures, we can represent this scenario with the linear model

$$Y_i = \mu_f + (\mu_m - \mu_f)\mathbb{I}(x_i = \text{"M"}) + \epsilon_i.$$

EXAMPLE: BLOOD PRESSURE MEASUREMENTS

Suppose that n_f females and n_m males each had their [systolic] blood pressures measured. Assume that the distributions of female and male blood pressures are both Normal with the same standard deviation σ_w . Let their expectations be denoted μ_f and μ_m respectively. With x_1, \dots, x_n representing sex (the only explanatory variable) and Y_1, \dots, Y_N representing blood pressures, we can represent this scenario with the linear model

$$Y_i = \mu_f + (\mu_m - \mu_f)\mathbb{I}(x_i = \text{"M"}) + \epsilon_i.$$

Now suppose that each of the subjects' blood pressure was measured up to four times, over the course of a year. The measurements of a single person's blood pressure can vary quite a bit when measured at different times. How could we adapt our previous blood pressure model to capture this?

EXAMPLE: BLOOD PRESSURE MEASUREMENTS

Suppose that n_f females and n_m males each had their [systolic] blood pressures measured. Assume that the distributions of female and male blood pressures are both Normal with the same standard deviation σ_w . Let their expectations be denoted μ_f and μ_m respectively. With x_1, \dots, x_n representing sex (the only explanatory variable) and Y_1, \dots, Y_N representing blood pressures, we can represent this scenario with the linear model

$$Y_i = \mu_f + (\mu_m - \mu_f)\mathbb{I}(x_i = \text{"M"}) + \epsilon_i.$$

Now suppose that each of the subjects' blood pressure was measured up to four times, over the course of a year. The measurements of a single person's blood pressure can vary quite a bit when measured at different times. How could we adapt our previous blood pressure model to capture this?

Define $z_i^{(j)}$ to be the indicator variable for whether observation i came from subject j . Then, we might want to modify our model to

$$Y_i = \mu_f + (\mu_m - \mu_f)\mathbb{I}(x_i = \text{"M"}) + b_1 z_i^{(1)} + \dots + b_m z_i^{(m)} + \epsilon_i.$$

with iid Normal *random effects* b_1, \dots, b_m .

EXAMPLE: BLOOD PRESSURE MEASUREMENTS

Suppose that n_f females and n_m males each had their [systolic] blood pressures measured. Assume that the distributions of female and male blood pressures are both Normal with the same standard deviation σ_w . Let their expectations be denoted μ_f and μ_m respectively. With x_1, \dots, x_n representing sex (the only explanatory variable) and Y_1, \dots, Y_N representing blood pressures, we can represent this scenario with the linear model

$$Y_i = \mu_f + (\mu_m - \mu_f)\mathbb{I}(x_i = \text{"M"}) + \epsilon_i.$$

Now suppose that each of the subjects' blood pressure was measured up to four times, over the course of a year. The measurements of a single person's blood pressure can vary quite a bit when measured at different times. How could we adapt our previous blood pressure model to capture this?

Define $z_i^{(j)}$ to be the indicator variable for whether observation i came from subject j . Then, we might want to modify our model to


$$Y_i = \mu_f + (\mu_m - \mu_f)\mathbb{I}(x_i = \text{"M"}) + b_1 z_i^{(1)} + \dots + b_m z_i^{(m)} + \epsilon_i.$$

with iid Normal *random effects* b_1, \dots, b_m .

To better understand what this model means, suppose observation 12 came from subject 5 who is a female: the corresponding equation simplifies to

$$Y_{12} = \mu_f + b_5 + \epsilon_{12}$$

MODEL DIAGRAM



Now, I'll turn to an R script to simulate blood pressure data according to the mechanism just described and draw some plots to visualize it. (We'll continue through this script at later points in the talk.)

If you'd like, you can download the file and execute the code along with me:

quantitations.com/static/symposium.r

HEALTH INSURANCE DECISION EXAMPLE

Context: Health Decision Corporation (HDC) provides health insurance to 25,850 members (15,125 females, 10,725 males).

HEALTH INSURANCE DECISION EXAMPLE

Context: Health Decision Corporation (HDC) provides health insurance to 25,850 members (15,125 females, 10,725 males).

Scenario: Suppose that HDC is considering a policy of measuring each member's blood pressure four times per year. Any time a measurement is at least 140, the member will undergo additional tests (at a cost of \$200 to HDC). Furthermore, if the measurement is at least 150, the member will also be provided with medication (at a cost of \$300 to HDC).

HEALTH INSURANCE DECISION EXAMPLE

Context: Health Decision Corporation (HDC) provides health insurance to 25,850 members (15,125 females, 10,725 males).

Scenario: Suppose that HDC is considering a policy of measuring each member's blood pressure four times per year. Any time a measurement is at least 140, the member will undergo additional tests (at a cost of \$200 to HDC). Furthermore, if the measurement is at least 150, the member will also be provided with medication (at a cost of \$300 to HDC).

Decision Problem: HDC wants to understand how much they should expect to spend on follow-up tests and medications next year if they implement this idea.

HEALTH INSURANCE DECISION EXAMPLE

Context: Health Decision Corporation (HDC) provides health insurance to 25,850 members (15,125 females, 10,725 males).

Scenario: Suppose that HDC is considering a policy of measuring each member's blood pressure four times per year. Any time a measurement is at least 140, the member will undergo additional tests (at a cost of \$200 to HDC). Furthermore, if the measurement is at least 150, the member will also be provided with medication (at a cost of \$300 to HDC).

Decision Problem: HDC wants to understand how much they should expect to spend on follow-up tests and medications next year if they implement this idea.

Data: HDC selected 11 members at random this year (5 female, 6 male) to have their blood pressures measured, between 1 and 4 times each.

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males.

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates.

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Frequentist: If males don't truly have higher blood pressure, then the probability that the data would be at least this extreme is at most .14.

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Frequentist: If males don't truly have higher blood pressure, then the probability that the data would be at least this extreme is at most .14.

Manager: ...

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Frequentist: If males don't truly have higher blood pressure, then the probability that the data would be at least this extreme is at most .14.

Manager: ... What? ...

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Frequentist: If males don't truly have higher blood pressure, then the probability that the data would be at least this extreme is at most .14.

Manager: ... What? ... Alright well, at least tell me how likely it is that this scheme will cost us more than \$600,000 dollars?

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Frequentist: If males don't truly have higher blood pressure, then the probability that the data would be at least this extreme is at most .14.

Manager: ... What? ... Alright well, at least tell me how likely it is that this scheme will cost us more than \$600,000 dollars?

Frequentist: If my parameter estimates are correct, the probability would be about .028.

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Frequentist: If males don't truly have higher blood pressure, then the probability that the data would be at least this extreme is at most .14.

Manager: ... What? ... Alright well, at least tell me how likely it is that this scheme will cost us more than \$600,000 dollars?

Frequentist: If my parameter estimates are correct, the probability would be about .028.

Manager: And do you think the parameter estimates *are* correct?

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Frequentist: If males don't truly have higher blood pressure, then the probability that the data would be at least this extreme is at most .14.

Manager: ... What? ... Alright well, at least tell me how likely it is that this scheme will cost us more than \$600,000 dollars?

Frequentist: If my parameter estimates are correct, the probability would be about .028.

Manager: And do you think the parameter estimates *are* correct?

Frequentist: No.

FREQUENTIST'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Frequentist: If males don't truly have higher blood pressure, then the probability that the data would be at least this extreme is at most .14.

Manager: ... What? ... Alright well, at least tell me how likely it is that this scheme will cost us more than \$600,000 dollars?

Frequentist: If my parameter estimates are correct, the probability would be about .028.

Manager: And do you think the parameter estimates *are* correct?

Frequentist: No.

[Uncomfortable silence...]

CLASSIC BAYES THEOREM EXAMPLE

Classic counter-intuitive example demonstrating Bayes' Theorem: Suppose the prevalence of a disease in the general population is 1 in 5000. A test for this disease has a 1% false positive rate and a 2% false negative rate. If this test gives you a positive result, find the probability that you have the disease.

CLASSIC BAYES THEOREM EXAMPLE

Classic counter-intuitive example demonstrating Bayes' Theorem: Suppose the prevalence of a disease in the general population is 1 in 5000. A test for this disease has a 1% false positive rate and a 2% false negative rate. If this test gives you a positive result, find the probability that you have the disease.

$$\begin{aligned}\mathbb{P}(\text{disease}|\text{positive}) &= \frac{\mathbb{P}(\text{disease})\mathbb{P}(\text{positive}|\text{disease})}{\mathbb{P}(\text{positive})} \\ &= \frac{(1/5000)(.98)}{(1/5000)(.98) + (4999/5000)(.01)} \\ &\approx 1/51\end{aligned}$$

CLASSIC BAYES THEOREM EXAMPLE

Classic counter-intuitive example demonstrating Bayes' Theorem: Suppose the prevalence of a disease in the general population is 1 in 5000. A test for this disease has a 1% false positive rate and a 2% false negative rate. If this test gives you a positive result, find the probability that you have the disease.

$$\begin{aligned}\mathbb{P}(\text{disease}|\text{positive}) &= \frac{\mathbb{P}(\text{disease})\mathbb{P}(\text{positive}|\text{disease})}{\mathbb{P}(\text{positive})} \\ &= \frac{(1/5000)(.98)}{(1/5000)(.98) + (4999/5000)(.01)} \\ &\approx 1/51\end{aligned}$$

It's not often highlighted, but this example also demonstrates the problem with interpreting p-values to decide whether a hypothesis is likely true or false:

CLASSIC BAYES THEOREM EXAMPLE

Classic counter-intuitive example demonstrating Bayes' Theorem: Suppose the prevalence of a disease in the general population is 1 in 5000. A test for this disease has a 1% false positive rate and a 2% false negative rate. If this test gives you a positive result, find the probability that you have the disease.

$$\begin{aligned}\mathbb{P}(\text{disease}|\text{positive}) &= \frac{\mathbb{P}(\text{disease})\mathbb{P}(\text{positive}|\text{disease})}{\mathbb{P}(\text{positive})} \\ &= \frac{(1/5000)(.98)}{(1/5000)(.98) + (4999/5000)(.01)} \\ &\approx 1/51\end{aligned}$$

It's not often highlighted, but this example also demonstrates the problem with interpreting p-values to decide whether a hypothesis is likely true or false: If a patient tests positive for the disease, the p-value for the null hypothesis (no disease) is .01.

CLASSIC BAYES THEOREM EXAMPLE

Classic counter-intuitive example demonstrating Bayes' Theorem: Suppose the prevalence of a disease in the general population is 1 in 5000. A test for this disease has a 1% false positive rate and a 2% false negative rate. If this test gives you a positive result, find the probability that you have the disease.

$$\begin{aligned}\mathbb{P}(\text{disease}|\text{positive}) &= \frac{\mathbb{P}(\text{disease})\mathbb{P}(\text{positive}|\text{disease})}{\mathbb{P}(\text{positive})} \\ &= \frac{(1/5000)(.98)}{(1/5000)(.98) + (4999/5000)(.01)} \\ &\approx 1/51\end{aligned}$$

It's not often highlighted, but this example also demonstrates the problem with interpreting p-values to decide whether a hypothesis is likely true or false: If a patient tests positive for the disease, the p-value for the null hypothesis (no disease) is .01.

P-values are useful when your objective is to control false-positive rates.

CLASSIC BAYES THEOREM EXAMPLE

Classic counter-intuitive example demonstrating Bayes' Theorem: Suppose the prevalence of a disease in the general population is 1 in 5000. A test for this disease has a 1% false positive rate and a 2% false negative rate. If this test gives you a positive result, find the probability that you have the disease.

$$\begin{aligned}\mathbb{P}(\text{disease}|\text{positive}) &= \frac{\mathbb{P}(\text{disease})\mathbb{P}(\text{positive}|\text{disease})}{\mathbb{P}(\text{positive})} \\ &= \frac{(1/5000)(.98)}{(1/5000)(.98) + (4999/5000)(.01)} \\ &\approx 1/51\end{aligned}$$

It's not often highlighted, but this example also demonstrates the problem with interpreting p-values to decide whether a hypothesis is likely true or false: If a patient tests positive for the disease, the p-value for the null hypothesis (no disease) is .01.

P-values are useful when your objective is to control false-positive rates. (Even then, it can fail! e.g. selective submission of experimental results.)

CLASSIC BAYES THEOREM EXAMPLE

Classic counter-intuitive example demonstrating Bayes' Theorem: Suppose the prevalence of a disease in the general population is 1 in 5000. A test for this disease has a 1% false positive rate and a 2% false negative rate. If this test gives you a positive result, find the probability that you have the disease.

$$\begin{aligned}\mathbb{P}(\text{disease}|\text{positive}) &= \frac{\mathbb{P}(\text{disease})\mathbb{P}(\text{positive}|\text{disease})}{\mathbb{P}(\text{positive})} \\ &= \frac{(1/5000)(.98)}{(1/5000)(.98) + (4999/5000)(.01)} \\ &\approx 1/51\end{aligned}$$

It's not often highlighted, but this example also demonstrates the problem with interpreting p-values to decide whether a hypothesis is likely true or false: If a patient tests positive for the disease, the p-value for the null hypothesis (no disease) is .01.

P-values are useful when your objective is to control false-positive rates. (Even then, it can fail! e.g. selective submission of experimental results.)

In most contexts, this is an objective we shouldn't care much about.

The philosophy behind Bayesian statistics is explained and worked out in detail in *Probability Theory: The Logic of Science* (2003) by Edwin Jaynes.

The philosophy behind Bayesian statistics is explained and worked out in detail in *Probability Theory: The Logic of Science* (2003) by Edwin Jaynes.

Main points providing foundation for Bayesian statistics:

- The only "sensible" way of quantifying beliefs is with probability distributions.

The philosophy behind Bayesian statistics is explained and worked out in detail in *Probability Theory: The Logic of Science* (2003) by Edwin Jaynes.

Main points providing foundation for Bayesian statistics:

- The only "sensible" way of quantifying beliefs is with probability distributions.
- In particular, the mathematically correct way of updating one's beliefs in light of new data is ...

The philosophy behind Bayesian statistics is explained and worked out in detail in *Probability Theory: The Logic of Science* (2003) by Edwin Jaynes.

Main points providing foundation for Bayesian statistics:

- The only "sensible" way of quantifying beliefs is with probability distributions.
- In particular, the mathematically correct way of updating one's beliefs in light of new data is ... **Bayes' Theorem**.

The philosophy behind Bayesian statistics is explained and worked out in detail in *Probability Theory: The Logic of Science* (2003) by Edwin Jaynes.

Main points providing foundation for Bayesian statistics:

- The only "sensible" way of quantifying beliefs is with probability distributions.
- In particular, the mathematically correct way of updating one's beliefs in light of new data is ... **Bayes' Theorem**.

Critics' objections against Bayesian statistics: A **prior belief** must be brought into the analysis:

The philosophy behind Bayesian statistics is explained and worked out in detail in *Probability Theory: The Logic of Science* (2003) by Edwin Jaynes.

Main points providing foundation for Bayesian statistics:

- The only "sensible" way of quantifying beliefs is with probability distributions.
- In particular, the mathematically correct way of updating one's beliefs in light of new data is ... **Bayes' Theorem**.

Critics' objections against Bayesian statistics: A **prior belief** must be brought into the analysis:

- can be *subjective*

BAYESIAN PHILOSOPHY

The philosophy behind Bayesian statistics is explained and worked out in detail in *Probability Theory: The Logic of Science* (2003) by Edwin Jaynes.

Main points providing foundation for Bayesian statistics:

- The only "sensible" way of quantifying beliefs is with probability distributions.
- In particular, the mathematically correct way of updating one's beliefs in light of new data is ... **Bayes' Theorem**.

Critics' objections against Bayesian statistics: A **prior belief** must be brought into the analysis:

- can be *subjective*
- can be *difficult to quantify*

Manager: I wonder if our average costs will be different for females versus males.

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates.

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

BAYESIAN'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Bayesian: The probability that males have higher average blood pressure is about .84.

BAYESIAN'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Bayesian: The probability that males have higher average blood pressure is about .84.

Manager: Got it!

BAYESIAN'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Bayesian: The probability that males have higher average blood pressure is about .84.

Manager: Got it! Can you also tell me how likely it is that this scheme will cost us more than \$600,000 dollars?

BAYESIAN'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Bayesian: The probability that males have higher average blood pressure is about .84.

Manager: Got it! Can you also tell me how likely it is that this scheme will cost us more than \$600,000 dollars?

Bayesian: The probability is about .087. In fact, here's the whole distribution of cost...

BAYESIAN'S CONVERSATION WITH MANAGEMENT

Manager: I wonder if our average costs will be different for females versus males. I have no idea what the average blood pressures are for males and females, but I want to know what our data indicates. Tell me, *do males seem to have higher blood pressure on average?*

Bayesian: The probability that males have higher average blood pressure is about .84.

Manager: Got it! Can you also tell me how likely it is that this scheme will cost us more than \$600,000 dollars?

Bayesian: The probability is about .087. In fact, here's the whole distribution of cost...

Manager: [unable to speak... weeping tears of joy]

END

Thank you, SageSure!

Any questions?