

Math 16:642:623  
Computational Finance  
Homework 5

NITISH KANABAR  
`nitish@eden.rutgers.edu`

February 23, 2010

I developed and tested my code on Mac OS X using GCC Version 4.2.1.

## Algorithms

### Sobol Sequence Generator

My Sobol sequence generator is based on code adapted from S. Joe and F. Y. Kuo which is located here: <http://web.maths.unsw.edu.au/~fkuo/sobol/>

A Sobol sequence is a type of  $(t, d)$ -sequence in the class of low-discrepancy sequences. The generator is implemented in the files Sobol.{cpp,h}. The algorithm of generating Sobol sequences is as described in Notes on generating Sobol' sequences, [?]. We begin with a set of primitive polynomials and direction numbers for dimensions 1 through 12 as defined by Joe and Kuo. The Sobol class encapsulates these numbers in the data-members  $a$  and  $m$ . The Sobol class is limited to generating Sobol sequences upto dimension=12. The Sobol constructor accepts two arguments - dimensionality and the dimension. The dimensionality indicates the total number of points that are required to be generated while the dimension refers to the number of points in each set. Hence the number of sets is dimensionality / dimension. The GetUniforms method generates sequences of uniforms while the GetGaussian method converts these uniforms into a gaussian sequence by the InverseTransform method.

### Park-Miller Generator

The Park-Miller generator is the standard Linear Congruential Generator using the following algorithm.

$$\begin{aligned}x_{i+1} &= ax_i \bmod m \\x_0 &= seed \\m &= 2^{31} - 1, a = 16807 \\u_{i+1} &= \frac{x_{i+1}}{m}\end{aligned}$$

I have used the Park-Miller generator implemented by Mark Joshi.

### Stratified Generator

The stratified generator accepts a Park-Miller generator as its inner-generator. A uniform from the inner generator is converted to the stratified sample using the following formula:

$$V := a_{i-1} + (a_i - a_{i-1})U$$

where  $U \sim \text{Unif}[0,1]$  is a Park-Miller uniform. The stratified normal random is obtained by converting the stratified uniform to the a gaussian using the Inverse Transform method.

## Closed Form Formula

The following closed-form formula was used for geometric options:

$$c = Se^{(b-r)(T-t)}N(d_1) - Ke^{-r(T-t)}N(d_2)$$

Where  $N(x)$  is the cumulative normal distribution function of:

$$d_1 = \frac{\ln(\frac{S}{K}) + (b + \frac{\sigma_a^2}{2})T}{\sigma_a\sqrt{T}}$$

$$d_2 = \frac{\ln(\frac{S}{K}) + (b - \frac{\sigma_a^2}{2})T}{\sigma_a\sqrt{T}}$$

$$\sigma_a = \frac{\sigma}{\sqrt{3}}$$

$$b = \frac{1}{2}(r - d - \frac{\sigma^2}{6})$$

## Results

Result obtained from Kerry Back's spread-sheet for the geometric call is: 3.811381662

Due to an upcoming mid-term I was unable to debug some issues with the Sobol and Stratified generators when used with path-dependent options. I have presented the results as implemented.

Closed-form vanilla call option = 10.0201

MC vanilla call, Park-Miller uniforms, antithetics = 9.97437

MC vanilla call, Park-Miller uniforms, stratified = 8.94626

QMC vanilla call, Sobol sequence = 10.0047

MC arithmetic Asian call, Park-Miller uniforms, antithetics = 23.1931

MC asian call, Park-Miller uniforms, stratified = 22.0335

QMC arithmetic asian call, Sobol sequence = 0

Closed-form geometric asian call = 3.38709

MC geometric asian call, Park-Miller uniforms, antithetics = nan

MC geometric call, Park-Miller uniforms, stratified = nan

QMC geometric asian call, Sobol sequence = 0

## References

- [1] Stephen Joe and Frances Y. Kuo: *Notes on generating Sobol' sequences*
- [2] Glasserman, Paul: *Monte Carlo Methods in Financial Engineering*