



quantumformalism.com

Zaiku Group
Quantum Formalism

Mathematical Tools of Quantum Mechanics: Introduction to Hilbert Spaces

Dr. M. Arnott

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

- (a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

- (b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*.
Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

- (c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

- (d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*.
Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*.

Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*.

Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*.

Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*.
Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*.
Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

- (a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

- (b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

- (c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

- (d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

- (a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

- (b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*.
Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

- (c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

- (d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

(a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

(b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

(c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

(d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

- (a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

- (b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

- (c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

- (d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

- (a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

- (b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

- (c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

- (d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

Important Types of Operator

Let $\mathcal{H}, \mathcal{K}, \mathcal{J}, \mathcal{L}$ be Hilbert spaces.

- (a) Identity - $I \in \mathcal{B}(\mathcal{H}) : x \mapsto x$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{H}; \mathcal{J})$, and any $S \in \mathcal{B}(\mathcal{J}; \mathcal{H})$, we have $IS = S$, $TI = T$.

Zero - $0 \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : x \mapsto 0$ for all $x \in \mathcal{H}$. For any $T \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and any $S \in \mathcal{B}(\mathcal{L}; \mathcal{H})$, we have $0S = 0$, $T0 = 0$.

Remark: The lack of subscript on these operators should not cause confusion - but be careful!

- (b) Hermitian - $T \in \mathcal{B}(\mathcal{H}) : T^* = T$; These operators are also known as *self-adjoint*. Examples: we always have that I and 0 are hermitian. The spectrum of a Hermitian operator is always a subset of \mathbb{R} . Examples: The standard Pauli operations are hermitian.

- (c) Isometry - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K}) : \|Tx\|_{\mathcal{K}} = \|x\|_{\mathcal{H}}$ for all $x \in \mathcal{H}$. Examples: rotations and reflections in \mathbb{R}^2 are always isometric.

An operator $U \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is an isometry if and only if for every $x, y \in \mathcal{H}$, we have that $\langle x, y \rangle = \langle Ux, Uy \rangle$.

- (d) Isomorphism (invertible) - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ such that $\exists S \in \mathcal{B}(\mathcal{K}; \mathcal{H})$ with $TS = I_{\mathcal{K}}$ and $ST = I_{\mathcal{H}}$. We write $S = T^{-1}$ in this case.

If T is invertible, then T^{-1} is unique (*exercise (easy) : prove this!*).

If there exists an isomorphism from \mathcal{H} to \mathcal{K} , we write $\mathcal{H} \cong \mathcal{K}$. If also this isomorphism is an isometry, we write $\mathcal{H} \equiv \mathcal{K}$. (*Exercise (medium) : show that \cong and \equiv are equivalence relations on the class of Hilbert spaces.*)

- (e) Unitary - $U \in \mathcal{B}(\mathcal{H})$ such that $U^{-1} = U^*$. Unitary operators are *isometric isomorphisms*, but there exist isometries that are not unitary (*exercise (easy) : prove this!*).

In quantum computing, the set of unitary operators on the state space \mathbb{C}^{2^n} form the possible operations applicable to the n qubits of the system.

The set of all unitary operators on a Hilbert space forms a *group* with operator composition and identity I (*exercise (medium) : prove this!*).

The spectrum of a unitary operator is always a subset of the unit circle of \mathbb{C} , i.e.

$$\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \{e^{i\theta} : 0 \leq \theta \leq 2\pi\}.$$

- (f) Normal - $T \in \mathcal{B}(\mathcal{H})$ is normal if it commutes with T^* , meaning that $T^*T = TT^*$. It is easy to see that unitary and Hermitian operators are normal. (*Exercise (medium) : an operator S is skew-Hermitian if $S^* = -S$. Show that skew-Hermitian operators are normal*).

- (e) Unitary - $U \in \mathcal{B}(\mathcal{H})$ such that $U^{-1} = U^*$. Unitary operators are *isometric isomorphisms*, but there exist isometries that are not unitary (*exercise (easy) : prove this!*).

In quantum computing, the set of unitary operators on the state space \mathbb{C}^{2^n} form the possible operations applicable to the n qubits of the system.

The set of all unitary operators on a Hilbert space forms a *group* with operator composition and identity I (*exercise (medium) : prove this!*).

The spectrum of a unitary operator is always a subset of the unit circle of \mathbb{C} , i.e.

$$\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \{e^{i\theta} : 0 \leq \theta \leq 2\pi\}.$$

- (f) Normal - $T \in \mathcal{B}(\mathcal{H})$ is normal if it commutes with T^* , meaning that $T^*T = TT^*$. It is easy to see that unitary and Hermitian operators are normal. (*Exercise (medium) : an operator S is skew-Hermitian if $S^* = -S$. Show that skew-Hermitian operators are normal*).

- (e) Unitary - $U \in \mathcal{B}(\mathcal{H})$ such that $U^{-1} = U^*$. Unitary operators are *isometric isomorphisms*, but there exist isometries that are not unitary (*exercise (easy) : prove this!*).

In quantum computing, the set of unitary operators on the state space \mathbb{C}^{2^n} form the possible operations applicable to the n qubits of the system.

The set of all unitary operators on a Hilbert space forms a *group* with operator composition and identity I (*exercise (medium) : prove this!*).

The spectrum of a unitary operator is always a subset of the unit circle of \mathbb{C} , i.e.

$$\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \{e^{i\theta} : 0 \leq \theta \leq 2\pi\}.$$

- (f) Normal - $T \in \mathcal{B}(\mathcal{H})$ is normal if it commutes with T^* , meaning that $T^*T = TT^*$. It is easy to see that unitary and Hermitian operators are normal. (*Exercise (medium) : an operator S is skew-Hermitian if $S^* = -S$. Show that skew-Hermitian operators are normal*).

- (e) Unitary - $U \in \mathcal{B}(\mathcal{H})$ such that $U^{-1} = U^*$. Unitary operators are *isometric isomorphisms*, but there exist isometries that are not unitary (*exercise (easy) : prove this!*).

In quantum computing, the set of unitary operators on the state space \mathbb{C}^{2^n} form the possible operations applicable to the n qubits of the system.

The set of all unitary operators on a Hilbert space forms a *group* with operator composition and identity I (*exercise (medium) : prove this!*).

The spectrum of a unitary operator is always a subset of the unit circle of \mathbb{C} , i.e.

$$\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \{e^{i\theta} : 0 \leq \theta \leq 2\pi\}.$$

- (f) Normal - $T \in \mathcal{B}(\mathcal{H})$ is normal if it commutes with T^* , meaning that $T^*T = TT^*$. It is easy to see that unitary and Hermitian operators are normal. (*Exercise (medium) : an operator S is skew-Hermitian if $S^* = -S$. Show that skew-Hermitian operators are normal*).

- (e) Unitary - $U \in \mathcal{B}(\mathcal{H})$ such that $U^{-1} = U^*$. Unitary operators are *isometric isomorphisms*, but there exist isometries that are not unitary (*exercise (easy) : prove this!*).

In quantum computing, the set of unitary operators on the state space \mathbb{C}^{2^n} form the possible operations applicable to the n qubits of the system.

The set of all unitary operators on a Hilbert space forms a *group* with operator composition and identity I (*exercise (medium) : prove this!*).

The spectrum of a unitary operator is always a subset of the unit circle of \mathbb{C} , i.e.

$$\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \{e^{i\theta} : 0 \leq \theta \leq 2\pi\}.$$

- (f) Normal - $T \in \mathcal{B}(\mathcal{H})$ is normal if it commutes with T^* , meaning that $T^*T = TT^*$. It is easy to see that unitary and Hermitian operators are normal. (*Exercise (medium) : an operator S is skew-Hermitian if $S^* = -S$. Show that skew-Hermitian operators are normal*).

- (e) Unitary - $U \in \mathcal{B}(\mathcal{H})$ such that $U^{-1} = U^*$. Unitary operators are *isometric isomorphisms*, but there exist isometries that are not unitary (*exercise (easy) : prove this!*).

In quantum computing, the set of unitary operators on the state space \mathbb{C}^{2^n} form the possible operations applicable to the n qubits of the system.

The set of all unitary operators on a Hilbert space forms a *group* with operator composition and identity I (*exercise (medium) : prove this!*).

The spectrum of a unitary operator is always a subset of the unit circle of \mathbb{C} , i.e.

$$\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \{e^{i\theta} : 0 \leq \theta \leq 2\pi\}.$$

- (f) Normal - $T \in \mathcal{B}(\mathcal{H})$ is normal if it commutes with T^* , meaning that $T^*T = TT^*$. It is easy to see that unitary and Hermitian operators are normal. (*Exercise (medium) : an operator S is skew-Hermitian if $S^* = -S$. Show that skew-Hermitian operators are normal*).

- (e) Unitary - $U \in \mathcal{B}(\mathcal{H})$ such that $U^{-1} = U^*$. Unitary operators are *isometric isomorphisms*, but there exist isometries that are not unitary (*exercise (easy) : prove this!*).

In quantum computing, the set of unitary operators on the state space \mathbb{C}^{2^n} form the possible operations applicable to the n qubits of the system.

The set of all unitary operators on a Hilbert space forms a *group* with operator composition and identity I (*exercise (medium) : prove this!*).

The spectrum of a unitary operator is always a subset of the unit circle of \mathbb{C} , i.e.

$$\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \{e^{i\theta} : 0 \leq \theta \leq 2\pi\}.$$

- (f) Normal - $T \in \mathcal{B}(\mathcal{H})$ is normal if it commutes with T^* , meaning that $T^*T = TT^*$. It is easy to see that unitary and Hermitian operators are normal. (*Exercise (medium) : an operator S is skew-Hermitian if $S^* = -S$. Show that skew-Hermitian operators are normal*).

- (e) Unitary - $U \in \mathcal{B}(\mathcal{H})$ such that $U^{-1} = U^*$. Unitary operators are *isometric isomorphisms*, but there exist isometries that are not unitary (*exercise (easy) : prove this!*).

In quantum computing, the set of unitary operators on the state space \mathbb{C}^{2^n} form the possible operations applicable to the n qubits of the system.

The set of all unitary operators on a Hilbert space forms a *group* with operator composition and identity I (*exercise (medium) : prove this!*).

The spectrum of a unitary operator is always a subset of the unit circle of \mathbb{C} , i.e.

$$\sigma(T) \subseteq \{\lambda \in \mathbb{C} : |\lambda| = 1\} = \{e^{i\theta} : 0 \leq \theta \leq 2\pi\}.$$

- (f) Normal - $T \in \mathcal{B}(\mathcal{H})$ is normal if it commutes with T^* , meaning that $T^*T = TT^*$. It is easy to see that unitary and Hermitian operators are normal. (*Exercise (medium) : an operator S is skew-Hermitian if $S^* = -S$. Show that skew-Hermitian operators are normal*).

Important Types of Operator

- (g) Rank-one - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with one-dimensional image. For these operators, there exist $u \in \mathcal{H}$, $v \in \mathcal{K}$ with $Tx = \langle x, u \rangle v$ for all $x \in \mathcal{H}$.
Finite-Rank - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with finite-dimensional image. Such operators are always expressible as linear combinations of rank-one operators.
- (h) Compact - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is compact if for every bounded sequence $(x_n)_{n \in \mathbb{N}} \subset \mathcal{H}$, there is a convergent subsequence to $(Tx_n)_{n \in \mathbb{N}}$. Equivalently, T is compact if the set $T(B_{\mathcal{H}})$ is compact, where $B_{\mathcal{H}} = \{h \in \mathcal{H} : \|h\| \leq 1\}$.
(Exercises - (medium) : Show that if T is compact, then so is STR for every $S \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and every $R \in \mathcal{B}(\mathcal{L}; \mathcal{H})$. Next, show that if T is compact, then so is T^* .) Every finite-rank operator is compact.
- (i) Projection - Recall that for every closed subspace \mathcal{L} of \mathcal{H} , there exists another subspace \mathcal{L}^{\perp} for which $\mathcal{L} \oplus \mathcal{L}^{\perp} = \mathcal{H}$. The projection $P_{\mathcal{L}} \in \mathcal{B}(\mathcal{H})$ is the function which maps a sum $l + l' \mapsto l$ for every $l \in \mathcal{L}$ and $l' \in \mathcal{L}^{\perp}$. Projections are idempotent, i.e., $P_{\mathcal{L}}P_{\mathcal{L}} = P_{\mathcal{L}}$.
(Exercises - (easy) : Show that $P_{\mathcal{L}}P_{\mathcal{L}^{\perp}} = 0$. Show that $\|P_{\mathcal{L}}\| = 1$ if $\{0\} \subsetneq \mathcal{L} \subset \mathcal{H}$.)
- (j) Tensor Product - If $T \in \mathcal{B}(\mathcal{H})$ and $S \in \mathcal{B}(\mathcal{K})$, we write $T \otimes S \in \mathcal{B}(\mathcal{H} \otimes \mathcal{K})$ for the operator which acts on pure tensors $h \otimes k$ for $h \in \mathcal{H}$ and $k \in \mathcal{K}$ by $h \otimes k \mapsto Th \otimes Tk$, and extend by linearity. Notice that not every bounded operator on $\mathcal{H} \otimes \mathcal{K}$ is formed in this way. (Intuition for QIS people : Not every gate on a n -qubit system can be deconstructed using only 1-qubit gates, else we would have no entangling gates.)

Important Types of Operator

- (g) Rank-one - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with one-dimensional image. For these operators, there exist $u \in \mathcal{H}$, $v \in \mathcal{K}$ with $Tx = \langle x, u \rangle v$ for all $x \in \mathcal{H}$.
Finite-Rank - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with finite-dimensional image. Such operators are always expressible as linear combinations of rank-one operators.
- (h) Compact - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is compact if for every bounded sequence $(x_n)_{n \in \mathbb{N}} \subset \mathcal{H}$, there is a convergent subsequence to $(Tx_n)_{n \in \mathbb{N}}$. Equivalently, T is compact if the set $T(B_{\mathcal{H}})$ is compact, where $B_{\mathcal{H}} = \{h \in \mathcal{H} : \|h\| \leq 1\}$.
(Exercises - (medium) : Show that if T is compact, then so is STR for every $S \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and every $R \in \mathcal{B}(\mathcal{L}; \mathcal{H})$. Next, show that if T is compact, then so is T^* .) Every finite-rank operator is compact.
- (i) Projection - Recall that for every closed subspace \mathcal{L} of \mathcal{H} , there exists another subspace \mathcal{L}^\perp for which $\mathcal{L} \oplus \mathcal{L}^\perp = \mathcal{H}$. The projection $P_{\mathcal{L}} \in \mathcal{B}(\mathcal{H})$ is the function which maps a sum $l + l' \mapsto l$ for every $l \in \mathcal{L}$ and $l' \in \mathcal{L}^\perp$. Projections are idempotent, i.e., $P_{\mathcal{L}}P_{\mathcal{L}} = P_{\mathcal{L}}$.
(Exercises - (easy) : Show that $P_{\mathcal{L}}P_{\mathcal{L}^\perp} = 0$. Show that $\|P_{\mathcal{L}}\| = 1$ if $\{0\} \subsetneq \mathcal{L} \subset \mathcal{H}$.)
- (j) Tensor Product - If $T \in \mathcal{B}(\mathcal{H})$ and $S \in \mathcal{B}(\mathcal{K})$, we write $T \otimes S \in \mathcal{B}(\mathcal{H} \otimes \mathcal{K})$ for the operator which acts on pure tensors $h \otimes k$ for $h \in \mathcal{H}$ and $k \in \mathcal{K}$ by $h \otimes k \mapsto Th \otimes Tk$, and extend by linearity. Notice that not every bounded operator on $\mathcal{H} \otimes \mathcal{K}$ is formed in this way. (Intuition for QIS people : Not every gate on a n -qubit system can be deconstructed using only 1-qubit gates, else we would have no entangling gates.)

Important Types of Operator

- (g) Rank-one - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with one-dimensional image. For these operators, there exist $u \in \mathcal{H}$, $v \in \mathcal{K}$ with $Tx = \langle x, u \rangle v$ for all $x \in \mathcal{H}$.
Finite-Rank - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with finite-dimensional image. Such operators are always expressible as linear combinations of rank-one operators.
- (h) Compact - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is compact if for every bounded sequence $(x_n)_{n \in \mathbb{N}} \subset \mathcal{H}$, there is a convergent subsequence to $(Tx_n)_{n \in \mathbb{N}}$. Equivalently, T is compact if the set $T(B_{\mathcal{H}})$ is compact, where $B_{\mathcal{H}} = \{h \in \mathcal{H} : \|h\| \leq 1\}$.
(Exercises - (medium) : Show that if T is compact, then so is STR for every $S \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and every $R \in \mathcal{B}(\mathcal{L}; \mathcal{H})$. Next, show that if T is compact, then so is T^* .) Every finite-rank operator is compact.
- (i) Projection - Recall that for every closed subspace \mathcal{L} of \mathcal{H} , there exists another subspace \mathcal{L}^\perp for which $\mathcal{L} \oplus \mathcal{L}^\perp = \mathcal{H}$. The projection $P_{\mathcal{L}} \in \mathcal{B}(\mathcal{H})$ is the function which maps a sum $l + l' \mapsto l$ for every $l \in \mathcal{L}$ and $l' \in \mathcal{L}^\perp$. Projections are idempotent, i.e., $P_{\mathcal{L}}P_{\mathcal{L}} = P_{\mathcal{L}}$.
(Exercises - (easy) : Show that $P_{\mathcal{L}}P_{\mathcal{L}^\perp} = 0$. Show that $\|P_{\mathcal{L}}\| = 1$ if $\{0\} \subsetneq \mathcal{L} \subset \mathcal{H}$.)
- (j) Tensor Product - If $T \in \mathcal{B}(\mathcal{H})$ and $S \in \mathcal{B}(\mathcal{K})$, we write $T \otimes S \in \mathcal{B}(\mathcal{H} \otimes \mathcal{K})$ for the operator which acts on pure tensors $h \otimes k$ for $h \in \mathcal{H}$ and $k \in \mathcal{K}$ by $h \otimes k \mapsto Th \otimes Tk$, and extend by linearity. Notice that not every bounded operator on $\mathcal{H} \otimes \mathcal{K}$ is formed in this way. (Intuition for QIS people : Not every gate on a n -qubit system can be deconstructed using only 1-qubit gates, else we would have no entangling gates.)

Important Types of Operator

- (g) Rank-one - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with one-dimensional image. For these operators, there exist $u \in \mathcal{H}$, $v \in \mathcal{K}$ with $Tx = \langle x, u \rangle v$ for all $x \in \mathcal{H}$.
Finite-Rank - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with finite-dimensional image. Such operators are always expressible as linear combinations of rank-one operators.
- (h) Compact - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is compact if for every bounded sequence $(x_n)_{n \in \mathbb{N}} \subset \mathcal{H}$, there is a convergent subsequence to $(Tx_n)_{n \in \mathbb{N}}$. Equivalently, T is compact if the set $T(B_{\mathcal{H}})$ is compact, where $B_{\mathcal{H}} = \{h \in \mathcal{H} : \|h\| \leq 1\}$.
(Exercises - (medium) : Show that if T is compact, then so is STR for every $S \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and every $R \in \mathcal{B}(\mathcal{L}; \mathcal{H})$. Next, show that if T is compact, then so is T^* .) Every finite-rank operator is compact.
- (i) Projection - Recall that for every closed subspace \mathcal{L} of \mathcal{H} , there exists another subspace \mathcal{L}^\perp for which $\mathcal{L} \oplus \mathcal{L}^\perp = \mathcal{H}$. The projection $P_{\mathcal{L}} \in \mathcal{B}(\mathcal{H})$ is the function which maps a sum $l + l' \mapsto l$ for every $l \in \mathcal{L}$ and $l' \in \mathcal{L}^\perp$. Projections are idempotent, i.e., $P_{\mathcal{L}}P_{\mathcal{L}} = P_{\mathcal{L}}$.
(Exercises - (easy) : Show that $P_{\mathcal{L}}P_{\mathcal{L}^\perp} = 0$. Show that $\|P_{\mathcal{L}}\| = 1$ if $\{0\} \subsetneq \mathcal{L} \subset \mathcal{H}$.)
- (j) Tensor Product - If $T \in \mathcal{B}(\mathcal{H})$ and $S \in \mathcal{B}(\mathcal{K})$, we write $T \otimes S \in \mathcal{B}(\mathcal{H} \otimes \mathcal{K})$ for the operator which acts on pure tensors $h \otimes k$ for $h \in \mathcal{H}$ and $k \in \mathcal{K}$ by $h \otimes k \mapsto Th \otimes Tk$, and extend by linearity. Notice that not every bounded operator on $\mathcal{H} \otimes \mathcal{K}$ is formed in this way. (Intuition for QIS people : Not every gate on a n -qubit system can be deconstructed using only 1-qubit gates, else we would have no entangling gates.)

Important Types of Operator

- (g) Rank-one - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with one-dimensional image. For these operators, there exist $u \in \mathcal{H}$, $v \in \mathcal{K}$ with $Tx = \langle x, u \rangle v$ for all $x \in \mathcal{H}$.
Finite-Rank - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with finite-dimensional image. Such operators are always expressible as linear combinations of rank-one operators.
- (h) Compact - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is compact if for every bounded sequence $(x_n)_{n \in \mathbb{N}} \subset \mathcal{H}$, there is a convergent subsequence to $(Tx_n)_{n \in \mathbb{N}}$. Equivalently, T is compact if the set $T(B_{\mathcal{H}})$ is compact, where $B_{\mathcal{H}} = \{h \in \mathcal{H} : \|h\| \leq 1\}$.
(Exercises - (medium) : Show that if T is compact, then so is STR for every $S \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and every $R \in \mathcal{B}(\mathcal{L}; \mathcal{H})$. Next, show that if T is compact, then so is T^* .) Every finite-rank operator is compact.
- (i) Projection - Recall that for every closed subspace \mathcal{L} of \mathcal{H} , there exists another subspace \mathcal{L}^\perp for which $\mathcal{L} \oplus \mathcal{L}^\perp = \mathcal{H}$. The projection $P_{\mathcal{L}} \in \mathcal{B}(\mathcal{H})$ is the function which maps a sum $l + l' \mapsto l$ for every $l \in \mathcal{L}$ and $l' \in \mathcal{L}^\perp$. Projections are idempotent, i.e., $P_{\mathcal{L}}P_{\mathcal{L}} = P_{\mathcal{L}}$.
(Exercises - (easy) : Show that $P_{\mathcal{L}}P_{\mathcal{L}^\perp} = 0$. Show that $\|P_{\mathcal{L}}\| = 1$ if $\{0\} \subsetneq \mathcal{L} \subset \mathcal{H}$.)
- (j) Tensor Product - If $T \in \mathcal{B}(\mathcal{H})$ and $S \in \mathcal{B}(\mathcal{K})$, we write $T \otimes S \in \mathcal{B}(\mathcal{H} \otimes \mathcal{K})$ for the operator which acts on pure tensors $h \otimes k$ for $h \in \mathcal{H}$ and $k \in \mathcal{K}$ by $h \otimes k \mapsto Th \otimes Tk$, and extend by linearity. Notice that not every bounded operator on $\mathcal{H} \otimes \mathcal{K}$ is formed in this way. (Intuition for QIS people : Not every gate on a n -qubit system can be deconstructed using only 1-qubit gates, else we would have no entangling gates.)

Important Types of Operator

- (g) Rank-one - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with one-dimensional image. For these operators, there exist $u \in \mathcal{H}$, $v \in \mathcal{K}$ with $Tx = \langle x, u \rangle v$ for all $x \in \mathcal{H}$.
Finite-Rank - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ with finite-dimensional image. Such operators are always expressible as linear combinations of rank-one operators.
- (h) Compact - $T \in \mathcal{B}(\mathcal{H}; \mathcal{K})$ is compact if for every bounded sequence $(x_n)_{n \in \mathbb{N}} \subset \mathcal{H}$, there is a convergent subsequence to $(Tx_n)_{n \in \mathbb{N}}$. Equivalently, T is compact if the set $T(B_{\mathcal{H}})$ is compact, where $B_{\mathcal{H}} = \{h \in \mathcal{H} : \|h\| \leq 1\}$.
(Exercises - (medium) : Show that if T is compact, then so is STR for every $S \in \mathcal{B}(\mathcal{K}; \mathcal{J})$ and every $R \in \mathcal{B}(\mathcal{L}; \mathcal{H})$. Next, show that if T is compact, then so is T^* .) Every finite-rank operator is compact.
- (i) Projection - Recall that for every closed subspace \mathcal{L} of \mathcal{H} , there exists another subspace \mathcal{L}^\perp for which $\mathcal{L} \oplus \mathcal{L}^\perp = \mathcal{H}$. The projection $P_{\mathcal{L}} \in \mathcal{B}(\mathcal{H})$ is the function which maps a sum $l + l' \mapsto l$ for every $l \in \mathcal{L}$ and $l' \in \mathcal{L}^\perp$. Projections are idempotent, i.e., $P_{\mathcal{L}}P_{\mathcal{L}} = P_{\mathcal{L}}$.
(Exercises - (easy) : Show that $P_{\mathcal{L}}P_{\mathcal{L}^\perp} = 0$. Show that $\|P_{\mathcal{L}}\| = 1$ if $\{0\} \subsetneq \mathcal{L} \subset \mathcal{H}$.)
- (j) Tensor Product - If $T \in \mathcal{B}(\mathcal{H})$ and $S \in \mathcal{B}(\mathcal{K})$, we write $T \otimes S \in \mathcal{B}(\mathcal{H} \otimes \mathcal{K})$ for the operator which acts on pure tensors $h \otimes k$ for $h \in \mathcal{H}$ and $k \in \mathcal{K}$ by $h \otimes k \mapsto Th \otimes Tk$, and extend by linearity. Notice that not every bounded operator on $\mathcal{H} \otimes \mathcal{K}$ is formed in this way. (Intuition for QIS people : Not every gate on a n -qubit system can be deconstructed using only 1-qubit gates, else we would have no entangling gates.)

Thank you for listening!