

Short Paper

Alice Anonymous^{a,*}, Bob Security^b, Cat Memes^b, Derek Zoolander

^a*Some Institute of Technology Street Address City State Postal Code*

^b*Another University Street Address City State Postal Code*

Abstract

This is the abstract.

It consists of two paragraphs.

Keywords: keyword1, keyword2

Please make sure that your manuscript follows the guidelines in the Guide for Authors of the relevant journal. It is not necessary to typeset your manuscript in exactly the same way as an article, unless you are submitting to a camera-ready copy (CRC) journal.

For detailed instructions regarding the elsevier article class, see <https://www.elsevier.com/authors/policies-and-guidelines/latex-instructions>

1. Bibliography styles

Here are two sample references: [Feynman and Vernon Jr. \(1963; Dirac, 1953\)](#).

By default, natbib will be used with the `authoryear` style, set in `classoption` variable in YAML. You can sets extra options with `natbiboptions` variable in YAML header. Example

```
natbiboptions: longnamesfirst,angle,semicolon
```

There are various more specific bibliography styles available at https://support.stmdocs.in/wiki/index.php?title=Model-wise_bibliographic_style_files. To use one of these, add it in the header using, for example, `biblio-style: model1-num-names`.

1.1. Using CSL

If `cite-method` is set to `citeproc` in `elsevier_article()`, then pandoc is used for citations instead of `natbib`. In this case, the `cs1` option is used to format the references. By default, this template will provide an appropriate style, but alternative `cs1` files are available from <https://www.zotero.org/styles?q=elsevier>. These can be downloaded and stored locally, or the url can be used as in the example header.

*Corresponding author

Email addresses: `alice@example.com` (Alice Anonymous), `bob@example.com` (Bob Security), `cat@example.com` (Cat Memes), `derek@example.com` (Derek Zoolander)

2. Equations

Here is an equation:

$$f_X(x) = \left(\frac{\alpha}{\beta}\right) \left(\frac{x}{\beta}\right)^{\alpha-1} e^{-\left(\frac{x}{\beta}\right)^\alpha}; \alpha, \beta, x > 0.$$

Inline equations work as well: $\sum_{i=2}^{\infty} \{\alpha_i^\beta\}$

3. Figures and tables

Figure 1 is generated using an R chunk.




Figure 1: A meaningless scatterplot

4. Tables coming from R

Tables can also be generated using R chunks, as shown in Table 1 example.

```
knitr::kable(head(mtcars)[,1:4])
```

Table 1: Caption centered above table

	mpg	cyl	disp	hp
Mazda RX4	21.0	6	160	110
Mazda RX4 Wag	21.0	6	160	110
Datsun 710	22.8	4	108	93
Hornet 4 Drive	21.4	6	258	110
Hornet Sportabout	18.7	8	360	175

	mpg	cyl	disp	hp
Valiant	18.1	6	225	105

References

- P. A. M. Dirac. The Lorentz transformation and absolute time. *Physica*, 19(1--12):888–896, 1953. doi: 10.1016/S0031-8914(53)80099-6.
- R. P Feynman and F. L Vernon Jr. The theory of a general quantum system interacting with a linear dissipative system. *Annals of Physics*, 24:118–173, 1963. doi: 10.1016/0003-4916(63)90068-X.