MONASH University

Expert advice from experts

Professor Marie Curie
Nobel Prize, PhD

Dr Pierre Curie
Nobel Prize, PhD

Report for
Acme Corporation

13 August 2026

MONASH
BUSINESS
SCHOOL

Department of Econometrics & Business Statistics

(03) 9905 2478
BusEco-Econometrics@monash.edu

ABN: 12 377 614 012

AACSB ACCREDITED EQUIS ACCREDITED ASSOCIATION AMBA ACCREDITED

1 Introduction

Kernel density estimation is a non-parametric way to estimate the probability density function of a random variable (Rice, 2007, p390). The kernel density estimate (KDE) is defined as:

$$\hat{f}(x) = \frac{1}{nh} \sum_{i=1}^{n} K\left(\frac{x - x_i}{h}\right), \tag{1}$$

where K is the kernel function, h is the bandwidth, and $x_1, x_2, \ldots, x_n$ are the observed data points. A common kernel function is the Gaussian kernel, defined as:

$$K(u) = \frac{1}{\sqrt{2\pi}} e^{-u^2/2}.$$

Figure 1 shows a KDE computed using Equation 1 with a Gaussian kernel, and h chosen using the Sheather-Jones method (Sheather & Jones, 1991).

```
library(tidyverse)
set.seed(2022 - 12 - 20)
df <- tibble(x = rnorm(200))
df |>
  ggplot(aes(x = x)) +
  geom_density(bw = "sj") +
  geom_rug()
```




Figure 1: *A kernel density estimate obtained using 200 simulated data points from a N(0,1) distribution.*

References

Rice, J. A. (2007). *Mathematical Statistics and Data Analysis* (3rd edition). Duxbury.

Sheather, S. J., & Jones, M. C. (1991). A reliable data-based bandwidth selection method for kernel density estimation. *Journal of the Royal Statistical Society, Series B*, *53*(3), 683–690. https://doi.org/10.1111/j.2517-6161.1991.tb01857.x