Chapter 1

Section 1.1: Introduction

Welcome to the first chapter of our textbook. This section introduces the basic concepts we will build on throughout the course.

Subsection 1.1.1: Motivation

Why do we study this subject? Because it appears in many real-world problems, from engineering to economics.

Subsection 1.1.2: Notation

We will use the following notation throughout:

- x denotes a scalar variable.
- $\mathbf{x}$ denotes a vector.
- A denotes a matrix.
- $\|\cdot\|$ denotes the Euclidean norm.

Section 1.2: Definitions

Definition 1.2.1. A *widget* is a finite collection of gears together with a rule describing how they mesh.

Definition 1.2.2. The *order* of a widget is the number of gears it contains.

Section 1.3: Theorems and Proofs

Theorem 1.3.1. Every widget of order $n \geq 1$ has at least $n - 1$ meshing pairs.

Proof. We argue by induction on n. The base case $n = 1$ is trivial, since a single gear has no pairs at all. Assume the statement holds for all widgets of order n, and let W be a widget of order $n + 1$. Remove any gear g from W; by the induction hypothesis the remaining widget has at least $n - 1$ meshing pairs. Reattaching g creates at least one new pair, so W has at least $(n - 1) + 1 = n$ meshing pairs. $\square$

Corollary 1.3.2. A widget of order n is connected if it has exactly $n - 1$ meshing pairs.

Lemma 1.3.3. If two widgets share a gear, their union is a widget.

Proof. Immediate from the definition of a widget. $\square$

Section 1.4: Examples

Example 1.4.1. The widget with gears $\{a, b, c\}$ and meshing pairs $\{a, b\}$, $\{b, c\}$ has order 3 and is connected.

Example 1.4.2. Adding a gear d that meshes only with a to the widget above yields a widget of order 4 with 3 meshing pairs.

Example 1.4.3. The empty collection of gears is not a widget, since it violates the order $n \geq 1$.

Section 1.5: Exercises

Exercise 1.5.1. Prove that every widget of order $n \geq 2$ has at least two gears that mesh with exactly one other gear.

Exercise 1.5.2. Show that the converse of Theorem 1.3.1 fails: exhibit a widget with more than $n - 1$ meshing pairs.

Exercise 1.5.3. Let W be a widget of order n with exactly $n - 1$ meshing pairs. Prove that W is a tree.

Exercise 1.5.4. Define a *subwidget* of W to be a widget whose gears and meshing pairs are all contained in those of W. How many subwidgets does a widget of order n have?

Exercise 1.5.5. Let W_1 and W_2 be widgets of orders n_1 and n_2 sharing exactly k gears. Bound the order of $W_1 \cup W_2$ in terms of n_1, n_2, and k.

Exercise 1.5.6. Prove Lemma 1.3.3.

Exercise 1.5.7. Let W be a widget of order $n \geq 3$. Show that removing any gear of degree 1 from W yields a widget.

Exercise 1.5.8. Define the *girth* of a widget to be the length of its shortest cycle of meshing pairs, or ∞ if it has none. Prove that every widget of order n has girth at most n.

Exercise 1.5.9. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.10. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.11. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.12. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.13. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.14. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.15. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.16. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.17. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.18. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.19. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Exercise 1.5.20. Let W be a widget of order n with girth g. Prove that W has at least $n - 1 + \lfloor g/2 \rfloor$ meshing pairs.

Section 1.6: Solutions

Solution to Exercise 1.5.1. Take a leaf of a spanning tree of W; it meshes with exactly one gear.

Solution to Exercise 1.5.2. Any widget of order 3 with all three pairs meshing has $3 > 2$ pairs.

Solution to Exercise 1.5.3. By Corollary 1.3.2, W is connected with $n - 1$ edges, hence a tree.

Solution to Exercise 1.5.4. Each subset of gears of size $k \geq 1$ that induces a connected meshing graph is a subwidget, so the count is $\sum_{k=1}^{n} c_k$ where c_k counts connected k-gear patterns.

Solution to Exercise 1.5.5. The shared k gears and meshing pairs are counted twice, so $\text{order}(W_1 \cup W_2) \leq n_1 + n_2 - k$.

Solution to Exercise 1.5.6. Restrict the meshing rule to the gears of W other than g; the result is a widget by the definition.

Solution to Exercise 1.5.7. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.8. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.9. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.10. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.11. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.12. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.13. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.14. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.15. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.16. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.17. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.18. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.19. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.

Solution to Exercise 1.5.20. Let C be a shortest cycle of meshing pairs in W, so $|C| = g$. Any gear not on C has at most two neighbors on C, so deleting it leaves C intact. Hence W contains a cycle of length g, and its girth is at most n.