

# A Statistical Comparison Between Proton Microinstabilities and Nonlinear Effects in Space Plasmas

Ramiz A. Qudsi<sup>1</sup>,

Bennett A. Maruca<sup>1,2</sup>, R. Bandyopadhyay<sup>1</sup>, A. Chasapis<sup>1</sup>, S. P. Gary<sup>3</sup>,  
R. Chhiber<sup>1</sup>, T. N. Parashar<sup>1</sup>, W. H. Matthaeus<sup>1,2</sup>, J. L. Burch<sup>4</sup>,  
T. E. Moore<sup>5</sup>, C. J. Pollock<sup>6</sup>, B. J. Giles<sup>5</sup>, W. R. Paterson<sup>5</sup>, J. Dorelli<sup>5</sup>,  
D. J. Gershman<sup>5</sup>, R. B. Torbert<sup>6</sup>, & R. J. Strangeway<sup>8</sup>

<sup>1</sup>Univ. Delaware    <sup>2</sup>Bartol Res. Inst.    <sup>3</sup>Space Sci. Inst.    <sup>4</sup>Southwest Res. Inst.  
<sup>5</sup>NASA/GSFC    <sup>6</sup>Denali Sci.    <sup>7</sup>Univ. New Hampshire    <sup>8</sup>Univ. California, LA

ahmadr@udel.edu

12 November 2019



Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy  $\Rightarrow$  Non-Maxwellian VDF

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy  $\Rightarrow$  Non-Maxwellian VDF  $\Rightarrow$  Free Energy

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy  $\Rightarrow$  Non-Maxwellian VDF  $\Rightarrow$  Free Energy  $\Rightarrow$  Unstable

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy  $\Rightarrow$  Non-Maxwellian VDF  $\Rightarrow$  Free Energy  $\Rightarrow$  Unstable  $\Rightarrow$  Micro-Instabilities

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy  $\Rightarrow$  Non-Maxwellian VDF  $\Rightarrow$  Free Energy  $\Rightarrow$  Unstable  $\Rightarrow$  Micro-Instabilities  $\Rightarrow$  Growth rate ( $\gamma_j$ )

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy  $\Rightarrow$  Non-Maxwellian VDF  $\Rightarrow$  Free Energy  $\Rightarrow$  Unstable  $\Rightarrow$  Micro-Instabilities  $\Rightarrow$  Growth rate ( $\gamma_j$ )

	Parallel ( $\mathbf{k} \parallel \mathbf{B}$ ) & Propagating ( $\omega_r > 0$ )	Oblique ( $\mathbf{k} \nparallel \mathbf{B}$ ) & Non-Propagating ( $\omega_r = 0$ )
$T_{\perp j} > T_{\parallel j}$ ( $R_j > 1$ )	Ion-cyclotron (Alfven mode)	Mirror (kinetic slow mode)
$T_{\perp j} < T_{\parallel j}$ ( $R_j < 1$ )	Parallel firehose (fast/whistler mode)	Oblique firehose (Alfven mode)

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy  $\Rightarrow$  Non-Maxwellian VDF  $\Rightarrow$  Free Energy  $\Rightarrow$  Unstable  $\Rightarrow$  Micro-Instabilities  $\Rightarrow$  Growth rate ( $\gamma_j$ )

	Parallel ( $\mathbf{k} \parallel \mathbf{B}$ ) & Propagating ( $\omega_r > 0$ )	Oblique ( $\mathbf{k} \nparallel \mathbf{B}$ ) & Non-Propagating ( $\omega_r = 0$ )
$T_{\perp j} > T_{\parallel j}$ ( $R_j > 1$ )	Ion-cyclotron (Alfven mode)	Mirror (kinetic slow mode)
$T_{\perp j} < T_{\parallel j}$ ( $R_j < 1$ )	Parallel firehose (fast/whistler mode)	Oblique firehose (Alfven mode)

$$\beta_{\parallel j} \equiv \frac{n_j k_B T_{\parallel j}}{B^2 / (2 \mu_0)}$$


# Ion Temperature-Anisotropy Instabilities

	Parallel ( $\mathbf{k} \parallel \mathbf{B}$ ) & Propagating ( $\omega_r > 0$ )	Oblique ( $\mathbf{k} \nparallel \mathbf{B}$ ) & Non-Prop. ( $\omega_r = 0$ )
$T_{\perp j} > T_{\parallel j}$ ( $R_j > 1$ )	Ion-cyclotron (Alfven mode)	Mirror (kinetic slow mode)
$T_{\perp j} < T_{\parallel j}$ ( $R_j < 1$ )	Parallel firehose (fast/whistler mode)	Oblique firehose (Alfven mode)

- Instabilities driven by  $R_j \neq 1$
- Separate modes for  $R_j > 1$  and  $< 1$
- Separate modes for  $\mathbf{k}$  parallel and oblique to  $\mathbf{B}$
- Value of  $\gamma$  strongly dependent on  $R_j$  and plasma beta:


$$\beta_{\parallel j} \equiv \frac{n_j k_B T_{\parallel j}}{B^2 / (2 \mu_0)}$$

- Example:  $\gamma_{\max}(\beta_{\parallel p}, R_p)$  for mirror instability




Maruca (PhD thesis, 2012)

# Instability Regulation of Temperature Anisotropy



- Joint distributions of  $(\beta_{\parallel j}, R_j)$ -data
  - Essentially, 2-D histograms
  - “Brazil plots”
- Popularized by Hellinger et al. (*GRL*, 2006)
  - Proton measurements *Wind*/SWE
  - Data bins smoothed into curves
  - Overlaid: contours of constant  $\gamma$  (theory)


# Instability Regulation of Temperature Anisotropy



- Alignment of data with  $\gamma$  contours
  - Excellent for oblique modes (right)
  - Worse for parallel modes (left) – despite theoretically stricter limits (especially at low- $\beta_{\parallel p}$ )
  - Cause unknown; possibly due to the shape of VDFs and the way IC resonates

# Instabilities and Heating


- Probability density of  $(\beta_{\parallel p}, R_p)$ 
  - Counts normalized by bin size
  - Bins smoothed into contours
- Temperature  $T_p$  over  $(\beta_{\parallel p}, R_p)$ -plane
  - Temperature enhancement in marginally unstable plasma
  - Heating (versus cooling) produces temperature anisotropy that drives instabilities
- Temperature components  $T_{\perp p}$  and  $T_{\parallel p}$ 
  - Enhancements in both at respective instability thresholds
  - Strongly preferential heating
  - No indications of cooling driving instabilities



Maruca et al. (PRL, 2011)

# Instabilities and Heating


- Probability density of  $(\beta_{\parallel p}, R_p)$ 
  - Counts normalized by bin size
  - Bins smoothed into contours
- Temperature  $T_p$  over  $(\beta_{\parallel p}, R_p)$ -plane
  - Temperature enhancement in marginally unstable plasma
  - Heating (versus cooling) produces temperature anisotropy that drives instabilities
- Temperature components  $T_{\perp p}$  and  $T_{\parallel p}$ 
  - Enhancements in both at respective instability thresholds
  - Strongly preferential heating
  - No indications of cooling driving instabilities



Maruca et al. (PRL, 2011)

# Temperature Anisotropy Instabilities in PIC simulation and Earth's Magnetosheath

# Instability Analysis of PIC Simulation




- 2.5-D fully kinetic PIC simulation
- Departures from LTE near current sheets (Greco et al., *PRL*, 2008; *PhRvE*, 2012)
- Linear Vlasov theory: growth rates of  $R_p \neq 1$  instabilities

- Distinct regions of  $\gamma_{\max} > 0$ 
  - More with parallel than oblique instabilities
  - Near (not co-local with) current sheets
- Turbulence generating anisotropic heating that drives instabilities?

# MMS/FPI Measurements in the Magnetosheath

- MMS/FPI burst-mode measurements of protons
  - Burst-mode cadence: 150 ms
  - 58,510 data from 6 distinct intervals
  - Intervals previously studied by Chasapis et al. (*ApJ*, 2017; *ApJL*, 2018)
  - Chosen for duration and turbulence activity
  - Four spacecraft used independently
  - For each interval, median( $R_p$ )  $\approx 1$
- Synchronization of proton and magnetic-field data
- Binning of data over  $(\beta_{\parallel p}, R_p)$ -plane




Maruca et al. (*ApJ*, 2018)


# Comparison of *MMS* Data and Linear Vlasov Theory

- Same data in both plots
- Low-count bins suppressed
- Normalization by bin size: probability density
- Contours of constant growth rate  $\gamma_{\max}$ : same code as Maruca et al. (*ApJ*, 2012)
- Very close alignment of data distribution to theoretical contours

Parallel Instabilities




Oblique Instabilities



Maruca et al. (*ApJ*, 2018)

# Instability Analysis of Time Series

- Multiple, longer periods of *MMS* data
- Growth rates of all 4 ion temperature anisotropy instabilities
- Distinct periods of  $\gamma_{\max} > 0$ 
  - Typical duration  $\approx$  few seconds
  - Similar results for parallel and oblique instabilities
  - Some alternation between  $R_p < 1$  and  $R_p > 1$  periods
- Frequency of  $\gamma_{\max} > 0$  periods varies widely



Are these instabilities important?

# Majority of solar wind is unstable ( $\sim 54\%$ )

- statistical assessment of solar wind stability at 1 AU against ion sources of free energy using Nyquist's instability criterion
- Considered multiple sources of free energy
- Less than 10% of the spectra have growth rates faster than  $\tau_{nl}$

	# Spectra	# Unstable	Mirror	CGL FH	Kinetic
Total	309	166	14	1	151
p, b, & $\alpha$	189	130	12	0	118
p & $\alpha$	114	33	2	1	30
p & b	5	3	0	0	3
p	1	0	0	0	0

Klein et al. (PRL, 2018)

# Majority of solar wind is unstable ( $\sim 54\%$ )

- statistical assessment of solar wind stability at 1 AU against ion sources of free energy using Nyquist's instability criterion
- Considered multiple sources of free energy
- Less than 10% of the spectra have growth rates faster than  $\tau_{nl}$

- $\tau_{nl}$  is an estimate for the nonlinear turbulent energy transfer time at the proton gyroscale

	# Spectra	# Unstable	Mirror	CGL FH	Kinetic
Total	309	166	14	1	151
p, b, & $\alpha$	189	130	12	0	118
p & $\alpha$	114	33	2	1	30
p & b	5	3	0	0	3
p	1	0	0	0	0

Klein et al. (PRL, 2018)

$$\tau_{nl} = (k_0 \rho_p)^{-1/3} \rho_p / V_A$$

## PIC simulation: Comparison between $\gamma$ and $\omega$

$$\tau_{\text{nl}}(r) = \ell / \delta b_\ell$$

where  $\delta b_\ell = |\hat{\ell} \cdot [\mathbf{b}(\mathbf{r} + \ell) - \mathbf{b}(\mathbf{r})]|$


$$\omega_{\text{nl}} = 2\pi / \tau_{\text{nl}}$$

# PIC simulation: Comparison between $\gamma$ and $\omega$

$$\tau_{\text{nl}}(r) = \ell / \delta b_\ell$$


where  $\delta b_\ell = |\hat{\ell} \cdot [\mathbf{b}(\mathbf{r} + \ell) - \mathbf{b}(\mathbf{r})]|$

$$\omega_{\text{nl}} = 2\pi/\tau_{\text{nl}}$$




Badyopadhyay et al. (in prep)

# Observations: Comparison between $\gamma$ and $\omega$



Badyopadhyay et al. (in prep)


## Observations: Comparison between $\gamma$ and $\omega$



Badyopadhyay et al. (in prep)


# Discussion and Conclusions

- Turbulence heats the plasma anisotropically giving rise to anisotropic distribution of instabilities in space plasma




# Discussion and Conclusions

- Turbulence heats the plasma anisotropically giving rise to anisotropic distribution of instabilities in space plasma
- The linear instability thresholds acts on plasma and restrict its excursion to extreme values of  $R_p$




# Discussion and Conclusions

- Turbulence heats the plasma anisotropically giving rise to anisotropic distribution of instabilities in space plasma
- The linear instability thresholds acts on plasma and restrict its excursion to extreme values of  $R_p$
- In simulation as well as observational data the non-linear times are almost always faster than the linear time scales



# Discussion and Conclusions

- Turbulence heats the plasma anisotropically giving rise to anisotropic distribution of instabilities in space plasma
- The linear instability thresholds acts on plasma and restrict its excursion to extreme values of  $R_p$
- In simulation as well as observational data the non-linear times are almost always faster than the linear time scales



# Questions?

Thank you!