

Intermittent distribution of ion temperature-anisotropy microinstabilities in the terrestrial magnetosheath

UD Plasma Group and MMS Team

University of Delaware

September 17, 2019

MMS-Telecon

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy \Rightarrow Non-Maxwellian VDF

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy \Rightarrow Non-Maxwellian VDF \Rightarrow Free Energy

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy \Rightarrow Non-Maxwellian VDF \Rightarrow Free Energy \Rightarrow Unstable

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy \Rightarrow Non-Maxwellian VDF \Rightarrow Free Energy \Rightarrow Unstable
 \Rightarrow Micro-Instabilities

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy \Rightarrow Non-Maxwellian VDF \Rightarrow Free Energy \Rightarrow Unstable
 \Rightarrow Micro-Instabilities \Rightarrow Growth rate (γ_j)

Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$

Anisotropy \Rightarrow Non-Maxwellian VDF \Rightarrow Free Energy \Rightarrow Unstable
 \Rightarrow Micro-Instabilities \Rightarrow Growth rate (γ_j)

	Parallel ($\mathbf{k} \parallel \mathbf{B}$) & Propagating ($\omega_r > 0$)	Oblique ($\mathbf{k} \nparallel \mathbf{B}$) & Non-Propagating ($\omega_r = 0$)
$T_{\perp j} > T_{\parallel j}$ ($R_j > 1$)	Ion-cyclotron (Alfven mode)	Mirror (kinetic slow mode)
$T_{\perp j} < T_{\parallel j}$ ($R_j < 1$)	Parallel firehose (fast/whistler mode)	Oblique firehose (Alfven mode)


Temperature Anisotropy:

$$R_j \equiv T_{\perp j} / T_{\parallel j}$$



Anisotropy \Rightarrow Non-Maxwellian VDF \Rightarrow Free Energy \Rightarrow Unstable
 \Rightarrow Micro-Instabilities \Rightarrow Growth rate (γ_j)

	Parallel ($\mathbf{k} \parallel \mathbf{B}$) & Propagating ($\omega_r > 0$)	Oblique ($\mathbf{k} \nparallel \mathbf{B}$) & Non-Propagating ($\omega_r = 0$)
$T_{\perp j} > T_{\parallel j}$ ($R_j > 1$)	Ion-cyclotron (Alfven mode)	Mirror (kinetic slow mode)
$T_{\perp j} < T_{\parallel j}$ ($R_j < 1$)	Parallel firehose (fast/whistler mode)	Oblique firehose (Alfven mode)



$$\beta_{\parallel j} \equiv \frac{n_j k_B T_{\parallel j}}{B^2 / (2 \mu_0)}$$



Hellinger et al. (*GRL*, 2006)



Maruca et al. (*ApJ*, 2018)





Marginally unstable plasma ($\gamma \gtrsim 0$) exhibits enhancements in:


- Magnetic fluctuations (Bale et al., *PRL*, 2009)
- Temperature (Maruca et al., *PRL*, 2011)
- Turbulent structures (PVI) (Osman et al., *PRL*, 2012; 2013)

Used similar method of γ calculation on a fully kinetic PIC simulation:

$$\beta_p = \beta_e = 0.6, R_p = 1, T_p = T_e$$









MMS observation in Magnetosheath


- Ion data from Fast Plasma Investigator (FPI) aboard MMS.
- In burst mode we get one proton distribution every 150 ms.
- Period analysed: Several burst modes from 2016 and 2018.
- Present results from 12/27/2016. Previously studied by Chasapis et al. (*ApJ*, 2017; *ApJL*, 2018).





- $\beta_{\parallel\rho}$ is high
- A lot more unstable VDFs
- Distribution of instabilities are still intermittent

Comparison between PIC and Observation



Conclusions

Conclusions

- Instability growth rates are intermittently distributed in the space/time

Conclusions

- Instability growth rates are intermittently distributed in the space/time
- Various studies (Osman et al. (*PRL*, 2011, 2012; Greco et al. (*PRL*, 2012) have shown intermittency to be associated with sharp gradients in magnetic field, thus implying presence of turbulence in those regions

Conclusions

- Instability growth rates are intermittently distributed in the space/time
- Various studies (Osman et al. (*PRL*, 2011, 2012; Greco et al. (*PRL*, 2012) have shown intermittency to be associated with sharp gradients in magnetic field, thus implying presence of turbulence in those regions
- Since they lie in the same configuration space, it appears that temperature anisotropy is driven by turbulence

Conclusions

- Instability growth rates are intermittently distributed in the space/time
- Various studies (Osman et al. (*PRL*, 2011, 2012; Greco et al. (*PRL*, 2012) have shown intermittency to be associated with sharp gradients in magnetic field, thus implying presence of turbulence in those regions
- Since they lie in the same configuration space, it appears that temperature anisotropy is driven by turbulence
- Locally plasma was still homogeneous enough for γ_p to limit anisotropy