

DPR: History and Physical Exam of the Gastrointestinal System Part I

Sonia Rivera-Martinez, DO, FACOFP
Associate Professor, Dept. of Family Medicine
srmartin@nyit.edu



Do.
Make.
Heal.


Innovate.
Reinvent the Future.

Session Objectives

- Identify and describe the components of a history for a common gastrointestinal complaint.
- Describe the components and sequence of the abdominal physical exam.
- Isolate and describe normal abdominal exam findings.
- Define, describe, and identify the different types of bowel sounds and their clinical significance
- Correlate abdominal exam findings with likely etiologies

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Case Presentation

- 44-year-old female with complaint of constipation which has been worsening over the past 3 to 4 months.
- What questions should you ask this patient?
- Any questions pertaining to osteopathic principles and practice?
- During your history taking consider the patient's discomfort in reporting certain complaints


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Definitions - Constipation

- Physicians:
 - Fewer than 3 bowel movements per week
- Patients:
 - Hard stools
 - Feeling of incomplete evacuation
 - Excessive straining
 - Sense of anorectal blockage during defecation
 - Need for manual maneuvers to defecate



Constipation - HPI

HPI Components	What should you ask patient?
Onset	When did it start? Older patient new-onset → cancer
Palliation/ Provocation	What makes it better/worse, What have they tried?
Quality	Lumpy or hard stools, incomplete evacuation, excessive straining. Which of these is most distressing to the patient?
Severity	Frequency, intervals between bowel movements Prolonged straining, digital evacuation → Pelvic floor dysfunction Need laxative to produce loose stools, how often and what dosage Sensation of incomplete evacuation
Timing	Constant, intermittently occurring How many bowel movements per week?

Source: Jonathan Gotfried, etal. Constipation. Merck Manual Professional, Updated April 2025
<https://www.merckmanuals.com/professional/gastrointestinal-disorders/symptoms-of-gastrointestinal-disorders/constipation?query=constipation>

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Constipation - HPI

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine

HPI Components	What should you ask patient?
Associated symptoms	Fever, nausea, abdominal pain Alternates with diarrhea → Irritable bowel syndrome Paradoxical diarrhea in elderly → fecal impaction Bloating and cramping → Irritable bowel syndrome Hematochezia → Colon cancer, diverticulosis, inflammatory bowel disease Weight loss ≥ 10 lbs → colon cancer
Other	Pain on DRE → Anal fissure, hemorrhoids Leakage of stool on DRE → Fecal impaction, rectal prolapse Medications → opioids, calcium channel blockers, iron supplements PMHX: Parkinson's, hypothyroidism, dementia

Source: Jonathan Gotfried, etal. Constipation. Merck Manual Professional, Updated April 2025
<https://www.merckmanuals.com/professional/gastrointestinal-disorders/symptoms-of-gastrointestinal-disorders/constipation?query=constipation>



Other Causes of Constipation

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Reduced physical activity
- Immobility
- Post- surgical
- Endocrine and metabolic disorders
- Medications
- Low fiber diet
- Dehydration
- Electrolyte disturbances
- Fecal impaction
- Intestinal obstruction



Source: K.Sadler, F.Arnold, Spencer D. Chronic Constipation in Adults. American Family Physician, 2022;106(3):299-306




Anatomy



NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine



Source: Bates' Guide to the Physical Examination and History Taking; 13th Ed. – Chapter 19




Abdominal Quadrant	Contents
Right Upper Quadrant (RUQ)	Liver, gallbladder, duodenum, pylorus, right kidney, hepatic flexure, ascending colon, transverse colon, lower right lung
Left Upper Quadrant (LUQ)	Tip of the medial liver lobe, spleen, stomach, left kidney, pancreas, splenic flexure, transverse colon, descending colons, left lower lung
Right Lower Quadrant (RLQ)	Appendix, cecum, ascending colon, bladder, right ovary, uterus, right spermatic cord, right ureter, small bowel
Left Lower Quadrant (LLQ)	Sigmoid colon, descending colon, bladder, left ovary, uterus, left spermatic cord, left ureter, small bowel



NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine




Abdominal Region	Contents
Right hypochondriac region	Liver, gallbladder, ascending colon, transverse colon, small intestine, hepatic flexure, right kidney
Epigastric region	Esophagus, stomach, liver, pancreas, spleen, small intestine, transverse colon, right and left (kidneys, adrenal, ureters)
Left hypochondriac region	Stomach, liver (tip), tail of pancreas, small intestine, transverse colon, descending colon, splenic flexure, left kidney, spleen
Right lumbar region	Liver (tip), gallbladder, small intestine, ascending colon, right kidney
Umbilical region	Stomach, pancreas, small intestine, transverse colon, right and left (kidneys, adrenal, ureters)



Abdominal Region	Contents
Left lumbar region	Small intestine, descending colon, left kidney (tip)
Right iliac region	Small intestine, appendix, cecum, ascending colon, right ovary and fallopian tube
Hypogastric region	Small intestine, sigmoid colon, rectum, right and left ovaries, right and left ureters, urinary bladder, uterus, right and left (ovaries and fallopian tubes), vas deferens, seminal vesicles, prostate
Left iliac region	Small intestine, descending colon, sigmoid colon, left ovary and fallopian tube




Angle formed by :
•Lower border of
12th rib
•And transverse
processes of the
upper lumbar
vertebrae



Defines region
to assess for
kidney
tenderness



Examination Techniques



1st - Inspect



2nd - Auscultate



3rd - Percuss



4th - Palpate



Abdominal Exam - Preparation

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Good Light
- Warm Room, hands, instruments
- Relaxed Patient:
 - Pillow or folded sheets under head and knees.
 - Patients arms at his/her sides
- Exposure – position drapes appropriately
- Empty Bladder
- Examine Area of Pain **Last**



Inspection



Inspect the Abdomen

- With the patient lying supine, the abdomen should be:
 - Completely exposed
 - Revealing all 4 quadrants
- Tell your patient that you are inspecting their abdomen.

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Inspect Abdominal Wall

- Contour
 - Symmetry, Masses
- Skin
 - Scars, striae
 - Dilated Veins, Discoloration
 - Rash or lesions
- Umbilicus
- Motion
- Pulsations

**Remember
to observe
from the
right side!**

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Inspect Abdominal Wall Contour

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



- Distention
 - Generalized
 - Localized
- Protuberant
- Scaphoid
- Flat




Inspect Abdomen for Masses

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Masses
 - Inspect from several angles
- Distinguish between:
 - Abdominal wall mass
 - Intra-abdominal mass



Inspect Abdomen for Scars

- Describe or diagram
- Include in the description:
 - Location
 - Size & Shape
 - Type (surgical/injury)
 - Healing stage
- Correlate with patient's history

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Inspect Abdomen for Striae

- Related to rupture of reticular dermis with rapid stretching of the skin
- Common in:
 - Cushing's syndrome
 - Obesity, Pregnancy
 - Ehlers-Danlos syndrome
 - Puberty, Ascites
 - Overuse of cortisone creams



**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




Inspect Abdomen for Dilated Veins

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- A few small visible veins may be normal.
- Abnormal dilation seen in:
 - Portal hypertension
 - Emaciation
 - IVC obstruction



Patient with caput medusa and ascites




Inspect Abdomen for Skin Lesions

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Describe
 - Location
 - Size
 - Color
 - Shape
 - Pattern and distribution



Inspect Abdomen for Discoloration

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Cullen's sign
 - Periumbilical ecchymosis
 - Hemoperitoneum
- Gray Turner's sign
 - Ecchymosis of the flank region
 - Retroperitoneal bleed



Inspect Abdomen for Jaundice

- Pre-hepatic
 - Red blood cells rupture faster than the liver can conjugate bilirubin
- Hepatic
 - Liver's inability to conjugate or excrete bilirubin
- Post-hepatic
 - Flow of bile into the intestine is blocked

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Inspect Abdominal Umbilicus

- Observe:
 - Contour
 - Location
 - Inflammation
 - Bulges



**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




Inspection of Abdomen for Respiratory Motion

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- **Absence of respiratory motion**
 - In all or part of abdominal wall suggests peritoneal irritation (acute abdomen)
- **Respiratory alternans or paradoxical**
 - Alternating outward and inward displacement - diaphragmatic muscle fatigue
 - Paradoxical inward displacement - COPD




Inspect Abdomen for Pulsations and Peristalsis

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Abdominal aortic aneurysm may be visible as a pulsatile mass
- Visible peristalsis is usually the result of intestinal obstruction



Visible peristalsis video:

<http://www.nejm.org/doi/full/10.1056/NEJMicm0910079>

Auscultation



Auscultation of the Abdomen

- Auscultate the abdomen **BEFORE** percussing or palpation as these maneuvers may alter frequency of bowel sounds.
- Place the diaphragm of your stethoscope gently on the abdomen
- Listen for bowel sounds: Note their frequency and character
- Listen for vascular bruits
- Auscultate the abdomen in **ALL 4 quadrants**

Auscultate
BEFORE
Palpation or
Percussion



Bowel Sounds - Normoactive

- Normal (normoactive):
estimated frequency of 5 to 34
“clicks or gurgles” per minute.
- Bowel sounds are produced by
peristalsis of the intestines
- Occurs intermittently:
 - Low-pitched
 - Chuckling



Bowel Sounds - Hypoactive

Reduction in the loudness, tone, or regularity of the sounds.

Sign that intestinal activity has slowed

- Normal during sleep; after the use of certain medications (opiates, anesthetics) and abdominal surgery.
- May be indicative:
 - Ileus (failure of peristalsis), Peritonitis
 - Bowel obstruction, Constipation
 - Electrolyte imbalances



Bowel Sounds - Hyperactive

- Increase in intestinal activity
- Sometimes can be heard without a stethoscope.
- Occurs with diarrhea and after eating.
- Disease states:
 - Crohn's disease, Ulcerative Colitis
 - GI bleeding, Enteritis
 - Early bowel obstruction
 - Laxative use



Bowel Sounds - Absent

- Must listen for a full 2-3 minutes before determining bowel sounds are absent
- Absence of bowels sounds occurs with:
 - Ileus, Bowel obstruction
 - Necrosis of bowel tissue
 - Peritonitis




Auscultation of Abdomen for Other Bowel Sounds

- **Borborygmi:** Prolonged gurgles of hyperperistalsis; stomach "growling" or "rumbling".
- **High-pitched tinkling sounds:** Suggestive of dilated bowel due to the presence of air and fluid under tension
- **Rushes of high-pitched** sounds that coincide with abdominal cramps: Suggestive of intestinal obstruction

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine




Auscultation of Abdomen for Vascular Bruits

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Bruits:
 - Resembles heart murmurs
 - Check aorta, renal, iliac and femoral arteries
- Presence of bruits are suggestive:
 - Renal artery stenosis (secondary cause of HTN)
 - Arterial insufficiency, Arterial occlusion




Auscultation of Abdomen for Friction Rubs

- Rare in occurrence
- May be found over liver or spleen
- Implies surface of organ is irregular
- Rub is caused by the grating of inflamed peritoneal surface against each other during respiration
- Involvement of tumor, abscess or infarction

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Percussion



Percussion of the Abdomen

- Percuss the abdomen in all 4 quadrants
- Assess for tympany, and dullness
- Usually tympany predominates
- Note any large area of dullness (may indicate mass or enlarged organ)

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




Percuss and Measure the Liver: Mid-clavicular Line

- Percuss the liver in a downward or upward direction.
- Start at an area below the umbilicus (midclavicular line) and percuss up from abdominal tympany until dullness is appreciated (lower border of the liver)
- Next, starting at the nipple line percuss from lung resonance down toward liver dullness (upper border of the liver)
- When dealing with a patient with large breasts, ask her or him to displace the breast as necessary.

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine



Normal Liver Span


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Percuss the Spleen



- Percuss the spleen on the anterior lateral, not directly on the front of the abdomen.
- Percuss the lowest interspace in the left anterior axillary line (tympanitic).
- Ask the patient to take a deep breath, and percuss again
- Should stay tympanitic
- If becomes dull upon inspiration suggests splenic enlargement



Percuss the Spleen

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Detect Abdominal Masses by Percussion

- **RUQ:** hepatomegaly, hydrops of the gallbladder, carcinoma of the head of the pancreas.
- **Epigastric:** pancreatic (pseudocyst or carcinoma), gastric malignancies, colon malignancies.
- **Flank:** kidney (cyst or tumor)
- **LUQ:** splenomegaly or carcinoma of the stomach or colon
- **RLQ:** appendiceal abscess, cecal carcinoma or ovarian tumor
- **LLQ:** diverticular abscess, sigmoid carcinoma, ovarian tumor
- **Central:** aortic aneurysms
- **Suprapubic:** distended bladder, uterine fibroid




Palpation




Palpate the Abdomen

- First, palpate the abdomen **lightly**.
 - To indentify tenderness, muscular resistance and superficial organs and masses.
- Then palpate **deeply** using finger pads, not finger tips.
 - To delineate masses (pregnant uterus, abdominal aortic aneurysm, or a distended bladder).
- **Note:** If patient complaint is “**abdominal pain**” make sure you examine the painful area **last**.




Palpate the Liver: *Bi-manual Compression*

- The left hand is placed under the right posterior ribs, and the right hand palpates the liver deeply



NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Palpate the Liver: *Two-Handed Deep Palpation*

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



- Push up towards the head from below with both hands



Palpate the Liver: Hooking Technique

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- “Hook” the liver from above with both hands



Palpate the Spleen

- Brace the posterior ribs with the left hand and palpate with the right hand
- Ask the patient to take a deep breath as the spleen is palpated



Palpation of Kidney

- Bimanual technique. Place one hand on patients back parallel to the 12th rib.
- Lift patients back.
- Place the other hand below the costal margin lateral and parallel to the rectus abdominus muscle.
- Ask patient to take a deep breath.
- At peak of inspiration attempt to capture patients kidney between your hands.

Bates' Guide to the Physical Examination and History Taking; 13th Ed. – Chapter 19

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine




Examiner stands on same side of kidney being palpated.



Palpation of Aortic Pulsations


NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Tenderness of the Abdomen

- **RUQ:** Cholecystitis, pancreatitis, or hepatitis, pulmonary pathology.
- **LUQ:** Peptic ulcer disease, splenic or pulmonary pathology
- **Epigastric:** Pancreatitis or peptic ulcer disease.
- **Periumbilical:** Appendicitis
- **Suprapubic:** Bladder/uterine pathology, cystitis, UTI
- **Flank:** Renal pathology, either pyelonephritis or perinephric abscess.
- **RLQ:** Appendicitis, cecal diverticulitis, perforated carcinoma, or ovarian pathology
- **LLQ:** Sigmoid diverticulitis or ovarian pathology



Preventative Measures – Colon Cancer

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Factors increasing risk
 - Excessive alcohol intake
 - Cigarette smoking
 - Obesity
- Factors decreasing risk
 - Regular physical activity
 - Aspirin
- USPSTF recommendations:
 - Begin screening at age 45 years and continuing until 75 years
- CDC screening tests
 - Stool tests – once a year
 - Sigmoidoscopy – every 5 years
 - Colonoscopy – every 10 years
 - CT Colonography – Every 5 years



Osteopathic Considerations – Lower bowel disorders

- Sympathetics
 - T10 -T11 – Right side of colon
 - T12 - L2 – Left side of colon
- Parasympathetics
 - Vagus – right side of colon
 - Pelvic splanchnic nerves – left side of colon
- Chapman points
 - Lateral sides of the thighs in the anterior half of the iliotibial bands



SOAP NOTE – Case Presentation

NEW YORK INSTITUTE
OF TECHNOLOGY

Subjective

cc: Constipation

HPI: 44-year-old female with complaint of constipation which has been worsening over the past 3 to 4 months. Has 1 to 2 bowel movements per week, strains to defecate, produced hard stools and many times feels she does not completely evacuate. Has tried increasing fiber and fluid in her diet, eats prunes and drink prune juice has not been helpful. Denies nausea, vomiting, diarrhea, hematochezia, abdominal pain, fever, chills, weight loss. Admits to some weight gain.

PMH: HTN, Vitiligo

PSH: BLT, C-Section x 2

Allergies: PCN (rash)

Meds: Diltiazem ER 120 mg daily, Multivitamin daily, Docusate sodium 100 mg daily

Social Hx: Denies past or present use of tobacco, alcohol or recreational drug use, 1–2 cups coffee/day, Married, sexually active with male husband, Pediatrician, eats usually a healthy diet, no stressors

Family Hx: Father & Mother A&W, no siblings, 2 daughters A&W

HCM: Mammography and pap smear this year, both normal, influenza vaccine 09/2019

ROS: Denies: fever, chills, chest pain, palpitations, SOB, cough, nausea, vomiting, diarrhea, hematochezia, melena, regurgitation, abdominal pain, joint stiffness, pain or swelling, rash, tremors, paralysis, depression or anxiety. Admits: fatigue, weight gain and skin dryness



SOAP NOTE – Case Presentation

NEW YORK INSTITUTE
OF TECHNOLOGY

Objective

VS: 100/60, HR 64, RR 12, Temp. 97.6, H 64" W 160 lbs BMI 27.5

Gen: Appears stated age, well groomed, NAD, overweight

Chest: CTA B/L, no w/r/r

Heart: S1S2, RRR, no m/r/g

Lungs: CTA B/L, no w/r/r

Abd; +BS hypoactive, soft, non-tender, non-distended, no hepatosplenomegaly, no palpable masses, no bruits, no dilated veins, no rebound or guarding, percussion sounds were tympanic in all 4 quadrants

Ext: No c/c/e, no varicosities, warm, +2 pulses x 4, joints with no synovitis, swelling or redness

Skin: No rash, good turgor, +dryness

Musc: Good range of motion upper and lower extremities

Neuro: Muscle strength 5/5 x 4, DTR's normal, negative Romberg

Rectal: No external lesions good sphincter tone, no rectal masses, +soft stool in rectal vault

OSE: L1FRS left

Labs: -Guaiac stool, UA negative

Assessment / Differential Diagnosis

Hypothyroidism

Irritable bowel syndrome

Functional constipation

Medication induced constipation

Hypertension

Overweight

Vitiligo

Plan

Labs: TSH, CBC, CMP, Guaiac stools x 3

Consider gastroenterology consult

Consider discontinuing Diltiazem once lab results are known

Continue current medications for now. Counsel patient on weight loss.

Discuss skin care. Use sun block lotions. Refer to dermatology.

Follow-up in 2 weeks.



Summary Slide

- Points:
 - **Consider** patient's discomfort with discussing their problems
 - **Observe** from the patient from the right side of the patient
 - **Explain** to the patient what you are doing
 - **Observe** the patient for signs of pain, don't rely solely on verbal cues



References

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine

- Bates Guide to Physical Examinations and History taking – 13th edition
- American Gastroenterological Association Medical Position Statement on Constipation. Gastroenterology 2013; 144:211-217
- McKean SC. Constipation. Principles and Practice of Hospital Medicine, 2017 - Chapter 80 Osteopathic considerations in systemic dysfunction, 2nd edition, 1994
- <https://www.cdc.gov/colorectal-cancer/screening/index.html>



Lecture Feedback Form: Link Below

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine



<https://comresearchdata.nyit.edu/redcap/surveys/?s=HRCY448FWYXREL4R>

